

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Commissioner
 US Department of Commerce
 United States Patent and Trademark
 Office, PCT
 2011 South Clark Place Room
 CP2/5C24
 Arlington, VA 22202
 ETATS-UNIS D'AMERIQUE
 in its capacity as elected Office

Date of mailing (day/month/year) 21 February 2001 (21.02.01)	
International application No. PCT/US00/17066	Applicant's or agent's file reference 15280-3991PC
International filing date (day/month/year) 16 June 2000 (16.06.00)	Priority date (day/month/year) 09 July 1999 (09.07.99)
Applicant SCHMIDT, Alexander, C. et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:
 10 January 2001 (10.01.01)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was
☐ was not

made before the expiration of 19-months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland	Authorized officer S. Mafla
Facsimile No.: (41-22) 740.14.35	Telephone No.: (41-22) 338.83.38

SWP

JWC/JJK cb

RECEIVED

From the
INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY

NOV 07 2001

PCT

SEATTLE
WWKMN

To:

KING, Jeffrey J. et al.
TOWNSEND AND TOWNSEND AND CREW LLP
Two Embarcadero Center, 8th Floor
San Francisco, CA 94111-3834
ETATS-UNIS D'AMERIQUENOTIFICATION OF TRANSMITTAL OF
THE INTERNATIONAL PRELIMINARY
EXAMINATION REPORT

(PCT Rule 71.1)

Date of mailing
(day/month/year) 15.10.2001Applicant's or agent's file reference
15280-399-1PC

IMPORTANT NOTIFICATION

International application No.
PCT/US00/17066 ✓International filing date (day/month/year)
16/06/2000Priority date (day/month/year)
09/07/1999 ✓

Applicant

THE GOVERNMENT OF THE UNITED STATES OF ... et al. ✓

1. The applicant is hereby notified that this International Preliminary Examining Authority transmits herewith the international preliminary examination report and its annexes, if any, established on the international application.
2. A copy of the report and its annexes, if any, is being transmitted to the International Bureau for communication to all the elected Offices.
3. Where required by any of the elected Offices, the International Bureau will prepare an English translation of the report (but not of any annexes) and will transmit such translation to those Offices.

4. REMINDER

11/9/02

The applicant must enter the national phase before each elected Office by performing certain acts (filing translations and paying national fees) within 30 months from the priority date (or later in some Offices) (Article 39(1)) (see also the reminder sent by the International Bureau with Form PCT/IB/301).

Where a translation of the international application must be furnished to an elected Office, that translation must contain a translation of any annexes to the international preliminary examination report. It is the applicant's responsibility to prepare and furnish such translation directly to each elected Office concerned.

For further details on the applicable time limits and requirements of the elected Offices, see Volume II of the PCT Applicant's Guide.

RECEIVED

RECEIVED
NOV 08 2001DOCKET DEPT.
WWKMN

NOV 08 2001

Woodcock Washburn Kurtz
Mackiewicz & Norris LLP

Name and mailing address of the IPEA/

European Patent Office
D-80298 Munich
Tel. +49 89 2399 - 0 Tx: 523656 epmu d
Fax: +49 89 2399 - 4465

Authorized officer

Hingel, W

Tel. +49 89 2399-8717



km

PATENT COOPERATION TREATY

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference 15280-3991PC	FOR FURTHER ACTION <small>see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.</small>	
International application No. PCT/US 00/17066	International filing date (day/month/year) 16/06/2000	(Earliest) Priority Date (day/month/year) 09/07/1999
Applicant THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 4 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing:

☒ contained in the international application in written form.

☒ filed together with the international application in computer readable form.

☐ furnished subsequently to this Authority in written form.

☐ furnished subsequently to this Authority in computer readable form.

☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.

☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☒ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of Invention is lacking** (see Box II).

4. With regard to the **title**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

☐ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

☒ None of the figures.

INTERNATIONAL SEARCH REPORT

International Application No

PC 00/17066

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/45 C12N5/10 C12N7/00 C07K14/115 A61K39/155

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	BAILLY J.E. ET AL.: "A recombinant human parainfluenza virus type 3 (PIV3) in which the nucleocapsid N protein has been replaced by that of bovine PIV3 is attenuated in primates." J. VIROL., vol. 74, no. 7, April 2000 (2000-04), pages 3188-3195, XP002149310 the whole document	1-83
X	WO 98 53078 A (DURBIN ANNA P ; TAO TAO (US); COLLINS PETER L (US); MURPHY BRIAN R) 26 November 1998 (1998-11-26) cited in the application the whole document see in particular p. 36, lines 15-30. --- -/--	1-83

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

5 October 2000

Date of mailing of the international search report

20/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Galli, I

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC 00/17066

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9853078 A	26-11-1998	AU 7797798 A BR 9809456 A EP 0994949 A	11-12-1998 13-06-2000 26-04-2000
WO 9314207 A	22-07-1993	AU 3340293 A CA 2126863 A EP 0621898 A FI 943211 A JP 7501707 T KR 208310 B NO 942530 A RU 2123047 C US 6033668 A US 5968776 A US 5998169 A US 6017539 A	03-08-1993 22-07-1993 02-11-1994 02-09-1994 23-02-1995 15-07-1999 05-09-1994 10-12-1998 07-03-2000 19-10-1999 07-12-1999 25-01-2000

INTERNATIONAL SEARCH REPORT

International Application No

PC 00/17066

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	VAN WYKE COELINGH K.L. ET AL.: "parainfluenza virus type 3 challenge." J. INFECT. DIS., vol. 157, no. 4, April 1988 (1988-04), pages 655-662, XP000946862 the whole document ---	1-83
A	TAO T ET AL: "A live attenuated chimeric recombinant parainfluenza virus (PIV) encoding the internal proteins of PIV type 3 and the surface glycoproteins of PIV type 1 induces complete resistance to PIV1 challenge and partial resistance to PIV3 challenge" VACCINE,GB,BUTTERWORTH SCIENTIFIC. GUILDFORD, vol. 17, no. 9-10, 5 March 1999 (1999-03-05), pages 1100-1108, XP004158231 ISSN: 0264-410X the whole document ---	1-83
A	WO 93 14207 A (CONNAUGHT LAB) 22 July 1993 (1993-07-22) the whole document -----	35-56

REC'D 22 OCT 2001

WIPO

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference 15280-399-1PC	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/US00/17066	International filing date (day/month/year) 16/06/2000	Priority date (day/month/year) 09/07/1999
International Patent Classification (IPC) or national classification and IPC C12N15/45		
Applicant THE GOVERNMENT OF THE UNITED STATES OF ... et al.		



1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2. This REPORT consists of a total of 5 sheets, including this cover sheet.

- ☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand 10/01/2001	Date of completion of this report 15.10.2001
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer Hillenbrand, G Telephone No. +49 89 2399 8428 

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/US00/17066

I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

Description, pages:

1-90 as originally filed

Claims, No.:

1-83 as originally filed

Drawings, sheets:

1-25 as originally filed

Sequence listing part of the description, pages:

1-20, as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☒ contained in the international application in written form.
- ☒ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/US00/17066

- ☐ the description, pages:
☐ the claims, Nos.:
☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):
(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

**V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability;
citations and explanations supporting such statement**

1. Statement

Novelty (N)	Yes:	Claims
	No:	Claims 1-83
Inventive step (IS)	Yes:	Claims
	No:	Claims 1-83
Industrial applicability (IA)	Yes:	Claims 1-57, 58-62 ?, 63-83
	No:	Claims

2. Citations and explanations
see separate sheet

VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:
see separate sheet

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/US00/17066

D1: WO 98 53078 A (DURBIN ANNA P ;TAO TAO (US); COLLINS PETER L (US);
MURPHY BRIAN R) 26 November 1998 (1998-11-26) cited in the application

Re Item V

Reasoned statement under Rule 66.2(a)(ii) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

Novelty (Article 33.2 PCT):

Having regard to document **D1** cited in the International Search Report the subject-matter of claims 1-83 lacks novelty.

D1 discloses inter alia infectious human-bovine chimeric PIVs (see the claims and page 36, lines 15-30 of the description). It is at present not visible which defined technical features of the claimed matter could differentiate the claimed matter from said prior art. Even if the applicant would render the claimed matter formally novel by the introduction of further additional features into the independent claims, it is not visible which surprising/advantageous effect could be involved with such a claimed matter. Thus, it appears that such matter would lack the required inventive activity.

Re Item VIII

Certain observations on the international application

For the assessment of the present claims 58-62 on the question whether they are industrially applicable, no unified criteria exist in the PCT Contracting States. The patentability can also be dependent upon the formulation of the claims. The EPO, for example, does not recognize as industrially applicable the subject-matter of claims to the use of a compound in medical treatment, but may allow, however, claims to a known compound for first use in medical treatment and the use of such a compound for the manufacture of a medicament for a new medical treatment.

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/US00/17066

The requirement that the claims shall be concise refers also to the claims in their entirety. The large number of claims on file (83!!!) makes it difficult, if not impossible, to determine the matter for which protection is sought, and places an undue burden on others seeking to establish the extent of the protection. Hence, claims 1-83 do not meet the requirements of Article 6 PCT.

In order to overcome this objection, it would appear appropriate to file an amended set of claims defining the relevant subject-matter in terms of a minimum number of independent claims in each category followed by a limited number of dependent claims covering features which are merely optional (Rule 6.4 PCT). Undue repetition of wording between one claim and another should be avoided.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 January 2001 (18.01.2001)

PCT

(10) International Publication Number
WO 01/04320 A1

(51) International Patent Classification?: C12N 15/45,
5/10, 7/00, C07K 14/115, A61K 39/155

20816 (US). BAILLY, Jane, E. [CA/CA]; 248 Braebrook
Avenue, Pointe Claire, Québec H9R 1V9 (CA). DURBIN,
Anna, P. [US/US]; 806 Hudson Avenue, Takoma Park, MD
20912 (US).

(21) International Application Number: PCT/US00/17066

(22) International Filing Date: 16 June 2000 (16.06.2000)

(74) Agents: KING, Jeffrey, J. et al.; Townsend and Townsend
and Crew LLP, 8th floor, Two Embarcadero Center, San
Francisco, CA 94111-3834 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/143,134 9 July 1999 (09.07.1999) US

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ,
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (for all designated States except US): THE
GOVERNMENT OF THE UNITED STATES OF
AMERICA, as represented by THE DEPARTMENT
OF HEALTH AND HUMAN SERVICES [US/US];
Suite 325, 6011 Executive Boulevard, Rockville, MD
20852 (US).

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SCHMIDT, Alexan-
der, C. [DE/US]; 1831 Belmont Road #203, NW, Wash-
ington, DC 20009 (US). SKIADOPOULOS, Mario, H.
[US/US]; 8303 Aqueduct Road, Potomac, MD 20854 (US).
COLLINS, Peter, L. [US/US]; Apartment #401, 12304
Village Square, Rockville, MD 20852 (US). MURPHY,
Brian, R. [US/US]; 5410 Tuscarawas Road, Bethesda, MD

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: ATTENUATED HUMAN-BOVINE CHIMERIC PARAINFLUENZA VIRUS (PIV) VACCINES

(57) Abstract: Chimeric human-bovine parainfluenza viruses (PIVs) are infectious and attenuated in humans and other mammals and useful individually or in combination in vaccine formulations for eliciting an anti-PIV immune response. Also provided are isolated polynucleotide molecules and vectors incorporating a chimeric PIV genome or antigenome which includes a partial or complete human or bovine PIV "background" genome or antigenome combined or integrated with one or more heterologous gene(s) or genome segment(s) of a different PIV. Chimeric human-bovine PIV of the invention include a partial or complete "background" PIV genome or antigenome derived from or patterned after a human or bovine PIV virus combined with one or more heterologous gene(s) or genome segment(s) of a different PIV virus to form the human-bovine chimeric PIV genome or antigenome. In certain aspects of the invention, chimeric PIV incorporate a partial or complete human PIV background genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a bovine PIV, whereby the resultant chimeric virus is attenuated by virtue of host-range restriction. In alternate embodiments, human-bovine chimeric PIV incorporate a partial or complete bovine PIV background genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a human PIV gene that encode a human PIV immunogenic protein, protein domain or epitope, for example encoded by PIV HN and/or F glycoprotein gene(s) or genome segment(s). Human-bovine chimeric PIV of the invention are also useful as vectors for developing vaccines against other pathogens. A variety of additional mutations and nucleotide modifications are provided within the human-bovine chimeric PIV of the invention to yield desired phenotypic and structural effects.

WO 01/04320 A1

ATTENUATED HUMAN-BOVINE CHIMERIC PARAINFLUENZA VIRUS (PIV) VACCINES

CROSS-REFERENCES TO RELATED APPLICATIONS

5

This application claims the benefit of United States Provisional Patent Application Serial No. 60/143,134, filed by Bailly et al. on July 9, 1999.

BACKGROUND OF THE INVENTION

10

Human parainfluenza virus type 3 (HPIV3) is a common cause of serious lower respiratory tract infection in infants and children less than one year of age. It is second only to respiratory syncytial virus (RSV) as a leading cause of hospitalization for viral lower respiratory tract disease in this age group (Collins et al., in B. N. Fields
15 Virology, p. 1205-1243, 3rd ed., vol. 1., Knipe et al., eds., Lippincott-Raven Publishers, Philadelphia, 1996; Crowe et al., Vaccine 13:415-421, 1995; Marx et al., J. Infect. Dis. 176:1423-1427, 1997, all incorporated herein by reference). Infections by this virus result in substantial morbidity in children less than 3 years of age. HPIV1 and HPIV2 are the principal etiologic agents of laryngotracheobronchitis (croup) and also can cause
20 severe pneumonia and bronchiolitis (Collins et al., 1996, *supra*). In a long term study over a 20-year period, HPIV1, HPIV2, and HPIV3 were identified as etiologic agents for 6.0, 3.2, and 11.5%, respectively, of hospitalizations for respiratory tract disease accounting in total for 18% of the hospitalizations, and, for this reason, there is a need for an effective vaccine (Murphy et al., Virus Res. 11:1-15, 1988). The parainfluenza viruses
25 have also been identified in a significant proportion of cases of virally-induced middle ear effusions in children with otitis media (Heikkinen et al., N. Engl. J. Med. 340:260-264, 1999, incorporated herein by reference). Thus, there is a need to produce a vaccine against these viruses that can prevent the serious lower respiratory tract disease and the otitis media that accompanies these HPIV infections. HPIV1, HPIV2, and HPIV3 are
30 distinct serotypes that do not elicit significant cross-protective immunity. The major protective antigens of PIVs are the hemeagglutinin (HN) and fusion (F) glycoproteins, which mediate viral attachment, penetration and release. Protection against reinfection is mediated primarily by virus-neutralizing antibodies.

Despite considerable efforts to develop effective vaccine therapies against HPIV, approved vaccine agents that prevent HPIV related illness have not yet been achieved. The most promising prospects to date are live attenuated vaccine viruses since these have been shown to be efficacious in non-human primates even in the presence of passively transferred antibodies, an experimental situation that simulates that present in the very young infant who possesses maternally acquired antibodies (Crowe et al., 1995, *supra*; and Durbin et al., J. Infect. Dis. 179:1345-1351, 1999a; each incorporated herein by reference). Two live attenuated PIV3 vaccine candidates, a temperature-sensitive (*ts*) derivative of the wild type PIV3 JS strain (designated PIV3cp45) and a bovine PIV3 (BPIV3) strain, are undergoing clinical evaluation (Karron et al., Pediatr. Infect. Dis. J. 15:650-654, 1996; Karron et al., 1995a, *supra*; Karron et al., 1995b, *supra*; each incorporated herein by reference). The BPIV3 vaccine candidate is attenuated, genetically stable and immunogenic in human infants and children. A second PIV3 vaccine candidate, JS cp45, is a cold-adapted mutant of the JS wildtype (wt) strain of HPIV3 (Karron et al., 1995b, *supra*; and Belshe et al., J. Med. Virol. 10:235-242, 1982a; each incorporated herein by reference). This live, attenuated, cold-passaged (cp) PIV3 vaccine candidate exhibits temperature-sensitive (*ts*), cold-adaptation (*ca*), and attenuation (*att*) phenotypes, which are stable after viral replication *in vivo*. The cp45 virus is protective against human PIV3 challenge in experimental animals and is attenuated, genetically stable, and immunogenic in seronegative human infants and children (Belshe et al., 1982a, *supra*; Belshe et al., Infect. Immun. 37:160-165, 1982b; Clements et al., J. Clin. Microbiol. 29:1175-1182, 1991; Crookshanks et al., J. Med. Virol. 13:243-249, 1984; Hall et al., Virus Res. 22:173-184, 1992; Karron et al., 1995b, *supra*; each incorporated herein by reference). Because these PIV3 candidate vaccine viruses are biologically derived there is no proven method for adjusting their level of attenuation as will likely be necessary for broad clinical application.

To facilitate development of PIV vaccine candidates, recombinant DNA technology has recently made it possible to recover infectious negative-strand RNA viruses from cDNA (for reviews, see Conzelmann, J. Gen. Virol. 77:381-389, 1996; Palese et al., Proc. Natl. Acad. Sci. U.S.A. 93:11354-11358, 1996; each incorporated herein by reference). In this context, rescue of recombinant viruses has been reported for infectious respiratory syncytial virus (RSV), rabies virus (RaV), simian virus 5 (SV5), rinderpest virus, Newcastle disease virus (NDV), vesicular stomatitis virus (VSV), measles virus (MeV), mumps virus (MuV) and Sendai virus (SeV) from cDNA-encoded

antigenomic RNA in the presence of essential viral proteins (see, e.g., Garcin et al., EMBO J. 14:6087-6094, 1995; Lawson et al., Proc. Natl. Acad. Sci. U.S.A. 92:4477-4481, 1995; Radecke et al., EMBO J. 14:5773-5784, 1995; Schnell et al., EMBO J. 13:4195-4203, 1994; Whelan et al., Proc. Natl. Acad. Sci. U.S.A. 92:8388-8392, 1995;

5 Hoffman et al., J. Virol. 71:4272-4277, 1997; Kato et al., Genes to Cells 1:569-579, 1996, Roberts et al., Virology 247:1-6, 1998; Baron et al., J. Virol. 71:1265-1271, 1997; International Publication No. WO 97/06270; Collins et al., Proc. Natl. Acad. Sci. USA 92:11563-11567, 1995; U.S. Patent Application No. 08/892,403, filed July 15, 1997 (corresponding to published International Application No. WO 98/02530 and priority

10 U.S. Provisional Application Nos. 60/047,634, filed May 23, 1997, 60/046,141, filed May 9, 1997, and 60/021,773, filed July 15, 1996); U.S. Patent Application Serial No. 09/291,894, filed on April 13, 1999; U.S. Provisional Patent Application Serial No. 60/129,006, filed April 13, 1999; U.S. Provisional Patent Application Serial No. 60/143,097, filed by Bucholz et al. on July 9, 1999; Juhasz et al., J. Virol. 71:5814-5819,

15 1997; He et al. Virology 237:249-260, 1997; Peters et al. J. Virol. 73:5001-5009, 1999; Whitehead et al., Virology 247:232-239, 1998a; Whitehead et al., J. Virol. 72:4467-4471, 1998b; Jin et al. Virology 251:206-214, 1998; Bucholz et al. J. Virol. 73:251-259, 1999; Whitehead et al., J. Virol. 73:3438-3442, 1999, and Clarke et al., J. Virol. 74:4831-4838, 2000; each incorporated herein by reference in its entirety for all purposes).

20 In more specific regard to the instant invention, a method for producing HPIV with a wt phenotype from cDNA was recently developed for recovery of infectious, recombinant HPIV3 JS strain (see, e.g., Durbin et al., Virology 235:323-332, 1997a; U.S. Patent Application Serial No. 09/083,793, filed May 22, 1998; U.S. Provisional Application Serial No. 60/047,575, filed May 23, 1997 (corresponding to International

25 Publication No. WO 98/53078), and U.S. Provisional Application Serial No. 60/059,385, filed September 19, 1997, each incorporated herein by reference). In addition, these disclosures allow for genetic manipulation of viral cDNA clones to determine the genetic basis of phenotypic changes in biological mutants, e.g., which mutations in the HPIV3 cp45 virus specify its ts, ca and att phenotypes, and which gene(s) or genome segment(s)

30 of BPIV3 specify its attenuation phenotype. Additionally, these and related disclosures render it feasible to construct novel PIV vaccine candidates having a wide range of different mutations and to evaluate their level of attenuation, immunogenicity and phenotypic stability (see also, U.S. Provisional Patent Application Serial No. 60/143,134,

filed by Bailly et al. on July 9, 1999; and U.S. Patent Application Serial No. 09/350,821, filed by Durbin et al. on July 9, 1999; each incorporated herein by reference).

Thus, infectious wild type recombinant PIV3 (r)PIV3, as well as a number of *ts* derivatives, have now been recovered from cDNA, and reverse genetics systems have been used to generate infectious virus bearing defined attenuating mutations and to study the genetic basis of attenuation of existing vaccine viruses. For example, the three amino acid substitutions found in the L gene of *cp45*, singularly or in combination, have been found to specify the *ts* and attenuation phenotypes. Additional *ts* and attenuating mutations are present in other regions of the PIV3 *cp45*. In addition a chimeric PIV1 vaccine candidate has been generated using the PIV3 cDNA rescue system by replacing the PIV3 HN and F open reading frames (ORFs) with those of PIV1 in a PIV3 full-length cDNA that contains the three attenuating mutations in L. The recombinant chimeric virus derived from this cDNA is designated rPIV3-1.*cp45*L (Skiadopoulos et al., J. Virol. 72:1762-1768, 1998; Tao et al., J. Virol. 72:2955-2961, 1998; Tao et al., Vaccine 17:1100-1108, 1999, incorporated herein by reference). rPIV3-1.*cp45*L was attenuated in hamsters and induced a high level of resistance to challenge with PIV1. A recombinant chimeric virus, designated rPIV3-1.*cp45*, has been produced that contains 12 of the 15 *cp45* mutations, i.e., excluding the mutations that occur in HN and F, and is highly attenuated in the upper and lower respiratory tract of hamsters (Skiadopoulos et al., Vaccine 18:503-510, 1999a).

BPIV3, which is antigenically-related to HPIV3, offers an alternative approach to the development of a live attenuated virus vaccine for HPIV1, HPIV2, and HPIV3. The first vaccine used in humans, live vaccinia virus believed to be of bovine origin, was developed by Jenner almost 200 years ago for the control of smallpox. During the ensuing two centuries, vaccinia virus was successful in controlling this disease and played an essential role in the final eradication of smallpox. In this "Jennerian" approach to vaccine development, an antigenically-related animal virus is used as a vaccine for humans. Animal viruses that are well adapted to their natural host often do not replicate efficiently in humans and hence are attenuated. At present, there is a lack of a thorough understanding regarding the genetic basis for this form of host range restriction. Evolution of a virus in its mammalian or avian host results in significant divergence of nucleotide (nt) and amino acid sequences from that of the corresponding sequences in the related human virus. This divergent sequence, consisting of a large number of sequence differences, specifies the host range attenuation phenotype. Having

an attenuation phenotype which is based on numerous sequence differences is a desirable property in a vaccine virus since it should contribute to the stability of the attenuation phenotype of the animal virus following its replication in humans.

The recently licensed quadrivalent rotavirus is an example of the Jennerian approach to vaccine development in which a nonhuman rotavirus strain, the rhesus rotavirus (RRV), was found to be attenuated in humans and protective against human serotype 3 to which it is antigenically highly related (Kapikian et al., Adv. Exp. Med. Biol. 327:59-69, 1992). Since there was a need for a multivalent vaccine that would induce resistance to each of the four major human rotavirus serotypes, the Jennerian approach was modified by constructing three reassortant viruses using conventional genetic techniques of gene reassortment in tissue culture. Each single gene reassortant virus contained 10 RRV genes plus a single human rotavirus gene that coded for the major neutralization antigen (VP7) of serotype 1, 2, or 4. The intent was to prepare single gene substitution RRV reassortants with the attenuation characteristics of this simian virus and the neutralization specificity of human rotavirus serotype 1, 2, or 4. The quadrivalent vaccine based on the host range restriction of the simian RRV in humans provided a high level of efficacy against human rotavirus infection in infants and young children (Perez-Schael et al., N. Engl. J. Med. 337:1181-1187, 1997). However, the vaccine virus retains mild reactogenicity in older seronegative infants lacking maternal antibody, therefore a second generation Jennerian vaccine, based on the UK strain of bovine rotavirus, is being developed to replace the RRV vaccine (Clements-Mann et al., Vaccine 17:2715-2725, 1999).

The Jennerian approach also is being explored to develop vaccines for parainfluenza type 1 virus and for hepatitis A virus which are attenuated and immunogenic in non-human primates (Emerson et al., J. Infect. Dis. 173:592-597, 1996; Hurwitz et al., Vaccine 15:533-540, 1997). The Jennerian approach was used for the development of a live attenuated vaccine for influenza A virus but it failed to produce a consistently attenuated vaccine for use in humans (Steinhoff et al., J. Infect. Dis. 163:1023-1028, 1991). As another example, reassortant viruses that contain two gene segments encoding the hemagglutinin and neuraminidase surface glycoproteins from a human influenza A virus and the six remaining gene segments from an avian influenza A virus were attenuated in humans (Clements et al., J. Clin. Microbiol. 27:219-222, 1989; Murphy et al., J. Infect. Dis. 152:225-229, 1985; and Snyder et al., J. Clin. Microbiol. 23:852-857, 1986). This indicated that one or more of the six gene segments of the avian

virus attenuated the avian-human influenza A viruses for humans. The genetic determinants of this attenuation were mapped using reassortant viruses possessing a single gene segment from an attenuating avian influenza A virus and the remaining genes from a human strain. It was shown that the nonstructural (NS), polymerase (PB1, PB2) and M genes contributed to the attenuation phenotype of avian influenza A viruses in humans (Clements et al., J. Clin. Microbiol. 30:655-662, 1992).

In another study, the severe host range restriction of bovine respiratory syncytial virus (BRSV) for replication in chimpanzees was only slightly alleviated by replacement of the BRSV F and G glycoproteins with their HRSV counterparts. This indicated that F and G are involved in this host range restriction, but that one or more additional bovine RSV genes are also involved (Buchholz et al., J. Virol. 74:1187-1199, 2000). This illustrates that more than one gene can contribute in unpredictable ways to the host range restriction phenotype of a mammalian or avian virus in primates.

The instant invention provides a new basis for attenuating a wild type or mutant parental virus for use as a vaccine against HPIV, in which attenuation is based completely or in part on host range effects, while at least one or more of the major neutralization and protective antigenic determinant(s) of the chimeric virus is homologous to the virus against which the vaccine is directed. The HN and F proteins of BPIV3 are each approximately 80% related by amino acid sequence to their corresponding HPIV3 proteins (Suzu et al., Nucleic Acids Res. 15:2945-2958, 1987, incorporated herein by reference) and 25% related by antigenic analysis (Coelingh et al., J. Virol. 64:3833-3843, 1990; Coelingh et al., J. Virol. 60:90-96, 1986; van Wyke Coelingh et al., J. Infect. Dis. 157:655-662, 1988, each incorporated herein by reference). Previous studies indicated that two strains of BPIV3, the Kansas (Ka) strain and the Shipping Fever (SF) prototype strain, were attenuated for the upper and lower respiratory tract of rhesus monkeys, and one of these, the Ka strain, was attenuated in chimpanzees (van Wyke Coelingh et al., 1988, *supra*, incorporated herein by reference). Immunization of nonhuman primates with the Ka virus induced antibodies reactive with HPIV3 and induced resistance to the replication of the human virus in the upper and the lower respiratory tract of monkeys (*id.*) Subsequent evaluation of the Ka strain in humans indicated that the virus was satisfactorily attenuated for seronegative infants, and it retained the attenuation phenotype following replication in fully susceptible infants and children (Karron et al., 1996, *supra*; and Karron et al., 1995a, *supra*; each incorporated herein by reference). Its major advantages therefore were that it was satisfactorily attenuated for fully susceptible

seronegative infants and children, and its attenuation phenotype was stable following replication in humans.

However, the level of serum hemagglutination-inhibiting antibodies reactive with HPIV3 induced in seronegative vaccinees who received $10^{5.0}$ tissue culture infectious dose₅₀ (TCID)₅₀ of the Ka strain of BPIV3 was 1:10.5, which was three-fold
5 lower than similar vaccinees who received a live attenuated HPIV3 vaccine (Karron et al., 1995a, *supra*; and Karron et al., 1995b, *supra*; each incorporated herein by reference). This lower level of antibodies to the human virus induced by BPIV3 reflected in large part the antigenic divergence between HPIV3 and BPIV3 (Karron et al., 1996, *supra*; and
10 Karron et al., 1995a, *supra*; each incorporated herein by reference). Studies to determine the efficacy of the Ka vaccine candidate against HPIV3 in humans have not been performed, but it is likely that this reduced level of antibodies reactive with HPIV3 will be reflected in a reduced level of protective efficacy.

Although it is clear that BPIV3 has host range genes that restrict
15 replication in the respiratory tract of rhesus monkeys, chimpanzees and humans, it remains unknown which of the bovine proteins or noncoding sequences contribute to this host range restriction of replication. It is possible that any of the BPIV3 proteins or noncoding sequences may confer a host range phenotype. It is not possible to determine in advance which genes or genome segments will confer an attenuation phenotype. This
20 can only be accomplished by systematic substitution of BPIV3 coding and non-coding sequences for their HPIV3 counterparts and by evaluation of the recovered HPIV3/BPIV3 chimeric viruses in seronegative rhesus monkeys or humans.

Despite the numerous advances toward development of effective vaccine agents against PIV serotypes 1, 2, and 3, there remains a clear need in the art for
25 additional tools and methods to engineer safe and effective vaccines to alleviate the serious health problems attributable to PIV, particularly illnesses among infants and children due to infection by HPIV. Among the remaining challenges in this context is the need for additional tools to generate suitably attenuated, immunogenic and genetically stable vaccine candidates for use in diverse clinical settings. To facilitate these goals,
30 existing methods for identifying and incorporating attenuating mutations into recombinant vaccine strains must be expanded. Furthermore, it is recognized that methods and compositions for designing vaccines against human PIV can be implemented as well to design novel vaccine candidates for veterinary use. Surprisingly,

the present invention fulfills these needs and provides additional advantages as described hereinbelow.

SUMMARY OF THE INVENTION

5

The present invention provides human-bovine chimeric parainfluenza viruses (PIVs) that are infectious and attenuated in humans and other mammals. In related aspects, the invention provides novel methods for designing and producing attenuated, human-bovine chimeric PIVs that are useful in various compositions to generate a desired immune response against PIV in a host susceptible to PIV infection. Included within these aspects of the invention are novel, isolated polynucleotide molecules and vectors incorporating such molecules that comprise a chimeric PIV genome or antigenome including a partial or complete human or bovine PIV “background” genome or antigenome combined or integrated with one or more heterologous gene(s) or genome segment(s) of a different PIV virus. Also provided within the invention are methods and compositions incorporating human-bovine chimeric PIV for prophylaxis and treatment of PIV infection.

The invention thus involves a method for developing live attenuated PIV vaccine candidates based on chimeras between HPIVs and BPIV3. Chimeras are generated using a cDNA-based virus recovery system. Recombinant viruses made from cDNA replicate independently and are propagated in the same manner as if they were biologically-derived viruses. Chimeric human-bovine PIV of the invention are recombinantly engineered to incorporate nucleotide sequences from both human and bovine PIV strains to produce an infectious, chimeric virus or subviral particle. In this manner, candidate vaccine viruses are recombinantly engineered to elicit an immune response against PIV in a mammalian host susceptible to PIV infection, including humans and non-human primates. Human-bovine chimeric PIV according to the invention may be engineered to elicit an immune response to a specific PIV, e.g., HPIV3, or a polyspecific response against multiple PIVs, e.g., HPIV1 and HPIV3. Additional chimeric viruses can be designed in accordance with the teachings herein which serve as vectors for antigens of non-PIV pathogens, for example respiratory syncytial virus (RSV) or measles virus.

Exemplary human-bovine chimeric PIV of the invention incorporate a chimeric PIV genome or antigenome comprising both human and bovine polynucleotide

sequences, as well as a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), and a large polymerase protein (L). Additional PIV proteins may be included in various combinations to provide a range of infectious subviral particles, up to a complete viral particle or a viral particle containing supernumerary proteins, antigenic determinants or other additional components.

Chimeric human-bovine PIV of the invention include a partial or complete "background" PIV genome or antigenome derived from or patterned after a human or bovine PIV strain or subgroup virus combined with one or more heterologous gene(s) or genome segment(s) of a different PIV strain or subgroup virus to form the human-bovine chimeric PIV genome or antigenome. In preferred aspects of the invention, chimeric PIV incorporate a partial or complete human PIV background genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a bovine PIV.

The partial or complete background genome or antigenome typically acts as a recipient backbone or vector into which are imported heterologous genes or genome segments of the counterpart, human or bovine PIV. Heterologous genes or genome segments from the counterpart, human or bovine PIV represent "donor" genes or polynucleotides that are combined with, or substituted within, the background genome or antigenome to yield a human-bovine chimeric PIV that exhibits novel phenotypic characteristics compared to one or both of the contributing PIVs. For example, addition or substitution of heterologous genes or genome segments within a selected recipient PIV strain may result in an increase or decrease in attenuation, growth changes, altered immunogenicity, or other desired phenotypic changes as compared with a corresponding phenotype(s) of the unmodified recipient and/or donor.

Genes and genome segments that may be selected for use as heterologous substitutions or additions within human-bovine chimeric PIV of the invention include genes or genome segments encoding a PIV N, P, C, D, V, M, F, SH (where appropriate), HN and/or L protein(s) or portion(s) thereof. In addition, genes and genome segments encoding non-PIV proteins, for example, an SH protein as found in mumps and SV5 viruses, may be incorporated within human-bovine PIV of the invention. Regulatory regions, such as the extragenic 3' leader or 5' trailer regions, and gene-start, gene-end, intergenic regions, or 3' or 5' non-coding regions, are also useful as heterologous substitutions or additions.

Preferred human-bovine chimeric PIV vaccine candidates of the invention

bear one or more of the major antigenic determinants of HPIV3 in a background which is attenuated by the substitution or addition of one or more BPIV3 genes or genome segments. The major protective antigens of PIVs are their HN and F glycoproteins, although other proteins can also contribute to a protective immune response. In certain
5 embodiments, the background genome or antigenome is an HPIV genome or antigenome, e.g., an HPIV3, HPIV2, or HPIV1 background genome or antigenome, to which is added or into which is substituted one or more BPIV gene(s) or genome segment(s), preferably from BPIV3. In one exemplary embodiment described below, an ORF of the N gene of a
10 antigenome may be a BPIV genome or antigenome which is combined with one or more genes or genome segments encoding a HPIV3, HPIV2, or HPIV1 glycoprotein, glycoprotein domain or other antigenic determinant.

In accordance with the methods of the invention, any BPIV gene or genome segment, singly or in combination with one or more other BPIV genes, can be
15 combined with HPIV sequences to give rise to a human-bovine chimeric PIV vaccine candidate. Any HPIV, including different strains of a particular HPIV serotype, e.g., HPIV3 will be a reasonable acceptor for attenuating BPIV gene(s). In general, the HPIV3 gene(s) or genome segment(s) selected for inclusion in a human-bovine chimeric PIV for use as a vaccine against human PIV will include one or more of the HPIV protective
20 antigens such as the HN or F glycoproteins.

In exemplary aspects of the invention, human-bovine chimeric PIV bearing one or more bovine gene(s) or genome segment(s) exhibits a high degree of host range restriction, e.g., in the respiratory tract of mammalian models of human PIV infection such as non-human primates. In exemplary embodiments a human PIV is
25 attenuated by the addition or substitution of one or more bovine gene(s) or genome segment(s) to a partial or complete human, e.g., HPIV3, PIV background genome or antigenome. In one example, the HPIV3 N gene is substituted by the BPIV3 N gene to yield a novel human-bovine chimeric PIV vaccine candidate.

Preferably, the degree of host range restriction exhibited by human-bovine
30 chimeric PIV vaccine candidates of the invention is comparable to the degree of host range restriction exhibited by the respective BPIV parent or "donor" strain. Preferably, the restriction should have a true host range phenotype, i.e., it should be specific to the host in question and should not restrict replication and vaccine preparation *in vitro* in a suitable cell line. In addition, human-bovine chimeric PIV bearing one or more bovine

gene(s) or genome segment(s) elicit a high level of resistance in hosts susceptible to PIV infection. Thus, the invention provides a new basis for attenuating a live virus vaccine against PIV, one which is based on host range effects due to the introduction of one or more gene(s) or genome segment(s) from a heterologous PIV, e.g., between HPIV3 and
5 BPIV3.

In related aspects of the invention, human-bovine chimeric PIV incorporates one or more heterologous gene(s) that encode an HPIV HN and/or F glycoprotein(s). Alternatively, the chimeric PIV may incorporate one or more genome segment(s) encoding an ectodomain (and alternatively a cytoplasmic domain and/or
10 transmembrane domain), or immunogenic epitope of an HPIV HN and/or F glycoprotein(s). These immunogenic proteins, domains and epitopes are particularly useful within human-bovine chimeric PIV because they generate novel immune responses in an immunized host. In particular, the HN and F proteins, and immunogenic domains and epitopes therein, provide major protective antigens.

In certain embodiments of the invention, addition or substitution of one or more immunogenic gene(s) or genome segment(s) from a human PIV subgroup or strain to or within a bovine background, or recipient, genome or antigenome yields a recombinant, chimeric virus or subviral particle capable of generating an immune response directed against the human donor virus, including one or more specific human
20 PIV subgroups or strains, while the bovine backbone confers an attenuated phenotype making the chimera a useful candidate for vaccine development. In one exemplary embodiment, one or more human PIV glycoprotein genes, e.g., HN and/or F, are added to or substituted within a partial or complete bovine genome or antigenome to yield an attenuated, infectious human-bovine chimera that elicits an anti-human PIV immune
25 response in a susceptible host.

In alternate embodiments, human-bovine chimeric PIV additionally incorporate a gene or genome segment encoding an immunogenic protein, protein domain or epitope from multiple human PIV strains, for example two HN or F proteins or immunogenic portions thereof each from a different HPIV, e.g., HPIV1 or HPIV2.
30 Alternatively, one glycoprotein or immunogenic determinant may be provided from a first HPIV, and a second glycoprotein or immunogenic determinant may be provided from a second HPIV by substitution without the addition of an extra glycoprotein- or determinant- encoding polynucleotide to the genome or antigenome. Substitution or addition of HPIV glycoproteins and antigenic determinants may also be achieved by

construction of a genome or antigenome that encodes a chimeric glycoprotein in the recombinant virus or subviral particle, for example having an immunogenic epitope, antigenic region or complete ectodomain of a first HPIV fused to a cytoplasmic domain of a heterologous HPIV. For example, a heterologous genome segment encoding a
5 glycoprotein ectodomain from a HPIV1 or HPIV2 HN or F glycoprotein may be joined with a genome segment encoding a corresponding HPIV3 HN or F glycoprotein cytoplasmic/endodomain in the background genome or antigenome.

In alternate embodiments a human-bovine chimeric PIV genome or antigenome may encode a substitute, extra, or chimeric glycoprotein or antigenic
10 determinant thereof in the recombinant virus or subviral particle, to yield a viral recombinant having both human and bovine glycoproteins, glycoprotein domains, or immunogenic epitopes. For example, a heterologous genome segment encoding a glycoprotein ectodomain from a human PIV HN or F glycoprotein may be joined with a genome segment encoding a corresponding bovine HN or F glycoprotein
15 cytoplasmic/endodomain in the background genome or antigenome. Alternatively, the human PIV HN or F glycoprotein or parts thereof may be joined with a genome segment encoding an HN or F glycoprotein or parts thereof from another PIV strain or serotype.

Thus, according to the methods of the invention, human-bovine chimeric PIV may be constructed by substituting the heterologous gene or genome segment for a
20 counterpart gene or genome segment in a partial PIV background genome or antigenome. Alternatively, the heterologous gene or genome segment may be added as a supernumerary gene or genome segment in combination with a complete (or partial if another gene or genome segment is deleted) PIV background genome or antigenome. For example, two human PIV HN or F genes or genome segments can be included, one each
25 from HPIV2 and HPIV3.

Often, a heterologous gene or genome segment is added near an intergenic position within a partial or complete PIV background genome or antigenome. Alternatively, the gene or genome segment can be placed in other noncoding regions of the genome, for example, within the 5' or 3' noncoding regions or in other positions
30 where noncoding nucleotides occur within the partial or complete genome or antigenome. In one aspect, noncoding regulatory regions contain cis-acting signals required for efficient replication, transcription, and translation, and therefore represent target sites for modification of these functions by introducing a heterologous gene or genome segment or other mutation as disclosed herein.

In more detailed aspects of the invention, attenuating mutations are introduced into cis-acting regulatory regions to yield, e.g., (1) a tissue specific attenuation (Gromeier et al., J. Virol. 73:958-964, 1999; Zimmermann et al., J. Virol. 71:4145-4149, 1997), (2) increased sensitivity to interferon (Zimmermann et al., 1997, *supra*), (3) 5 temperature sensitivity (Whitehead et al., 1998a, *supra*), (4) a general restriction in level of replication (Men et al., J. Virol. 70:3930-3937, 1996; Muster et al., Proc. Natl. Acad. Sci. USA 88:5177-5181, 1991), and/or (5) host specific restriction of replication (Cahour et al., Virology 207:68-76, 1995). These attenuating mutations can be achieved in various ways to produce an attenuated human-bovine chimeric PIV of the invention, for 10 example by point mutations, swaps of sequences between related viruses, or nucleotide deletions.

In yet additional alternative methods provided herein, a heterologous gene or genome segment may be added or substituted at a position corresponding to a wild-type gene order position of a counterpart gene or genome segment within the partial or 15 complete PIV background genome or antigenome. In other embodiments, the heterologous gene or genome segment is added or substituted at a position that is more promoter-proximal or promoter-distal compared to a wild-type gene order position of a counterpart gene or genome segment within the background genome or antigenome, to enhance or reduce expression, respectively, of the heterologous gene or genome segment.

20 In general aspects of the invention, bovine genes or genome segments may be added to or substituted within a human PIV background to form an attenuated, human-bovine chimeric PIV. Alternatively, the chimera may be comprised of one or more human gene(s) or genome segment(s) added to or substituted within a bovine PIV background to form an attenuated PIV vaccine candidate. In this context, a chimeric PIV 25 genome or antigenome is formed of a partial or complete bovine PIV background genome or antigenome combined with a heterologous gene or genome segment from a human PIV. In preferred aspects, one or more bovine PIV gene(s) or genome segment(s) is substituted for a counterpart gene(s) or genome segment(s) within a human PIV background genome or antigenome. In alternate embodiments, one or more human PIV 30 glycoprotein genes, e.g., HN and/or F or a genome segment encoding a cytoplasmic domain, transmembrane domain, ectodomain or immunogenic epitope of a human PIV glycoprotein gene is substituted for a counterpart gene or genome segment within the bovine PIV background genome or antigenome. For example, both human PIV

glycoprotein genes HN and F may be substituted to replace counterpart HN and F glycoprotein genes in a bovine PIV background genome or antigenome.

In a parallel fashion, the chimeric human-bovine PIV of the invention can be readily designed as "vectors" to incorporate antigenic determinants from different pathogens, including more than one PIV strain or group (e.g., both human PIV3 and human PIV1), respiratory syncytial virus (RSV), measles and other pathogens (see, e.g., U.S. Provisional Patent Application Serial No. 60/170,195, filed December 10, 1999 by Murphy et al., incorporated herein by reference).

In more detailed aspects of the invention, human-bovine chimeric PIV are comprised of a partial or complete BPIV background genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a human PIV. Within these aspects, one or more of the HPIV glycoprotein genes HN and F, or one or more genome segments encoding a cytoplasmic domain, transmembrane domain, ectodomain or immunogenic epitope of the HN and/or F genes, may be added to a BPIV background genome or antigenome or substituted for one or more counterpart genes or genome segments within the BPIV background genome or antigenome to yield the chimeric construct. Often, both HPIV glycoprotein genes HN and F will be substituted to replace counterpart HN and F glycoprotein genes in the BPIV background genome or antigenome, as exemplified by the recombinant chimeric virus rBPIV3-F_HHN_H described below. This is a desirable construct because it combines the antigenic determinants of the human PIV with the host range restricting elements of the bovine PIV.

In combination with the host range phenotypic effects provided in the human-bovine chimeric PIV of the invention, it is often desirable to adjust the attenuation phenotype by introducing additional mutations that increase or decrease attenuation of the chimeric virus. Thus, in additional aspects of the invention, attenuated, human-bovine chimeric PIV are produced in which the chimeric genome or antigenome is further modified by introducing one or more attenuating mutations specifying an attenuating phenotype in the resultant virus or subviral particle. These can include mutations in RNA regulatory sequences or in encoded proteins. These attenuating mutations may be generated *de novo* and tested for attenuating effects according to a rational design mutagenesis strategy. Alternatively, the attenuating mutations may be identified in existing biologically derived mutant PIV and thereafter incorporated into a human-bovine chimeric PIV of the invention.

Introduction of attenuating and other desired phenotype-specifying mutations into chimeric bovine-human PIV of the invention may be achieved by transferring a heterologous gene or genome segment, e.g., a gene encoding an L protein or portion thereof, into a bovine or human PIV background genome or antigenome.

5 Alternatively, the mutation may be present in the selected background genome or antigenome, and the introduced heterologous gene or genome segment may bear no mutations or may bear one or more different mutations. Typically, the human or bovine background or "recipient" genome or antigenome is modified at one or more sites corresponding to a site of mutation in a heterologous virus (e.g., a heterologous bovine or
10 human PIV or a non-PIV negative stranded RNA virus) to contain or encode the same or a conservatively related mutation (e.g., a conservative amino acid substitution) as that identified in the donor virus (see, PCT/US00/09695 filed April 12, 2000 and its priority U.S. Provisional Patent Application Serial No. 60/129,006, filed April 13, 1999, incorporated herein by reference). In one exemplary embodiment, a bovine background
15 or "recipient" genome or antigenome is modified at one or more sites corresponding to a site of mutation in HPIV3 JS *cp45*, as enumerated below, to contain or encode the same or a conservatively related mutation as that identified in the *cp45* "donor."

Preferred mutant PIV strains for identifying and incorporating attenuating mutations into bovine-human chimeric PIV of the invention include cold passaged (cp),
20 cold adapted (ca), host range restricted (hr), small plaque (sp), and/or temperature sensitive (ts) mutants, for example the JS HPIV3 cp 45 mutant strain. In exemplary embodiments, one or more attenuating mutations occur in the polymerase L protein, e.g., at a position corresponding to Tyr₉₄₂, Leu₉₉₂, or Thr₁₅₅₈ of JS wild type HPIV3. Alternatively, attenuating mutations in the N protein may be selected and incorporated in
25 a human-bovine chimeric PIV, for example which encode amino acid substitution(s) at a position corresponding to residues Val₉₆ or Ser₃₈₉ of JS. Alternative or additional mutations may encode amino acid substitution(s) in the C protein, e.g., at a position corresponding to Ile₉₆ of JS and in the M protein, e.g., at a position corresponding to Pro₁₉₉ (for example a Pro₁₉₉ to Thr mutation). Yet additional mutations for adjusting
30 attenuation of a human-bovine chimeric PIV of the invention are found in the F protein, e.g., at a position corresponding to Ile₄₂₀ or Ala₄₅₀ of JS, and in the HN protein, e.g., at a position corresponding to residue Val₃₈₄ of JS.

Attenuating mutations from biologically derived PIV mutants for incorporation into human-bovine chimeric PIV of the invention also include mutations in

noncoding portions of the PIV genome or antigenome, for example in a 3' leader sequence. Exemplary mutations in this context may be engineered at a position in the 3' leader of a recombinant virus at a position corresponding to nucleotide 23, 24, 28, or 45 of JS cp45. Yet additional exemplary mutations may be engineered in the N gene start sequence, for example by changing one or more nucleotides in the N gene start sequence, e.g., at a position corresponding to nucleotide 62 of JS cp45.

From JS cp45 and other biologically derived PIV and non-PIV mutants, a large "menu" of attenuating mutations is provided, each of which mutations can be combined with any other mutation(s) for adjusting the level of attenuation in a recombinant PIV bearing a genome or antigenome that is a chimera of human and bovine gene(s) or genome segment(s). For example, mutations within recombinant PIV of the invention include one or more, and preferably two or more, mutations of JS cp45. Desired human-bovine chimeric PIV of the invention selected for vaccine use often have at least two and sometimes three or more attenuating mutations to achieve a satisfactory level of attenuation for broad clinical use. Preferably, recombinant human-bovine chimeric PIV incorporate one or more attenuating mutation(s) stabilized by multiple nucleotide substitutions in a codon specifying the mutation.

Additional mutations which can be adopted or transferred to human-bovine chimeric PIV of the invention may be identified in non-PIV nonsegmented negative stranded RNA viruses and incorporated in PIV mutants of the invention. This is readily accomplished by mapping the mutation identified in a heterologous negative stranded RNA virus to a corresponding, homologous site in a recipient PIV genome or antigenome and mutating the existing sequence in the recipient to the mutant genotype (either by an identical or conservative mutation), as described in PCT/US00/09695 filed April 12, 2000 and its priority U.S. Provisional Patent Application Serial No. 60/129,006, filed April 13, 1999, incorporated herein by reference.

In addition to recombinant human-bovine chimeric PIV, the invention provides related cDNA clones, vectors and particles, each of which incorporate HPIV and BPIV sequences and, optionally, one or more of the additional, phenotype-specific mutations set forth herein. These are introduced in selected combinations, e.g., into an isolated polynucleotide which is a recombinant cDNA genome or antigenome, to produce a suitably attenuated, infectious virus or subviral particle upon expression, according to the methods described herein. This process, coupled with routine phenotypic evaluation, provides human-bovine chimeric PIV having such desired characteristics as attenuation,

temperature sensitivity, altered immunogenicity, cold-adaptation, small plaque size, host range restriction, genetic stability, etc. In particular, vaccine candidates are selected which are attenuated and yet are sufficiently immunogenic to elicit a protective immune response in the vaccinated mammalian host.

5 In yet additional aspects of the invention, human-bovine chimeric PIV, with or without additional mutations adopted, e.g., from a biologically derived attenuated mutant virus, are constructed to have additional nucleotide modification(s) to yield a desired phenotypic, structural, or functional change. Typically, the selected nucleotide modification will specify a phenotypic change, for example a change in growth
10 characteristics, attenuation, temperature-sensitivity, cold-adaptation, plaque size, host range restriction, or immunogenicity. Structural changes in this context include introduction or ablation of restriction sites into PIV encoding cDNAs for ease of manipulation and identification.

In preferred embodiments, nucleotide changes within the genome or
15 antigenome of a human-bovine chimeric PIV include modification of a viral gene by partial or complete deletion of the gene or reduction or ablation (knock-out) of its expression. These modifications can be introduced within the human or bovine background genome or antigenome, or may be introduced into the chimeric genome or antigenome by incorporation within the heterologous gene(s) or genome segment(s)
20 added or substituted therein. Target genes for mutation in this context include any of the PIV genes, including the nucleocapsid protein N, phosphoprotein P, large polymerase subunit L, matrix protein M, hemagglutinin-neuraminidase protein HN, small hydrophobic SH protein, where applicable, fusion protein F, and the products of the C, D and V open reading frames (ORFs). To the extent that the recombinant virus remains
25 viable and infectious, each of these proteins can be selectively deleted, substituted or rearranged, in whole or in part, alone or in combination with other desired modifications, to achieve novel deletion or knock out mutants. For example, one or more of the C, D, and/or V genes may be deleted in whole or in part, or its expression reduced or ablated (e.g., by introduction of a stop codon, by a mutation in an RNA editing site, by a mutation
30 that alters the amino acid specified by an initiation codon, or by a frame shift mutation in the targeted ORF(s). In one embodiment, a mutation can be made in the editing site that prevents editing and ablates expression of proteins whose mRNA is generated by RNA editing (Kato et al., *EMBO J.* 16:578-587, 1997a and Schneider et al., *Virology* 227:314-322, 1997, each incorporated herein by reference). Alternatively, one or more of the C,

D, and/or V ORF(s) can be deleted in whole or in part to alter the phenotype of the resultant recombinant clone to improve growth, attenuation, immunogenicity or other desired phenotypic characteristics (see, U.S. Patent Application Serial No. 09/350,821, filed by Durbin et al. on July 9, 1999, incorporated herein by reference).

5 Alternative nucleotide modifications in human-bovine chimeric PIV of the invention include a deletion, insertion, addition or rearrangement of a cis-acting regulatory sequence for a selected gene in the recombinant genome or antigenome. As with other such modifications described herein, these modifications can be introduced within the human or bovine background genome or antigenome, or may be introduced
10 into the chimeric genome or antigenome by incorporation within the heterologous gene(s) or genome segment(s) added or substituted therein. In one example, a cis-acting regulatory sequence of one PIV gene is changed to correspond to a heterologous regulatory sequence, which may be a counterpart cis-acting regulatory sequence of the same gene in a different PIV, or a cis-acting regulatory sequence of a different PIV gene.
15 For example, a gene end signal may be modified by conversion or substitution to a gene end signal of a different gene in the same PIV strain. In other embodiments, the nucleotide modification may comprise an insertion, deletion, substitution, or rearrangement of a translational start site within the recombinant genome or antigenome, e.g., to ablate an alternative translational start site for a selected form of a protein.

20 In addition, a variety of other genetic alterations can be produced in a human-bovine chimeric PIV genome or antigenome, alone or together with one or more attenuating mutations adopted from a biologically derived mutant PIV. For example, genes or genome segments from non-PIV sources may be inserted in whole or in part. Alternatively, the order of genes can be changed, or a PIV genome promoter replaced
25 with its antigenome counterpart. Different or additional modifications in the recombinant genome or antigenome can be made to facilitate manipulations, such as the insertion of unique restriction sites in various non-coding regions or elsewhere. Nontranslated gene sequences can be removed to increase capacity for inserting foreign sequences.

30 In yet additional aspects, polynucleotide molecules or vectors encoding the human-bovine chimeric PIV genome or antigenome can be modified to encode non-PIV sequences, e.g., a cytokine, a T-helper epitope, a restriction site marker, or a protein or immunogenic epitope of a microbial pathogen (e.g., virus, bacterium, parasite, or fungus) capable of eliciting a protective immune response in an intended host. In one such embodiment, human-bovine chimeric PIV are constructed that incorporate a gene or

genome segment from a respiratory syncytial virus (RSV), for-example a gene encoding an antigenic protein (e.g., an F or G protein), immunogenic domain or epitope of RSV.

In related aspects of the invention, compositions (e.g., isolated polynucleotides and vectors incorporating a PIV-encoding cDNA) and methods are provided for producing an isolated infectious human-bovine chimeric PIV. Included within these aspects of the invention are novel, isolated polynucleotide molecules and vectors incorporating such molecules that comprise a human-bovine chimeric PIV genome or antigenome. Also provided is the same or different expression vector comprising one or more isolated polynucleotide molecules encoding N, P, and L proteins. These proteins also can be expressed directly from the genome or antigenome cDNA. The vector(s) is/are preferably expressed or coexpressed in a cell or cell-free lysate, thereby producing an infectious human-bovine chimeric PIV viral particle or subviral particle.

The above methods and compositions for producing human-bovine chimeric PIV yield infectious viral or subviral particles, or derivatives thereof. A recombinant infectious virus is comparable to the authentic PIV virus particle and is infectious as is. It can directly infect fresh cells. An infectious subviral particle typically is a subcomponent of the virus particle which can initiate an infection under appropriate conditions. For example, a nucleocapsid containing the genomic or antigenomic RNA and the N, P, and L proteins is an example of a subviral particle which can initiate an infection if introduced into the cytoplasm of cells. Subviral particles provided within the invention include viral particles which lack one or more protein(s), protein segment(s), or other viral component(s) not essential for infectivity.

In other embodiments the invention provides a cell or cell-free lysate containing an expression vector which comprises an isolated polynucleotide molecule comprising a human-bovine chimeric PIV genome or antigenome as described above, and an expression vector (the same or different vector) which comprises one or more isolated polynucleotide molecules encoding the N, P, and L proteins of PIV. One or more of these proteins also can be expressed from the genome or antigenome cDNA. Upon expression the genome or antigenome and N, P and L combine to produce an infectious human-bovine chimeric PIV virus or subviral particle.

The human-bovine chimeric PIVs of the invention are useful in various compositions to generate a desired immune response against PIV in a host susceptible to PIV infection. Human-bovine chimeric PIV recombinants are capable of eliciting a

protective immune response in an infected mammalian host, yet are sufficiently attenuated so as not to cause unacceptable symptoms of severe respiratory disease in the immunized host. In addition, the human-bovine chimeric PIV recombinants should replicate with sufficient efficiency *in vitro* to make vaccine preparation feasible. The
5 attenuated virus or subviral particle may be present in a cell culture supernatant, isolated from the culture, or partially or completely purified. The virus may also be lyophilized, and can be combined with a variety of other components for storage or delivery to a host, as desired.

The invention further provides novel vaccines comprising a
10 physiologically acceptable carrier and/or adjuvant and an isolated attenuated human-bovine chimeric PIV virus or subviral particle. In preferred embodiments, the vaccine is comprised of a human-bovine chimeric PIV having at least one, and preferably two or more additional mutations or other nucleotide modifications as described above to achieve a suitable balance of attenuation and immunogenicity. The vaccine can be
15 formulated in a dose of 10^3 to 10^7 PFU of attenuated virus. The vaccine may comprise attenuated human-bovine chimeric PIV that elicits an immune response against a single PIV strain or against multiple PIV strains or groups. In this regard, human-bovine chimeric PIV can be combined in vaccine formulations with other PIV vaccine strains, or with other viral vaccine viruses such as RSV.

20 In related aspects, the invention provides a method for stimulating the immune system to elicit an immune response against PIV in a mammalian subject. The method comprises administering a formulation of an immunologically sufficient amount of a human-bovine chimeric PIV in a physiologically acceptable carrier and/or adjuvant. In one embodiment, the immunogenic composition is a vaccine comprised of a human-
25 bovine chimeric PIV having at least one, and preferably two or more attenuating mutations or other nucleotide modifications specifying a desired phenotype as described above. The vaccine can be formulated in a dose of 10^3 to 10^7 PFU of attenuated virus. The vaccine may comprise attenuated human-bovine chimeric PIV virus that elicits an immune response against a single PIV, against multiple PIVs, e.g., HPIV1 and HPIV3, or
30 against one or more PIV(s) and a non-PIV pathogen such as RSV. In this context, human-bovine chimeric PIV can elicit a monospecific immune response or a polyspecific immune response against multiple PIVs, or against one or more PIV(s) and a non-PIV pathogen such as RSV. Alternatively, human-bovine chimeric PIV having different immunogenic characteristics can be combined in a vaccine mixture or administered

separately in a coordinated treatment protocol to elicit more effective protection against one PIV, against multiple PIVs, or against one or more PIV(s) and a non-PIV pathogen such as RSV. Preferably the immunogenic composition is administered to the upper respiratory tract, e.g., by spray, droplet or aerosol.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates cloning of the N coding region of bovine PIV strains Ka or SF into HPIV3. In Fig. 1A-C, the BPIV3 N open reading frame (ORF) replaces its
 10 corresponding HPIV3 sequence in the full-length rJS antigenomic cDNA (Durbin et al., 1997a, *supra*). BPIV3 Ka and SF N genes were first amplified by RT-PCR using standard molecular biological techniques from virion RNA and subcloned as 1.9 kb fragments into pBluescript to give pBS-KaN or pBS-SFN, respectively. The HPIV3 rJS N gene was subcloned as a 1.9 kb MluI/EcoRI fragment into pUC 119 from a plasmid
 15 containing the 5' half of the rJS HPIV3 antigenome (Durbin et al., 1997a, *supra*; U.S. Patent Application Serial No. 09/083,793, filed May 22, 1998; U.S. Provisional Application No. 60/047,575, filed May 23, 1997 (corresponding to International Publication No. WO 98/53078), and U.S. Provisional Application No. 60/059,385, filed September 19, 1997, each incorporated herein by reference) to give pUC119JSN. Each N
 20 gene was modified by site-directed mutagenesis to place an NcoI and AflII site at the translational start and stop sites, respectively. The Ka and SF N genes are identical in the translational start and stop site regions and, therefore, identical mutagenesis reactions were performed on both BPIV3 N genes as depicted in 1A. Fig. 1B--Following AflII/NcoI digestion, a 1.5 kb fragment from pBS-KaN or pBS-SFN representing the
 25 BPIV3 N coding region was introduced into the NcoI/AflII window of the HPIV3 N subclone pUC119JSN-NcoI/AflII as a replacement for its HPIV3 counterpart. Fig. 1C--Each chimeric subclone was then subjected to site-directed mutagenesis to restore the sequence present in HPIV3 rJS before the translation start codon or after the stop codon and BPIV3 coding sequence immediately after the start codon and before the stop codon.
 30 This yielded pUC119B/HKaN and pUC119B/HSFN, which were used to import the BPIV3 N gene into the HPIV3 cDNA clone as shown in Fig. 2. Figure 1, Panel A, CAAAAATGTTG (SEQ ID NO. 10); GCAACTAATCGA (SEQ ID NO. 11); TAACCATGGTGA (SEQ ID NO. 12); GCACTTAAGCAC (SEQ ID NO. 13). Figure 1,

Panel C, TAACCATGGTGA (SEQ ID NO. 12); GCACTTAAGCAC (SEQ ID NO. 13); CAAAAATGTTGA (SEQ ID NO. 14); GCAACTAGTCGA (SEQ ID NO. 15).

Figure 2 illustrates insertion of the HPIV3/BPIV3 (strain Ka or SF) chimeric N gene into the HPIV3 antigenomic cDNA. In Fig. 2A, the BPIV3 N ORF of Ka or SF flanked by HPIV3 sequence was subcloned as an MluI/EcoRI fragment from pUC119B/HKaN or pUC119B/HSFN and inserted into pLeft+2G (Durbin et al., 1997a, *supra*). The pLeft+2G plasmid contains the 5' half of the HPIV3 rJS antigenome from nt 1-7437 (genome sense) behind a T7 promoter. The location of two G residues that were inserted between T7 promoter and HPIV3 sequence to improve transcription is indicated by an asterisk. Fig. 2B--An XhoI/NgoMl fragment of pRight (Durbin et al., 1997a, *supra*; U.S. Patent Application Serial No. 09/083,793, filed May 22, 1998; U.S. Provisional Application No. 60/047,575, filed May 23, 1997 (corresponding to International Publication No. WO 98/53078), and U.S. Provisional Application No. 60/059,385, filed September 19, 1997, each incorporated herein by reference) containing the 3' end of the HPIV3 antigenome flanked by the hepatitis delta virus ribozyme and T7 terminator was cloned into the XhoI/NgoMl window of the modified pLeft plasmid resulting in plasmids pB/HPIV3KaN and pB/HPIV3SFN. Each of these chimeric constructs contains the complete positive sense sequence of the HPIV3 antigenomic RNA except for the N coding region which has been replaced by its BPIV3 Ka or SF counterpart.

Figure 3 provides nucleotide sequences of HPIV3, BPIV3 and chimeric viruses of the invention around N translation start (A) and stop (B) codons. The position of the individual ORFs is described in the respective Genbank reports (#AF178654 for BPIV3 Ka, #AF178655 for BPIV3 SF and #Z11515) and included herein by reference. The sequences (positive-sense) flanking the translational start (A) and stop (B) codons (each underlined) in the N gene are shown for the parental recombinant HPIV3 JS (rJS), the parental biologically-derived BPIV3 Ka and SF viruses (Ka and SF), and the chimeric cKa and cSF viruses. Host-specific residues in the cKa and cSF virus sequences and their counterparts in rJS (before the start codon and after the stop codon) and SF or Ka (start codon through stop codon, inclusive) are in boldface type. Plaque-purified chimeric virus was amplified by RT-PCR from virion RNA and sequenced using the Taq Dye Deoxy Terminator Cycle kit (ABI, Foster City, CA). This confirmed that the predicted sequences were present in each chimeric virus. Figure 3A. rJS, GGAAGCTCTATAATTTCAAAAATGTTGAGCCTATTTGATAC (SEQ ID NO. 16). Figure 3A. cKa and cSF,

GGAAGCTCTATAATTTCAAAAATGTTGAGTCTATTTCGACAC (SEQ ID NO. 17).

Figure 3A. Ka and SF,

GAAATCCTAAGACTGTAATCATGTTGAGTCTATTTCGACAC (SEQ ID NO. 18).

Figure 3B. rJS, TTAACGCATTTGGAAGCAACTAATCGAATCAACATTTTAA (SEQ

5 ID NO. 19). Figure 3B. cKa and cSF,

TCAGTGCATTCGGAAGCAACTAGTCGAATCAACATTTTAA (SEQ ID NO. 20).

Figure 3B. Ka and SF,

TCAGTGCATTCGGAAGCAACTAGTCACAAAGAGATGACCA (SEQ ID NO. 21).

Figure 4 details the structure of the BPIV3/HPIV3 chimeric viruses of the
 10 invention, and their confirmation by TaqI digestion of RT-PCR products generated from
 virus RNA. In Fig. 4A the genomes of the chimeric cKa and cSF viruses are shown
 schematically (not to scale) relative to that of HPIV3 and BPIV3 parent viruses. Ka- and
 SF-specific regions are indicated by light and dark shading respectively. Arrows above
 the rJS genome indicate the locations of primers used for RT-PCR amplification of
 15 chimeric and parent viruses for the purposes of diagnostic TaqI digestion. These primers
 were directed to regions conserved between HPIV3 and BPIV3 so that they could be used
 for the amplification of HPIV3, BPIV3 and chimeric BPIV3/HPIV3 viruses. In Fig. 4B
 the expected sizes of TaqI digestion products for each virus are shown for a 1898-bp PCR
 product amplified from RNA with the primer pair illustrated in Fig. 4A. This PCR
 20 product is illustrated at the top in Fig. 4B, and the N ORF is indicated as a filled
 rectangle. TaqI fragments unique to each virus and which therefore serve in virus
 identification are indicated with an asterisk. Fig. 4C provides TaqI profiles of PCR
 products containing the PIV3 N coding region of chimeric cKa (left) or cSF (right)
 flanked by those of the HPIV3 and BPIV3 parent viruses. Unique TaqI fragments
 25 diagnostic of virus identity and corresponding to those identified in (4B) are indicated
 with an asterisk. Calculated lengths (bp) of DNA gel bands are indicated.

Figure 5 provides multicycle growth curves of parental and chimeric
 viruses in MDBK (A) or LLC-MK2 (B) cells. Monolayers of bovine MDBK (A) or
 simian LLC-MK2 (B) cells in wells (9.6 cm² each) of a 6 well plate were infected
 30 individually at a multiplicity of infection of 0.01 with the indicated parental or chimeric
 virus. Three replicate infections were performed for each virus. Samples were taken at
 the indicated time points, stored at -70°C, and titered by TCID₅₀ assay in parallel.
 Growth curves are constructed using the average of 3 replicate samples at each time

point. The lower limit of virus detectability was $10^{1.5}$ TCID₅₀/ml, which is indicated by a dotted line.

Figures 6A-6G set forth the complete positive sense nucleotide sequence (SEQ ID NO. 22) of the bovine PIV3 Ka strain.

5 Figures 7A-7G set forth the complete positive sense nucleotide sequence (SEQ ID NO. 23) of the bovine PIV3 SF strain.

Figure 8A provides a schematic depiction of the genomes of chimeric rHPIV3-F_BHN_B and rBPIV3-F_HHN_H viruses, and of their parent viruses, rHPIV3 JS and BPIV3 Ka (not to scale). The F and HN genes were exchanged in a single restriction
10 fragment between rHPIV3 and rBPIV3 using *Sgr*AI and *Bsi*WI sites that had been introduced in front of the M and HN gene end sequences, respectively.

Figure 8B depicts assembly of an antigenomic cDNA for BPIV3 Ka. A full length cDNA was constructed to encode the complete antigenomic sequence of BPIV3 Ka (GenBank accession #AF178654). The cDNA was assembled from subclones
15 derived from reverse transcription (RT) of viral (v)RNA and polymerase chain reaction (PCR) amplification. Multiple subclones of the antigenome were sequenced, and only clones matching the consensus sequence of BPIV3 Ka were used for assembly of the full length clone, with the exception of nt 21 and nt 23, which differ from the published sequence but occur with similar frequency in the virus population..

20 Figure 8C illustrates features of parental and chimeric bovine-human PIV genomes. The genomes of the chimeric rHPIV3 F_BHN_B and rBPIV3 F_HHN_H viruses and those of their parent viruses rHPIV3 JS and BPIV3 Ka are shown schematically (not to scale). Two unique restriction enzyme recognition sites, *Sgr*AI and *Bsi*WI, were introduced near the M and HN gene ends, respectively. The recombinant HPIV3 and
25 BPIV3 viruses bearing these introduced restriction sites were designated rHPIV3s and rBPIV3s as indicated in Figure 8C2. Glycoprotein genes were exchanged between rHPIV3 JS and rBPIV3 Ka. The nucleotide sequence that was mutagenized is shown below each cDNA construct, with the position of the first nucleotide of each sequence indicated. The introduced *Sgr*AI and *Bsi*WI restriction sites are underlined and
30 nucleotides that differ between HPIV3 and BPIV3 and thus identify the origin of the gene inserts are depicted in bold print. Figure 8C, Panel 1, rHPIV3 JS, TCCACCGGTGCA (SEQ ID NO. 4), TAGACAAAAGGG (SEQ ID NO. 24). Figure 8C, Panel 1, rBPIV3 Ka, TCCAACATTGCA (SEQ ID NO. 2); AAGATATAAAGA (SEQ ID NO. 25). Figure 8C, Panel 2 rHPIV3s, CGCACCGGTGTA (SEQ ID NO. 5); TAGACGTACGGG

- (SEQ ID NO. 26). Figure 8C, Panel 2. rBPIV3s, TCCACCGGTGCA (SEQ ID NO. 3); AAGACGTACGGA (SEQ ID NO. 27). Figure 8C, Panel 3. rHPIV3 F_BHN_B, CGCACCGGTGCA (SEQ ID NO. 28); AAGACGTACGGG (SEQ ID NO. 29). Figure 8C, Panel 3. rBPIV3 F_HHN_H, AAGACGTACGGG (SEQ ID NO. 30);
- 5 TAGACGTACGGA (SEQ ID NO. 31).

Figure 9 provides a confirmation of the identity of recombinant viruses by RT-PCR of viral RNA and *Eco* RI digestion. RT-PCR products of viral RNA were prepared with a primer pair that recognized conserved regions on either side of the F and HN genes in both BPIV3 and HPIV3. Digestion with *Eco* RI resulted in a unique pattern

10 of restriction fragments for each of the four viruses. In the schematic diagram on the left, horizontal lines symbolize the amplified viral sequences and vertical bars show the positions of *Eco* RI sites. The expected size of each restriction fragment is indicated above the line. The numbers below each line correspond to the sequence position in the antigenomic RNA of BPIV3 Ka, HPIV3 JS (GenBank accession #AF178654 and

15 Z11575), or of the indicated chimeric derivative. On the right, a 1% agarose gel of the *Eco* RI digestion of PCR products is shown, confirming the identity of parental and chimeric viruses. The asterisks indicate gel bands that contain two restriction fragments that comigrate due to close similarity in size.

Figure 10 depicts multicycle replication of chimeric and parental viruses in

20 simian LLC-MK2 cells. Multicycle replication (the input inoculum had an MOI of 0.01) of the three chimeric viruses rHPIV3-F_BHN_B, rBPIV3-F_HHN_H and rHPIV3-N_B (also referred to as cKa) is compared with the replication of their parental viruses BPIV3 Ka and rHPIV3. The virus titers are shown as mean log₁₀ TCID₅₀/ml ± standard error of triplicate samples. The lower limit of detection of this assay is 10 TCID₅₀, as indicated by

25 the dotted horizontal line.

Figure 11 documents mean titers of chimeric and parental viruses in nasopharyngeal swabs of infected rhesus monkeys over the course of infection. Virus titers are shown as mean TCID₅₀/ml in LLC-MK2 cells ± standard error for groups of 4 or 6 monkeys infected with the same virus. This illustrates the same experiment as shown in

30 Table 3. In panel A, mean titers of rHPIV3-F_BHN_B are compared to rHPIV3 and BPIV3 Ka titers. In panel B, mean rBPIV3-F_HHN_H titers are compared to those of BPIV3 Ka and rHPIV3, which, for the last two viruses, are the same values in panel A but are presented separately to facilitate comparison. Day 5 titers were excluded from the figures

because they were much lower than day 4 and day 6 titers, most likely due to technical problems during the sample collection.

DESCRIPTION OF THE SPECIFIC EMBODIMENTS

5

The present invention provides recombinant parainfluenza virus (PIV) cloned as a chimera of human and bovine PIV genomic or antigenomic sequences to yield a human-bovine chimeric PIV. The chimeric construction of human-bovine PIV yields a viral particle or subviral particle that is infectious in mammals, particularly humans, and
10 useful for generating immunogenic compositions for clinical or veterinary use. Also provided within the invention are novel methods and compositions for designing and producing attenuated, human-bovine chimeric PIV, as well as methods and compositions for the prophylaxis and treatment of PIV infection.

Chimeric human-bovine PIV of the invention are recombinantly
15 engineered to incorporate nucleotide sequences from both human and bovine PIV strains to produce an infectious, chimeric virus or subviral particle. In this manner, candidate vaccine viruses are recombinantly engineered to elicit an immune response against PIV in a mammalian host susceptible to PIV infection, including humans and non-human primates. Human-bovine chimeric PIV according to the invention may elicit an immune
20 response to a specific PIV, e.g., HPIV3, or a polyspecific response against multiple PIVs, e.g., HPIV1 and HPIV3.

Exemplary human-bovine chimeric PIV of the invention incorporate a chimeric PIV genome or antigenome comprising both human and bovine polynucleotide sequences, as well as a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein
25 (P), and a large polymerase protein (L). Additional PIV proteins may be included in various combinations to provide a range of infectious subviral particles, up to a complete viral particle or a viral particle containing supernumerary proteins, antigenic determinants or other additional components.

Chimeric human-bovine PIV of the invention include a partial or complete
30 "background" PIV genome or antigenome derived from or patterned after a human or bovine PIV strain or serotype virus combined with one or more heterologous gene(s) or genome segment(s) of a different PIV strain or serotype virus to form the human-bovine chimeric PIV genome or antigenome. In certain aspects of the invention, chimeric PIV incorporate a partial or complete human PIV (HPIV) background genome or antigenome

combined with one or more heterologous gene(s) or genome segment(s) from a bovine PIV. In alternate aspects of the invention, chimeric PIV incorporate a partial or complete bovine PIV (BPIV) background genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a human PIV.

5 The partial or complete background genome or antigenome typically acts as a recipient backbone or vector into which are imported heterologous genes or genome segments of the counterpart, human or bovine PIV. Heterologous genes or genome segments from the counterpart, human or bovine PIV represent "donor" genes or polynucleotides that are combined with, or substituted within, the background genome or
10 antigenome to yield a human-bovine chimeric PIV that exhibits novel phenotypic characteristics compared to one or both of the contributing PIVs. For example, addition or substitution of heterologous genes or genome segments within a selected recipient PIV strain may result in an increase or decrease in attenuation, growth changes, altered immunogenicity, or other desired phenotypic changes as compared with a corresponding
15 phenotype(s) of the unmodified recipient and/or donor. Genes and genome segments that may be selected for use as heterologous inserts or additions within human-bovine chimeric PIV of the invention include genes or genome segments encoding a PIV N, P, C, D, V, M, SH, where applicable, F, HN and/or L protein(s) or portion(s) thereof. Regulatory regions, such as the extragenic leader or trailer or intergenic regions, are also
20 useful as heterologous inserts or additions.

 The heterologous gene(s) or genome segment(s) may be added or substituted at a position corresponding to a wild-type gene order position of the counterpart gene(s) or genome segment(s) within the partial or complete PIV background genome or antigenome, which counterpart gene or genome segment is thereby replaced or
25 displaced (e.g., to a more promotor-distal position). In yet additional embodiments, the heterologous gene or genome segment is added or substituted at a position that is more promoter-proximal or promotor-distal compared to a wild-type gene order position of the counterpart gene or genome segment within the background genome or antigenome, which enhances or reduces, respectively, expression of the heterologous gene or genome
30 segment.

 The introduction of heterologous immunogenic proteins, domains and epitopes to produce human-bovine chimeric PIV is particularly useful to generate novel immune responses in an immunized host. Addition or substitution of an immunogenic gene or genome segment from one, donor PIV within a recipient genome or antigenome

of a different PIV can generate an immune response directed against the donor subgroup or strain, the recipient subgroup or strain, or against both the donor and recipient subgroup or strain. To achieve this purpose, human-bovine chimeric PIV may also be constructed that express a chimeric protein, e.g., an immunogenic glycoprotein having a cytoplasmic tail and/or transmembrane domain specific to one PIV fused to an ectodomain of a different PIV to provide, e.g., a human-bovine fusion protein, or a fusion protein incorporating domains from two different human PIVs. In a preferred embodiment, a human-bovine chimeric PIV genome or antigenome encodes a chimeric glycoprotein in the recombinant virus or subviral particle having both human and bovine glycoprotein domains or immunogenic epitopes. For example, a heterologous genome segment encoding a glycoprotein ectodomain from a human PIV HN or F glycoprotein may be joined with a polynucleotide sequence (i.e., a genome segment) encoding the corresponding bovine HN or F glycoprotein cytoplasmic and transmembrane domains to form the human-bovine chimeric PIV genome or antigenome.

In other embodiments, human-bovine chimeric PIV useful in a vaccine formulation can be conveniently modified to accommodate antigenic drift in circulating virus. Typically the modification will be in the HN and/or F proteins. This might involve the introduction of one or more point mutations; it might also involve an entire HN or F gene, or a genome segment encoding a particular immunogenic region thereof, from one PIV strain or group is incorporated into a chimeric PIV genome or antigenome cDNA by replacement of a corresponding region in a recipient clone of a different PIV strain or group, or by adding one or more copies of the gene, such that multiple antigenic forms are represented. Progeny virus produced from the modified PIV clone can then be used in vaccination protocols against emerging PIV strains.

Replacement of a human PIV coding sequence or non-coding sequence (e.g., a promoter, gene-end, gene-start, intergenic or other cis-acting element) with a heterologous counterpart yields chimeric PIV having a variety of possible attenuating and other phenotypic effects. In particular, host range and other desired effects arise from substituting a bovine or murine PIV (MPIV) protein, protein domain, gene or genome segment imported within a human PIV background, wherein the bovine or murine gene does not function efficiently in a human cell, e.g., from incompatibility of the heterologous sequence or protein with a biologically interactive human PIV sequence or protein (i.e., a sequence or protein that ordinarily cooperates with the substituted sequence or protein for viral transcription, translation, assembly, etc.) or, more typically

in a host range restriction, with a cellular protein or some other aspect of the cellular milieu which is different between the permissive and less permissive host. In exemplary embodiments, bovine PIV sequences are selected for introduction into human PIV based on known aspects of bovine and human PIV structure and function.

5 HPIV3 is a member of the *Respirovirus* genus of the *Paramyxoviridae* family in the order *Mononegavirales* (Collins et al., 1996, *supra*). HPIV3 is the best characterized of the HPIVs and represents the prototype HPIV. Its genome is a single strand of negative-sense RNA 15462 nucleotides (nt) in length (Galinski et al., Virology 165:499-510, 1988; and Stokes et al., Virus Res. 25:91-103, 1992; each incorporated
10 herein by reference). At least eight proteins are encoded by the PIV3 genome: the nucleocapsid protein N, the phosphoprotein P, the C and D proteins of unknown functions, the matrix protein M, the fusion glycoprotein F, the hemagglutinin-neuraminidase glycoprotein HN, and the large polymerase protein L (Collins et al., 1996, *supra*). A protein containing the V ORF in the P gene might also be produced (Durbin et
15 al., Virology 261:319-333, 1999)

 The M, HN, and F proteins are envelope-associated, and the latter two are surface glycoproteins which, as is the case with each PIV, are the major neutralization and protective antigens (Collins et al., 1996, *supra*). The significant sequence divergence between comparable PIV HN or F proteins among the PIVs is thought to be the basis for
20 the type specificity of the protective immunity (Collins et al., 1996, *supra*; Cook et al., Amer. Jour. Hyg. 77:150-159, 1963; Ray et al., J. Infect. Dis. 162:746-749, 1990; each incorporated herein by reference).

 The HPIV3 genes are each transcribed as a single mRNA that encodes a single protein, with the exception of the P mRNA which contains four ORFs, namely P,
25 C, D and V (Galinski et al., Virology 186:543-550, 1992; and Spriggs et al., J. Gen. Virol. 67:2705-2719, 1986; each incorporated herein by reference). The P and C proteins are translated from separate, overlapping ORFs in the mRNA. Whereas all paramyxoviruses encode a P protein, only members of the genus *Respirovirus* and *Morbillivirus* encode a C protein. Individual viruses vary in the number of proteins expressed from the C ORF and
30 in its importance in replication of the virus *in vitro* and *in vivo*. Sendai virus (SeV) expresses four independently initiated proteins from the C ORF: C', C, Y1, and Y2, whose translational start sites appear in that order in the mRNA (Curran, et al., Enzyme 44:244-249, 1990; Lamb et al., in The Paramyxoviruses, D. Kingsbury, ed., pp. 181-214, Plenum Press, New York, 1991; incorporated herein by reference), whereas HPIV3 and

measles virus (MeV) express only a single C protein (Bellini et al., J. Virol. 53:908-919, 1985; Sanchez et al., Virology 147:177-86, 1985; and Spriggs et al., 1986, *supra*; each incorporated herein by reference).

The PIV3 D protein is a fusion protein of the P and D ORFs, and is
5 expressed from the P gene by the process of co-transcriptional RNA editing in which two nontemplated G residues are added to the P mRNA at the RNA editing site (Galinski et al., 1992, *supra*; and Pelet et al., EMBO J. 10:443-448, 1991; each incorporated herein by reference). BPIV3, the only other paramyxovirus which expresses a D protein, uses RNA editing to express this protein as well as a second protein; the V protein.

10 Nearly all members of the genera *Respirovirus*, *Rubulavirus*, and *Morbillivirus* express a V protein. The one member which clearly does not is HPIV1, which lacks an intact V ORF (Matsuoka et al., J. Virol. 65:3406-3410, 1991, incorporated herein by reference). The V ORF is characterized by the presence of a cysteine-rich domain that is highly conserved (Cattaneo et al., Cell 56:759-764, 1989; Park et al., J.
15 Virol. 66:7033-7039, 1992; Thomas et al., Cell 54:891-902, 1988; and Vidal et al., J. Virol. 64:239-246, 1990; each incorporated herein by reference). The V ORF is maintained in each of the HPIV3 viruses sequenced to date suggesting that this ORF is expressed and retains function for this virus (Galinski et al., Virology 155:46-60, 1986; Spriggs et al., 1986, *supra*; and Stokes et al., 1992, *supra*; incorporated herein by
20 reference).

The BPIV3 V protein is expressed when one nontemplated G residue is added at the RNA editing site (Pelet et al., 1991, *supra*; incorporated herein by reference). However, in the case of HPIV3, two translation stop codons lie between the editing site and the V ORF, and it is not clear whether HPIV3 represents another example in which
25 this ORF is not expressed, or whether it is expressed by some other mechanism. One possibility is that HPIV3 editing also occurs at a second, downstream site in the P gene, although this did not appear to occur in cell culture (Galinski et al., 1992, *supra*). Alternatively, it might be that ribosomes gain access to the V ORF by ribosomal frameshifting. This would be comparable to the situation with the P locus of MV. MV
30 expresses C, P, and V proteins, but also expresses a novel R protein which is synthesized by frameshifting from the P ORF to the V ORF (Liston et al., J. Virol. 69:6742-6750, 1995, incorporated herein by reference). Genetic evidence suggests that the V ORF of HPIV3 is functional (Durbin et al., 1999, *supra*).

Although the means by which HPIV3 expresses its V protein is unclear, the extreme conservation of the its V ORF in different strains suggests that it is indeed expressed. The function of the V protein is not well defined, but V-minus MV and SeV recombinants have been recovered that replicate efficiently *in vitro* but exhibit reduced replication *in vivo* (Delenda, et al., Virology 228:55-62, 1997; Delenda et al., Virology 242:327-337, 1998; Kato et al., 1997a, *supra*; Kato et al., J. Virol. 71:7266-7272, 1997b; and Valsamakis et al., J. Virol. 72:7754-7761, 1998; each incorporated herein by reference).

The viral genome of PIV also contains extragenic leader and trailer regions, possessing all or part of the promoters required for viral replication and transcription, as well as non-coding and intergenic regions. Thus, the PIV genetic map is represented as 3' leader-N-P/C/D/V-M-F-HN-L-5' trailer. Some viruses, such as simian virus 5 and mumps virus, have a gene located between F and HN that encodes a small hydrophobic (SH) protein of unknown function. Transcription initiates at the 3' end and proceeds by a sequential stop-start mechanism that is guided by short conserved motifs found at the gene boundaries. The upstream end of each gene contains a gene-start (GS) signal, which directs initiation of its respective mRNA. The downstream terminus of each gene contains a gene-end (GE) motif which directs polyadenylation and termination. Exemplary sequences have been described for the human PIV3 strains JS (GenBank accession number Z11575, incorporated herein by reference) and Washington (Galinski M.S., in The Paramyxoviruses, Kingsbury, D.W., ed., pp. 537-568, Plenum Press, New York, 1991, incorporated herein by reference), and for the bovine PIV3 strain 910N (GenBank accession number D80487, incorporated herein by reference).

As used herein, "PIV gene" generally refers to a portion of the PIV genome encoding an mRNA and typically begins at the upstream end with a gene-start (GS) signal and ends at the downstream end with the gene-end (GE) signal. The term PIV gene also includes what is described as "translational open reading frame", or ORF, particularly in the case where a protein, such as C, is expressed from an additional ORF rather than from a unique mRNA. To construct human-bovine chimeric PIV of the invention, one or more PIV gene(s) or genome segment(s) may be deleted, inserted or substituted in whole or in part. This means that partial or complete deletions, insertions and substitutions may include open reading frames and/or cis-acting regulatory sequences of any one or more of the PIV genes or genome segments. By "genome segment" is

meant any length of continuous nucleotides from the PIV genome, which might be part of an ORF, a gene, or an extragenic region, or a combination thereof.

The instant invention involves a method for developing live attenuated PIV vaccine candidates based on chimeras between HPIVs and BPIV3. Chimeras are constructed through a cDNA-based virus recovery system. Recombinant viruses made from cDNA replicate independently and are propagated in the same manner as if they were biologically-derived viruses. Preferred human-bovine chimeric PIV vaccine candidates of the invention bear one or more of the major antigenic determinants of one or more human PIV(s), e.g., HPIV1, HPIV2, and/or HPIV3, in a background which is attenuated by the substitution or addition of one or more BPIV genes or genome segments. The major protective antigens of PIVs are their HN and F glycoproteins, although other proteins can also contribute to a protective immune response.

Thus, the invention provides a new basis for attenuating a wild type or mutant parental virus for use as a vaccine against PIV, one which is based on host range effects due to the introduction of one or more gene(s) or genome segment(s) between HPIV and BPIV. There are numerous nucleotide and amino acid sequence differences between BPIV and HPIV, which are reflected in host range differences. For example, between HPIV3 and BPIV3 the percent amino acid identity for each of the following proteins is: N (86%), P (65%), M (93%), F (83%), HN (77%), and L (91%). The host range difference is exemplified by the highly permissive growth of HPIV3 in rhesus monkeys, compared to the restricted replication of two different strains of BPIV3 in the same animal (van Wyke Coelingh et al., 1988, *supra*). Although the basis of the host range differences between HPIV3 and BPIV3 remains to be determined, it is likely that they will involve more than one gene and multiple amino acid differences. The involvement of multiple genes and possibly cis-acting regulatory sequences, each involving multiple amino acid or nucleotide differences, gives a very broad basis for attenuation, one which cannot readily be altered by reversion. This is in contrast to the situation with other live attenuated HPIV3 viruses which are attenuated by one or several point mutations. In this case, reversion of any individual mutation may yield a significant reacquisition of virulence or, in a case where only a single residue specified attenuation, complete reacquisition of virulence.

In exemplary embodiments of the invention described hereinbelow, the background genome or antigenome is an HPIV3 genome or antigenome, and the heterologous gene or genome segment is a N ORF derived from, alternatively, a Ka or SF

strain of BPIV3 (which are 99% related in amino acid sequence). The N ORF of the HPIV3 background antigenome is substituted by the counterpart BPIV3 N ORF yielding a novel recombinant human-bovine chimeric PIV cDNA clone. Replacement of the HPIV3 N ORF of HPIV3 with that of BPIV3 Ka or SF results in a protein with
5 approximately 70 amino acid differences (depending on the strain involved) from that of HPIV3 N. N is one of the more conserved proteins, and substitution of other proteins such as P, singly or in combination, would result in many more amino acid differences. The involvement of multiple genes and genome segments each conferring multiple amino acid or nucleotide differences provides a broad basis for attenuation which is highly
10 stable to reversion.

This mode of attenuation contrasts sharply to current HPIV vaccine candidates that are attenuated by one or more point mutations, where reversion of an individual mutation may yield a significant or complete reacquisition of virulence. In addition, several known attenuating point mutations in HPIV typically yield a temperature
15 sensitive phenotype. One problem with attenuation associated with temperature sensitivity is that the virus can be overly restricted for replication in the lower respiratory tract while being under attenuated in the upper respiratory tract. This is because there is a temperature gradient within the respiratory tract, with temperature being higher (and more restrictive) in the lower respiratory tract and lower (less restrictive) in the upper
20 respiratory tract. The ability of an attenuated virus to replicate in the upper respiratory tract can result in complications including congestion, rhinitis, fever and otitis media, whereas overattenuation in the lower respiratory tract can reduce immunogenicity. Thus, attenuation achieved solely by temperature sensitive mutations may not be ideal. In contrast, host range mutations present in human-bovine chimeric PIV of the invention
25 will not in most cases confer temperature sensitivity. Therefore, the novel method of PIV attenuation provided by the invention will be more stable genetically and phenotypically and less likely to be associated with residual virulence in the upper respiratory tract compared to other known PIV vaccine candidates.

Surprisingly, both the Ka and SF HPIV3/BPIV3 chimeric recombinants
30 involving the N ORF replacement were viable. Since the N protein of Ka or SF strain BPIV3 differs in 70 of 515 amino acid residues, respectively, from that of the JS strain of HPIV3. It was therefore unexpected that a bovine N protein with this level of amino acid sequence divergence could efficiently interact with the HPIV3 RNA, or with other HPIV3 proteins that constitute the functional replicase/transcriptase. Equally surprising was the

finding that the Ka and SF chimeric viruses replicated as efficiently in cell culture as either HPIV3 or BPIV3 parent indicating that the chimeric recombinants did not exhibit gene incompatibilities that restricted replication *in vitro*. This property of efficient replication *in vitro* is important since it permits efficient manufacture of this biological.

5 Also surprising is the observation, based on the studies hereinbelow, that the Ka and the SF HPIV3/BPIV3 chimeric recombinants (termed cKa and cSF), bearing only one bovine gene, are nearly equivalent to their BPIV3 parents in the degree of host range restriction in the respiratory tract of the rhesus monkey. In particular, the cKa and cSF viruses exhibited approximately a 60-fold or 30-fold reduction, respectively, in
10 replication in the upper respiratory tract of rhesus monkeys compared to replication of HPIV3. Based on this finding, it is possible that other BPIV3 genes will also confer desired levels of host range restriction within human-bovine chimeric PIV of the invention. Thus, according to the methods herein, a list of attenuating determinants will be readily identified in heterologous genes and genome segments of both HPIV and BPIV
15 that will confer, in appropriate combination, an optimal level of host range restriction and immunogenicity on human-bovine chimeric PIV selected for vaccine use. In preferred vaccine recombinants, attenuation marked by replication in the lower and/or upper respiratory tract in an accepted animal model for PIV replication in humans, e.g., hamsters or rhesus monkeys, may be reduced by at least about two-fold, more often about
20 5-fold, 10-fold, or 20-fold, and preferably 50-100-fold and up to 1,000-fold or greater overall (e.g., as measured between 3-8 days following infection) compared to growth of the corresponding wild-type or mutant parental PIV strain.

 Confirming the unexpected nature and advantages provided by the human-bovine chimeric PIV of the invention, both the cKa and cSF induced a high level of
25 protection against HPIV3 challenge in the respiratory tract of rhesus monkeys, despite the exceptional degree of restriction of replication exhibited by these viruses in this model for human PIV infection and protection. In particular, previous infection with either chimeric virus induced a high level of resistance to replication of the rJS challenge virus in both the upper and lower respiratory tract. Infection of monkeys with cKa elicited a high degree
30 of protection as indicated by an approximate 300-fold reduction of replication of wild type HPIV3 (rJS) in the upper respiratory tract, and an approximate 1000-fold reduction in the lower tract compared to uninoculated control monkeys. Monkeys infected with cSF manifested a 2000-fold reduction of replication of rJS in the upper respiratory tract, and a 1000-fold reduction in the lower tract compared to uninoculated control monkeys.

The levels of protection elicited by cKa or cSF were comparable to those seen in monkeys previously infected with either the bovine or the human PIV parent. Thus, infection with human-bovine chimeric PIV of the invention provides a high level of protection in the upper and lower respiratory tract of monkeys, and both chimeric viruses
5 represent promising vaccine candidates. In other preferred vaccine recombinants, the immunogenic activity of human-bovine chimeric PIV will be balanced against the level of attenuation to achieve useful vaccine candidates, and will typically be marked by a reduction of replication of challenge virus, e.g., rJS in the lower and/or upper respiratory tract by about 50-100-fold, 100-500-fold, preferably about 500-2,000-fold and up to
10 3,000-fold or greater overall (e.g., as measured between 3-8 days post-challenge). Thus, the recombinant vaccine viruses of the invention maintain immunogenicity while exhibiting concomitant reductions in replication and growth. This surprising assemblage of phenotypic traits is highly desired for vaccine development.

The observation that the N gene from two independent strains of BPIV3
15 confers an attenuation phenotype on HPIV3 for the rhesus monkey indicates that this is likely a property shared by N genes of other BPIV strains. Accordingly, within the methods of the invention any BPIV gene or genome segment, singly or in combination with one or more other BPIV gene(s) or genome segment(s), can be combined with HPIV sequences to produce an attenuated HPIV3/BPIV3 chimeric recombinant virus suitable
20 for use as a vaccine virus. In preferred embodiments, all HPIVs, including HPIV1, HPIV2, HPIV3 and variant strains thereof, are useful recipients for attenuating BPIV gene(s) and/or genome segment(s). In general, the HPIV genes selected for inclusion in a HPIV3/BPIV3 chimeric virus will include one or more of the protective antigens, such as the HN or F glycoproteins.

25 Alternative human-bovine chimeric PIV of the invention will contain protective antigenic determinants of HPIV1 or HPIV2. This may be achieved, for example, by expression of an HN and/or F gene of HPIV1 or HPIV2 as an extra gene(s) in an attenuated HPIV3/BPIV3 chimeric recombinant. Alternatively, it is possible to use a HPIV3/HPIV1 or a HPIV3/HPIV2 antigenic chimeric virus, in which the HPIV1 or
30 HPIV2 HN and/or F genes replace their PIV3 counterpart(s) (Skiadopoulos et al., 1999a, *supra*; Tao et al., 1999, *supra*; and U.S. Patent Application Serial No. 09/083,793, filed May 22, 1998; each incorporated herein by reference), as a recipient or background virus for one or more heterologous, attenuating bovine gene(s) or genome segment(s), for example a Ka or SF N gene or genome segment. Such antigenic chimeric viruses will be

attenuated by the bovine N gene, but will induce immunity to the HPIV1 or HPIV2 virus. In this context, a chimeric PIV1 vaccine candidate has been generated using the PIV3 cDNA rescue system by replacing the PIV3 HN and F open reading frames (ORFs) with those of PIV1 in a PIV3 full-length cDNA that contains the three attenuating mutations in L. The recombinant chimeric virus derived from this cDNA is designated rPIV3-1.cp45L (Skiadopoulos et al., 1998, *supra*; Tao et al., 1998, *supra*; Tao et al., 1999, *supra*). rPIV3-1.cp45L was attenuated in hamsters and induced a high level of resistance to challenge with PIV1. A recombinant chimeric virus, designated rPIV3-1cp45, has also been produced that contains 12 of the 15 cp45 mutations, i.e., excluding the mutations in HN and F, and is highly attenuated in the upper and lower respiratory tract of hamsters (Skiadopoulos et al., 1999a, *supra*).

Still other HPIV/BPIV chimeric recombinants will incorporate two or more BPIV genes or genome segments, in any combination, up to and including all of the BPIV genome other than selected genes or antigenic determinants selected from HN or F gene(s) and genome segment(s), which could come from a human HPIV1, HPIV2, or HPIV3 virus. Yet additional embodiments of the invention are directed to human-bovine chimeric PIV incorporating attenuating genes from other animal PIVs, such as murine PIV1, the canine SV5 PIV2 virus, or another avian or mammalian PIV in combination with a HPIV backbone, alternatively including a chimeric HPIV backbone, from HPIV1, HPIV2, and/or HPIV3.

In other detailed aspects of the invention, human-bovine chimeric PIV are employed as vectors for protective antigens of heterologous pathogens, including other PIVs and non-PIV viruses and non-viral pathogens. Within these aspects, the bovine-human chimeric genome or antigenome comprises a partial or complete PIV "vector genome or antigenome" combined with one or more heterologous genes or genome segments encoding one or more antigenic determinants of one or more heterologous pathogens (see, e.g., U.S. Provisional Patent Application Serial No. 60/170,195, filed December 10, 1999 by Murphy et al., incorporated herein by reference). The heterologous pathogen in this context may be a heterologous PIV and the heterologous gene(s) or genome segment(s) can be selected to encode one or more PIV N, P, C, D, V, M, F, SH (where applicable), HN and/or L protein(s), as well as protein domains, fragments, and immunogenic regions or epitopes. PIV vector vaccines thus constructed may elicit a polyspecific immune response and may be administered simultaneously or in a coordinate administration protocol with other vaccine agents.

In exemplary embodiments of the invention, human-bovine chimeric PIV may comprise a vector genome or antigenome that is a partial or complete HPIV genome or antigenome, which is combined with or is modified to incorporate one or more heterologous genes or genome segments encoding antigenic determinant(s) of one or more heterologous PIV(s), including heterologous HPIVs selected from HPIV1, HPIV2, or HPIV3. In more detailed aspects, the vector genome or antigenome is a partial or complete HPIV3 genome or antigenome and the heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are of one or more heterologous HPIV(s). Typically, the chimeric genome or antigenome incorporates one or more gene(s) or genome segment(s) of a BPIV that specifies attenuation.

In exemplary aspects of the invention, the bovine-human chimeric PIV incorporates one or more HPIV1 or HPIV2 genes or genome segments that encode(s) one or more HN and/or F glycoproteins or antigenic domains, fragments or epitopes thereof within a partial or complete HPIV3 vector genome or antigenome. In more detailed aspects, both HPIV1 genes encoding HN and F glycoproteins are substituted for counterpart HPIV3 HN and F genes to form a chimeric HPIV3-1 vector genome or antigenome. Such recombinant constructs can be used to produce vaccine virus directly, or can be further modified by addition or incorporation of one or more genes or gene segments encoding one or more antigenic determinants. Such constructs for the production of vaccine viruses typically incorporate one or more heterologous gene(s) or genome segment(s) of a BPIV that specifies attenuation, for example an open reading frame (ORF) encoding an attenuating BPIV protein, such as N. Certain human-bovine chimeric PIV of the invention may be employed as vectors for generating specific vaccines to HPIV2, for example wherein a transcription unit comprising an open reading frame (ORF) of an HPIV2 HN gene is added to or incorporated within a chimeric HPIV3-1 vector genome or antigenome and the chimeric construct is attenuated by incorporation of a BPIV gene or genome segment.

Within related aspects of the invention, the vector genome or antigenome is a partial or complete BPIV genome or antigenome, and the heterologous genes or genome segments encoding the antigenic determinant(s) is/are of one or more HPIV(s). Typically, the determinant(s) is/are selected from HPIV1, HPIV2 or HPIV3 HN and F glycoproteins, but antigenic domains, fragments and epitopes of these and other antigenic proteins are also useful. In certain embodiments, one or more genes or genome segments encoding one or more antigenic determinant(s) of HPIV2 is/are added to or substituted

within the partial or complete BPIV vector genome or antigenome. Alternatively, a plurality of heterologous genes or genome segments encoding antigenic determinants of multiple HPIVs may be added to or incorporated within the partial or complete BPIV vector genome or antigenome.

5 In yet additional aspects of the invention, human-bovine chimeric PIV are provided as vectors for a range of non-PIV pathogens (see, e.g., U.S. Provisional Patent Application Serial No. 60/170,195, filed December 10, 1999 by Murphy et al., incorporated herein by reference). The vector genome or antigenome for use within these aspects of the invention may comprise a partial or complete BPIV or HPIV genome or
10 antigenome, and the heterologous pathogen may be selected from measles virus, subgroup A and subgroup B respiratory syncytial viruses, mumps virus, human papilloma viruses, type 1 and type 2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses.

15 For example, a HPIV or BPIV vector genome or antigenome for constructing bovine-human chimeric PIV of the invention may incorporate heterologous antigenic determinant(s) selected from the measles virus HA and F proteins, or antigenic domains, fragments and epitopes thereof. In exemplary embodiments, a transcription unit comprising an open reading frame (ORF) of a measles virus HA gene is added to or
20 incorporated within a BPIV or HPIV3 vector genome or antigenome.

Alternatively bovine-human chimeric PIV of the invention may be used as vectors to incorporate heterologous antigenic determinant(s) from respiratory syncytial virus (RSV), for example by incorporating one or more genes or genome segments that encode(s) RSV F and/or G glycoprotein or immunogenic domain(s) or epitope(s) thereof.
25 In this context, the cloning of RSV cDNA and other disclosure is provided in U.S. Provisional Patent Application No. 60/007,083, filed September 27, 1995; U.S. Patent Application No. 08/720,132, filed September 27, 1996; U.S. Provisional Patent Application No. 60/021,773, filed July 15, 1996; U.S. Provisional Patent Application No. 60/046,141, filed May 9, 1997; U.S. Provisional Patent Application No. 60/047,634, filed
30 May 23, 1997; U.S. Patent Application No. 08/892,403, filed July 15, 1997 (corresponding to International Publication No. WO 98/02530); U.S. Patent Application Serial No. 09/291,894, filed on April 13, 1999; U.S. Provisional Patent Application Serial No. 60/129,006, filed on April 13, 1999; Collins, et al., 1995, *supra*; Bukreyev, et al., *J. Virol.* 70:6634-6641, 1996; Juhasz et al., 1997, *supra*; Durbin et al., 1997a, *supra*; He et

al., 1997, *supra*; Baron et al., 1997, *supra*; Whitehead et al., 1998a, *supra*; Whitehead et al., 1998b, *supra*; Jin et al., 1998, *supra*; and Whitehead et al., 1999, *supra*; and Bukreyev et al., Proc. Natl. Acad. Sci. USA 96:2367-2372, 1999, each incorporated herein by reference in its entirety for all purposes).

5 According to this aspect of the invention, human-bovine chimeric PIV are provided which incorporate at least one antigenic determinant from a heterologous PIV or non-PIV pathogen. For example, one or more individual gene(s) or genome segment(s) of HPIV3 may be replaced with counterpart gene(s) or genome segment(s) from human RSV, or an RSV gene or genome segment can be inserted or added as an supernumerary
10 gene. Alternatively, a selected, heterologous genome segment, e.g. encoding a cytoplasmic tail, transmembrane domain or ectodomain of an RSV glycoprotein, is substituted for a counterpart genome segment in, e.g., the same gene in HPIV3 or within a different gene in HPIV3, or added within a non-coding sequence of the HPIV3 genome or antigenome to yield a chimeric PIV-RSV glycoprotein. In one embodiment, a genome
15 segment from an F gene of human RSV is substituted for a counterpart HPIV3 genome segment to yield constructs encoding chimeric proteins, e.g. fusion proteins having a cytoplasmic tail and/or transmembrane domain of PIV fused to an ectodomain of RSV to yield a novel attenuated virus, and/or a multivalent vaccine immunogenic against both PIV and RSV.

20 As noted above, it is often desirable to adjust the attenuation phenotype in human-bovine chimeric PIV of the invention by introducing additional mutations that increase or decrease attenuation or otherwise alter the phenotype of the chimeric virus. Detailed descriptions of the materials and methods for producing recombinant PIV from cDNA, and for making and testing the full range of mutations and nucleotide
25 modifications set forth herein as supplemental aspects of the present invention, provided in, e.g., Durbin et al., 1997a, *supra*; U.S. Patent Application Serial No. 09/083,793, filed May 22, 1998; U.S. Provisional Application No. 60/047,575, filed May 23, 1997 (corresponding to International Publication No. WO 98/53078), and U.S. Provisional Application No. 60/059,385, filed September 19, 1997; each incorporated herein by
30 reference. In particular, these documents describe methods and procedures for mutagenizing, isolating and characterizing PIV to obtain attenuated mutant strains (e.g., temperature sensitive (ts), cold passaged (cp) cold-adapted (ca), small plaque (sp) and host-range restricted (hr) mutant strains) and for identifying the genetic changes that specify the attenuated phenotype. In conjunction with these methods, the foregoing

documents detail procedures for determining replication, immunogenicity, genetic stability and protective efficacy of biologically derived and recombinantly produced attenuated human PIV in accepted model systems, including murine and non-human primate model systems. In addition, these documents describe general methods for developing and testing immunogenic compositions, including monovalent and bivalent vaccines, for prophylaxis and treatment of PIV infection. Methods for producing infectious recombinant PIV by construction and expression of cDNA encoding a PIV genome or antigenome coexpressed with essential PIV proteins are also described in the above-incorporated documents, which include description of the following exemplary plasmids that may be employed to produce infectious PIV viral clones: p3/7(131) (ATCC 97990); p3/7(131)2G (ATCC 97889); and p218(131) (ATCC 97991); each deposited under the terms of the Budapest Treaty with the American Type Culture Collection (ATCC) of 10801 University Boulevard, Manassas, Virginia 20110-2209, U.S.A., and granted the above identified accession numbers.

Also disclosed in the above-incorporated references are methods for constructing and evaluating infectious recombinant PIV that are modified to incorporate phenotype-specific mutations identified in biologically-derived PIV mutants, e.g., cold passaged (cp), cold adapted (ca), host range restricted (hr), small plaque (sp), and/or temperature sensitive (ts) mutants, for example the JS HPIV3 cp 45 mutant strain. Other mutations may be attenuating without an auxiliary marker phenotype. Mutations identified in these mutants can be readily adopted in human-bovine chimeric PIV. In exemplary embodiments, one or more attenuating mutations occur in the polymerase L protein, e.g., at a position corresponding to Tyr₉₄₂, Leu₉₉₂, or Thr₁₅₅₈ of JS cp45. Preferably, these mutations are incorporated in human-bovine chimeric PIV of the invention by an identical, or conservative, amino acid substitution as identified in the biological mutant. Thus, PIV recombinants may incorporate a mutation wherein Tyr₉₄₂ is replaced by His, Leu₉₉₂ is replaced by Phe, and/or Thr₁₅₅₈ is replaced by Ile. Substitutions that are conservative to these replacement amino acids are also useful to achieve a desired mutant phenotype

Other exemplary mutations adopted from a biologically derived PIV mutant include one or more mutations in the N protein, including specific mutations at a position corresponding to residues Val₉₆ or Ser₃₈₉ of JS cp45. In more detailed aspects, these mutations are represented as Val₉₆ to Ala or Ser₃₈₉ to Ala or substitutions that are conservative thereto. Also useful within recombinant PIV of the invention are amino acid

substitution in the C protein, e.g., a mutation at a position corresponding to Ile₉₆ of JS *cp45*, preferably represented by an identical or conservative substitution of Ile₉₆ to Thr. Further exemplary mutations adopted from biologically derived PIV mutants include one mutation in the M gene such as Pro₁₉₉ in JS *cp45*, or more mutations in the F protein, including mutations adopted from JS *cp45* at a position corresponding to residues Ile₄₂₀ or Ala₄₅₀ of JS *cp45*, preferably represented by acid substitutions Ile₄₂₀ to Val or Ala₄₅₀ to Thr or substitutions conservative thereto. Other human-bovine chimeric PIV within the invention adopt one or more amino acid substitutions in the HN protein, as exemplified hereinbelow by a recombinant PIV adopting a mutation at a position corresponding to residue Val₃₈₄ of JS, preferably represented by the substitution Val₃₈₄ to Ala.

Yet additional examples within this aspect of the invention include human-bovine chimeric PIV which incorporate one or more mutations in noncoding portions of the PIV genome or antigenome, for example in a 3' leader sequence. Exemplary mutations in this context may be engineered at a position in the 3' leader of a recombinant virus at a position corresponding to nucleotide 23, 24, 28, or 45 of JS *cp45*. Yet additional exemplary mutations may be engineered in the N gene start sequence, for example by changing one or more nucleotides in the N gene start sequence, e.g., at a position corresponding to nucleotide 62 of JS *cp45*. In more detailed aspects human-bovine chimeric PIV incorporate a T to C change at nucleotide 23, a C to T change at nucleotide 24, a G to T change at nucleotide 28, and/or a T to A change at nucleotide 45. Additional mutations in extragenic sequences are exemplified by a A to T change in N gene start sequence at a position corresponding to nucleotide 62 of JS.

These foregoing exemplary mutations which can be engineered in a human-bovine chimeric PIV of the invention have been successfully engineered and recovered in recombinant PIV, as represented by the recombinant PIV clones designated rcp45, rcp45 L, rcp45 F, rcp45 M, rcp45 HN, rcp45 C, rcp45 F, rcp45 3'N, 3'NL, and rcp45 3'NCMFHN (Durbin et al., 1997a, *supra*; Skiadopolos et al., 1998, *supra*; Skiadopolos et al., *J. Virol.* **73**:1374-1381, 1999b; U.S. Patent Application Serial No. 09/083,793, filed May 22, 1998; U.S. Provisional Application No. 60/047,575, filed May 23, 1997 (corresponding to International Publication No. WO 98/53078), and U.S. Provisional Application No. 60/059,385, filed September 19, 1997, each incorporated herein by reference). In addition, the above-incorporated references describe construction of chimeric PIV recombinants, e.g., having the HN and F genes of HPIV1 substituted into a partial HPIV3 background genome or antigenome, which is further modified to bear one

or more of the attenuating mutations identified in HPIV3 JS cp45. One such chimeric recombinant incorporates all of the attenuating mutations identified in the L gene of cp45. It has since been shown that all of the cp45 mutations outside of the heterologous (HPIV1) HN and F genes can be incorporated in a HPIV3-l recombinant to yield an
5 attenuated, chimeric vaccine candidate.

From JS cp45 and other biologically derived PIV mutants, a large "menu" of attenuating mutations is provided, each of which can be combined with any other mutation(s) for adjusting the level of attenuation, immunogenicity and genetic stability in a recombinant PIV bearing C, D, and/or V deletion or knock out mutation(s). In this
10 context, many recombinant PIVs of the invention will include one or more, and preferably two or more, mutations from biologically derived PIV mutants, e.g., any one or combination of mutations identified in JS cp45. Preferred PIV recombinants within the invention will incorporate a plurality and up to a full complement of the mutations present in JS cp45 or other biologically derived mutant PIV strains. Preferably, these mutations
15 are stabilized against reversion in human-bovine chimeric PIV by multiple nucleotide substitutions in a codon specifying each mutation.

Additional mutations that may be incorporated in human-bovine chimeric PIV of the invention are mutations, e.g., attenuating mutations, identified in heterologous PIV or more distantly related nonsegmented negative stranded RNA viruses. In
20 particular, attenuating and other desired mutations identified in one negative stranded RNA virus may be "transferred", e.g., introduced by mutagenesis in a corresponding position within the genome or antigenome of the human-bovine chimeric PIV. Briefly, desired mutations in one heterologous negative stranded RNA virus are transferred to the PIV recipient (e.g., bovine or human PIV, respectively). This involves mapping the
25 mutation in the heterologous virus, thus identifying by sequence alignment the corresponding site in the recipient RSV, and mutating the native sequence in the PIV recipient to the mutant genotype (either by an identical or conservative mutation), as described in PCT/US00/09695 filed April 12, 2000 and its priority U.S. Provisional Patent Application Serial No. 60/129,006, filed April 13, 1999, incorporated herein by
30 reference). As this disclosure teaches, it is preferable to modify the recipient genome or antigenome to encode an alteration at the subject site of mutation that corresponds conservatively to the alteration identified in the heterologous mutant virus. For example, if an amino acid substitution marks a site of mutation in the mutant virus compared to the corresponding wild-type sequence, then a similar substitution should be engineered at the

corresponding residue(s) in the recombinant virus. Preferably the substitution will involve an identical or conservative amino acid to the substitute residue present in the mutant viral protein. However, it is also possible to alter the native amino acid residue at the site of mutation non-conservatively with respect to the substitute residue in the mutant protein (e.g., by using any other amino acid to disrupt or impair the function of the wild-type residue).

Negative stranded RNA viruses from which exemplary mutations are identified and transferred into human-bovine chimeric PIV of the invention include other PIVs (e.g., HPIV1, HPIV2, HPIV3, HPIV4A, HPIV4B and BPIV3), RSV, Sendai virus (SeV), Newcastle disease virus (NDV), simian virus 5 (SV5), measles virus (MeV), rinderpest virus, canine distemper virus (CDV), rabies virus (RaV) and vesicular stomatitis virus (VSV), among others.

A variety of exemplary mutations for use within the invention are disclosed in the above-incorporated reference, including but not limited to an amino acid substitution of phenylalanine at position 521 of the RSV L protein corresponding to and therefore transferable to a substitution of phenylalanine (or a conservatively related amino acid) at position 456 of the HPIV3 L protein. In the case of mutations marked by deletions or insertions, these can be introduced as corresponding deletions or insertions into the recombinant virus, either within the background genome or antigenome or within the heterologous gene or genome segment incorporated therein. However the particular size and amino acid sequence of the deleted or inserted protein fragment can vary.

Yet additional human-bovine PIV vaccine candidates within the invention can be achieved by modifying the chimeric PIV genome or antigenome to encode an analogous mutation to an attenuating mutation identified in Sendai virus (SeV). In one example, the attenuating mutation comprises an amino acid substitution of phenylalanine at position 170 of the C protein of SeV. The PIV genome or antigenome is modified to encode an alteration of a conserved residue that corresponds conservatively to the alteration marking the attenuating mutation in the heterologous, SeV mutant. In one embodiment, the mutation is incorporated within a recombinant HPIV3 protein and comprises an amino acid substitution of phenylalanine at position 164 of the C protein of HPIV3.

Various target proteins are amenable to introduction of attenuating mutations from one negative stranded RNA virus at a corresponding site within chimeric human-bovine PIV of the invention. Throughout the order Mononegavirales, five target

proteins are strictly conserved and show moderate to high degrees of sequence identity for specific regions or domains. In particular, all known members of the order share a homologous constellation of five proteins: a nucleocapsid protein (N); a nucleocapsid phosphoprotein (P), a nonglycosylated matrix (M) protein, at least one surface glycoprotein (HN, F, H, or G) and a large polymerase (L) protein. These proteins all represent useful targets for incorporating attenuating mutations by altering one or more conserved residues in a protein of the recombinant virus at a site corresponding to the site of an attenuating mutation identified in the heterologous, mutant virus.

In this context, the methods for transferring heterologous mutations into chimeric human-bovine PIV of the invention are based on identification of an attenuating mutation in a first negative stranded RNA virus. The mutation, identified in terms of mutant versus wild-type sequence at the subject amino acid position(s) marking the site of the mutation, provides an index for sequence comparison against a homologous protein in the chimeric virus (either in the background genome or antigenome or in the heterologous gene or gene segment added or substituted therein) that is the target for recombinant attenuation. The attenuating mutation may be previously known or may be identified by mutagenic and reverse genetics techniques applied to generate and characterize biologically-derived mutant virus. Alternatively, attenuating mutations of interest may be generated and characterized *de novo*, e.g., by site directed mutagenesis and conventional screening methods.

Each attenuating mutation identified in a negative stranded RNA virus provides an index for sequence comparison against a homologous protein in one or more heterologous negative stranded virus(es). In this context, existing sequence alignments may be analyzed, or conventional sequence alignment methods may be employed to yield sequence comparisons for analysis, to identify corresponding protein regions and amino acid positions between the protein bearing the attenuating mutation and a homologous protein of a different virus that is the target recombinant virus for attenuation. Where one or more residues marking the attenuating mutation have been altered from a "wild-type" identity that is conserved at the corresponding amino acid position(s) in the target human-bovine chimeric virus protein, the genome or antigenome of the target virus is recombinantly modified to encode an amino acid deletion, substitution, or insertion to alter the conserved residue(s) in the target virus protein and thereby confer an analogous, attenuated phenotype on the recombinant virus.

Within this rational design method for constructing attenuated recombinant negative stranded viruses, the wild-type identity of residue(s) at amino acid positions marking an attenuating mutation in one negative stranded RNA virus may be conserved strictly, or by conservative substitution, at the corresponding amino acid position(s) in the target, human-bovine chimeric virus protein. Thus, the corresponding residue(s) in the target virus protein may be identical, or may be conservatively related in terms of amino acid side-group structure and function, to the wild-type residue(s) found to be altered by the attenuating mutation in the heterologous, mutant virus. In either case, analogous attenuation in the recombinant virus may be achieved according to the methods of the invention by modifying the recombinant genome or antigenome of the target virus to encode the amino acid deletion, substitution, or insertion to alter the conserved residue(s).

In this context, it is preferable to modify the genome or antigenome to encode an alteration of the conserved residue(s) that corresponds conservatively to the alteration marking the attenuating mutation in the heterologous, mutant virus. For example, if an amino acid substitution marks a site of mutation in the mutant virus compared to the corresponding wild-type sequence, then a substitution should be engineered at the corresponding residue(s) in the recombinant virus. Preferably the substitution will be identical or conservative to the substitute residue present in the mutant viral protein. However, it is also possible to alter the native amino acid residue at the site of mutation non-conservatively with respect to the substitute residue in the mutant protein (e.g., by using any other amino acid to disrupt or impair the identity and function of the wild-type residue). In the case of mutations marked by deletions or insertions, these can be transferred as corresponding deletions or insertions into the recombinant virus, however the particular size and amino acid sequence of the deleted or inserted protein fragment can vary.

Within alternative aspects of the invention, mutations thus transferred from heterologous mutant negative stranded viruses may confer a variety of phenotypes within human-bovine chimeric PIV of the invention, in addition to or associated with the desired, an attenuated phenotype. Thus, exemplary mutations incorporated within recombinant proteins of the virus may confer temperature sensitive (*ts*), cold-adapted (*ca*), small plaque (*sp*), or host range restricted (*hr*) phenotypes, or a change in growth or immunogenicity, in addition to or associated with the attenuated phenotype.

Attenuating mutations in biologically derived PIV and other nonsegmented negative stranded RNA viruses for incorporation within human-bovine chimeric PIV may

occur naturally or may be introduced into wild-type PIV strains by well known mutagenesis procedures. For example, incompletely attenuated parental PIV strains can be produced by chemical mutagenesis during virus growth in cell cultures to which a chemical mutagen has been added, by selection of virus that has been subjected to
5 passage at suboptimal temperatures in order to introduce growth restriction mutations, or by selection of a mutagenized virus that produces small plaques (sp) in cell culture, as described in the above incorporated references.

By "biologically derived PIV" is meant any PIV not produced by recombinant means. Thus, biologically derived PIV include all naturally occurring PIV,
10 including, e.g., naturally occurring PIV having a wild-type genomic sequence and PIV having allelic or mutant genomic variations from a reference wild-type RSV sequence, e.g., PIV having a mutation specifying an attenuated phenotype. Likewise, biologically derived PIV include PIV mutants derived from a parental PIV by, *inter alia*, artificial mutagenesis and selection procedures.

15 As noted above, production of a sufficiently attenuated biologically derived PIV mutant can be accomplished by several known methods. One such procedure involves subjecting a partially attenuated virus to passage in cell culture at progressively lower, attenuating temperatures. For example, partially attenuated mutants are produced by passage in cell cultures at suboptimal temperatures. Thus, a cp mutant or
20 other partially attenuated PIV strain is adapted to efficient growth at a lower temperature by passage in culture. This selection of mutant PIV during cold-passage substantially reduces any residual virulence in the derivative strains as compared to the partially attenuated parent.

Alternatively, specific mutations can be introduced into biologically
25 derived PIV by subjecting a partially attenuated parent virus to chemical mutagenesis, e.g., to introduce ts mutations or, in the case of viruses which are already ts, additional ts mutations sufficient to confer increased attenuation and/or stability of the ts phenotype of the attenuated derivative. Means for the introduction of ts mutations into PIV include replication of the virus in the presence of a mutagen such as 5-fluorouridine according to
30 generally known procedures. Other chemical mutagens can also be used. Attenuation can result from a ts mutation in almost any PIV gene, although a particularly amenable target for this purpose has been found to be the polymerase (L) gene.

The level of temperature sensitivity of replication in exemplary attenuated PIV for use within the invention is determined by comparing its replication at a

permissive temperature with that at several restrictive temperatures. The lowest temperature at which the replication of the virus is reduced 100-fold or more in comparison with its replication at the permissive temperature is termed the shutoff temperature. In experimental animals and humans, both the replication and virulence of PIV correlate with the mutant's shutoff temperature.

The JS cp45 HPIV3 mutant has been found to be relatively stable genetically, highly immunogenic, and satisfactorily attenuated. Nucleotide sequence analysis of this biologically derived and recombinant viruses incorporating various individual and combined mutations found therein, indicates that each level of increased attenuation is associated with specific nucleotide and amino acid substitutions. The above-incorporated references also disclose how to routinely distinguish between silent incidental mutations and those responsible for phenotype differences by introducing the mutations, separately and in various combinations, into the genome or antigenome of infectious PIV clones. This process coupled with evaluation of phenotype characteristics of parental and derivative virus identifies mutations responsible for such desired characteristics as attenuation, temperature sensitivity, cold-adaptation, small plaque size, host range restriction, etc.

Mutations thus identified are compiled into a "menu" and are then introduced as desired, singly or in combination, to adjust a human-bovine chimeric PIV to an appropriate level of attenuation, immunogenicity, genetic resistance to reversion from an attenuated phenotype, etc., as desired. In accordance with the foregoing description, the ability to produce infectious PIV from cDNA permits introduction of specific engineered changes within human-bovine chimeric PIV. In particular, infectious, recombinant PIVs are employed for identification of specific mutation(s) in biologically derived, attenuated PIV strains, for example mutations which specify ts, ca, att and other phenotypes. Desired mutations are thus identified and introduced into recombinant, human-bovine chimeric PIV vaccine strains. The capability of producing virus from cDNA allows for routine incorporation of these mutations, individually or in various selected combinations, into a full-length cDNA clone, whereafter the phenotypes of rescued recombinant viruses containing the introduced mutations can be readily determined.

By identifying and incorporating specific, biologically derived mutations associated with desired phenotypes, e.g., a cp or ts phenotype, into infectious PIV clones, the invention provides for other, site-specific modifications at, or within close proximity

to, the identified mutation. Whereas most attenuating mutations produced in biologically derived PIV are single nucleotide changes, other "site specific" mutations can also be incorporated by recombinant techniques into biologically derived or recombinant PIV. As used herein, site-specific mutations include insertions, substitutions, deletions or rearrangements of from 1 to 3, up to about 5-15 or more altered nucleotides (e.g., altered from a wild-type PIV sequence, from a sequence of a selected mutant PIV strain, or from a parent recombinant PIV clone subjected to mutagenesis). Such site-specific mutations may be incorporated at, or within the region of, a selected, biologically derived point mutation. Alternatively, the mutations can be introduced in various other contexts within a PIV clone, for example at or near a cis-acting regulatory sequence or nucleotide sequence encoding a protein active site, binding site, immunogenic epitope, etc. Site-specific PIV mutants typically retain a desired attenuating phenotype, but may additionally exhibit altered phenotypic characteristics unrelated to attenuation, e.g., enhanced or broadened immunogenicity, and/or improved growth. Further examples of desired, site-specific mutants include recombinant PIV designed to incorporate additional, stabilizing nucleotide mutations in a codon specifying an attenuating point mutation. Where possible, two or more nucleotide substitutions are introduced at codons that specify attenuating amino acid changes in a parent mutant or recombinant PIV clone, yielding a biologically derived or recombinant PIV having genetic resistance to reversion from an attenuated phenotype. In other embodiments, site-specific nucleotide substitutions, additions, deletions or rearrangements are introduced upstream or downstream, e. g., from 1 to 3, 5-10 and up to 15 nucleotides or more 5' or 3', relative to a targeted nucleotide position, e.g., to construct or ablate an existing cis-acting regulatory element.

In addition to single and multiple point mutations and site-specific mutations, changes to the human-bovine chimeric PIV disclosed herein include deletions, insertions, substitutions or rearrangements of one or more gene(s) or genome segment(s). Particularly useful are deletions involving one or more gene(s) or genome segment(s), which deletions have been shown to yield additional desired phenotypic effects for adjusting the characteristics of human-bovine chimeric PIV within the invention. Thus, U.S. Patent Application Serial No. 09/350,821, filed by Durbin et al. on July 9, 1999) describes methods and compositions whereby expression of one or more HPIV genes, exemplified by the C, D, and/or V ORFs, is reduced or ablated by modifying the PIV genome or antigenome to incorporate a mutation that alters the coding assignment of an

initiation codon or mutation(s) that introduce one or one or more stop codon(s).

Alternatively, one or more of the C, D, and/or V ORFs can be deleted in whole or in part to render the corresponding protein(s) partially or entirely non-functional or to disrupt protein expression altogether. Recombinant PIV having such mutations in C, D, and/or V, or other non-essential gene(s), possess highly desirable phenotypic characteristics for vaccine development. For example, these modifications may specify one or more desired phenotypic changes including (i) altered growth properties in cell culture, (ii) attenuation in the upper and/or lower respiratory tract of mammals, (iii) a change in viral plaque size, (iv) a change in cytopathic effect, and (v) a change in immunogenicity. One such exemplary "knock out" mutant lacking C ORF expression, designated rC-KO, was able to induce a protective immune response against wild type HPIV3 challenge in a non-human primate model despite its beneficial attenuation phenotype.

Thus, in more detailed aspects of the instant invention, human-bovine chimeric PIV incorporate deletion or knock out mutations in a C, D, and/or V ORF(s) which alters or ablates expression of the selected gene(s) or genome segment(s). This can be achieved, e.g., by introducing a frame shift mutation or termination codon within a selected coding sequence, altering translational start sites, changing the position of a gene or introducing an upstream start codon to alter its rate of expression, changing GS and/or GE transcription signals to alter phenotype, or modifying an RNA editing site (e.g., growth, temperature restrictions on transcription, etc.). In more detailed aspects of the invention, human-bovine chimeric PIVs are provided in which expression of one or more gene(s), e.g., a C, D, and/or V ORF(s), is ablated at the translational or transcriptional level without deletion of the gene or of a segment thereof, by, e.g., introducing multiple translational termination codons into a translational open reading frame (ORF), altering an initiation codon, or modifying an editing site. These forms of knock-out virus will often exhibit reduced growth rates and small plaque sizes in tissue culture. Thus, these methods provide yet additional, novel types of attenuating mutations which ablate expression of a viral gene that is not one of the major viral protective antigens. In this context, knock-out virus phenotypes produced without deletion of a gene or genome segment can be alternatively produced by deletion mutagenesis, as described, to effectively preclude correcting mutations that may restore synthesis of a target protein. Several other gene knock-outs for the C, D, and/or V ORF(s) deletion and knock out mutants can be made using alternate designs and methods that are well known in the art (as described, for example, in (Kretschmer et al., *Virology* 216:309-316, 1996; Radecke

et al., Virology 217:418-421, 1996; and Kato et al., 1987a, *supra*; and Schneider et al., 1997, *supra*; each incorporated herein by reference).

These and other nucleotide modifications in human-bovine chimeric PIV may alter small numbers of bases (e.g., from 15-30 bases, up to 35-50 bases or more),
5 large blocks of nucleotides (e.g., 50-100, 100-300, 300-500, 500-1,000 bases), or nearly complete or complete genes (e.g., 1,000-1,500 nucleotides, 1,500-2,500 nucleotides, 2,500-5,000, nucleotides, 5,000-6,500 nucleotides or more) in the donor or recipient genome or antigenome, depending upon the nature of the change (i.e., a small number of bases may be changed to insert or ablate an immunogenic epitope or change a small
10 genome segment, whereas large block(s) of bases are involved when genes or large genome segments are added, substituted, deleted or rearranged.

In related aspects, the invention provides for supplementation of mutations adopted into a recombinant PIV clone from biologically derived PIV, e.g., cp and ts mutations, with additional types of mutations involving the same or different genes in a
15 further modified PIV clone. Each of the PIV genes can be selectively altered in terms of expression levels, or can be added, deleted, substituted or rearranged, in whole or in part, alone or in combination with other desired modifications, to yield a human-bovine chimeric PIV exhibiting novel vaccine characteristics. Thus, in addition to or in combination with attenuating mutations adopted from biologically derived PIV mutants,
20 the present invention also provides a range of additional methods for attenuating or otherwise modifying the phenotype of human-bovine chimeric PIV based on recombinant engineering of infectious PIV clones. A variety of alterations can be produced in an isolated polynucleotide sequence encoding a targeted gene or genome segment, including a donor or recipient gene or genome segment in a chimeric PIV genome or antigenome
25 for incorporation into infectious clones. More specifically, to achieve desired structural and phenotypic changes in recombinant PIV, the invention allows for introduction of modifications which delete, substitute, introduce, or rearrange a selected nucleotide or plurality of nucleotides from a parent genome or antigenome, as well as mutations which delete, substitute, introduce or rearrange whole gene(s) or genome segment(s), within a
30 human-bovine chimeric PIV clone.

Thus provided are modifications in the human-bovine chimeric PIV which simply alter or ablate expression of a selected gene, e.g., by introducing a termination codon within a selected PIV coding sequence or altering its translational start site or RNA editing site, changing the position of a PIV gene relative to an operably linked promoter,

introducing an upstream start codon to alter rates of expression, modifying (e.g., by changing position, altering an existing sequence, or substituting an existing sequence with a heterologous sequence) GS and/or GE transcription signals to alter phenotype (e.g., growth, temperature restrictions on transcription, etc.), and various other deletions, substitutions, additions and rearrangements that specify quantitative or qualitative changes in viral replication, transcription of selected gene(s), or translation of selected RNA(s). In this context, any PIV gene or genome segment which is not essential for growth can be ablated or otherwise modified in a recombinant PIV to yield desired effects on virulence, pathogenesis, immunogenicity and other phenotypic characters. As for coding sequences, noncoding, leader, trailer and intergenic regions can be similarly deleted, substituted or modified and their phenotypic effects readily analyzed, e.g., by the use of minireplicons and recombinant PIV.

In addition, a variety of other genetic alterations can be produced in a PIV genome or antigenome for incorporation into human-bovine chimeric PIV, alone or together with one or more attenuating mutations adopted from a biologically derived mutant PIV, e.g., to adjust growth, attenuation, immunogenicity, genetic stability or provide other advantageous structural and/or phenotypic effects. These additional types of mutations are also disclosed in the foregoing incorporated references and can be readily engineered into human-bovine chimeric PIV of the invention.

In addition to these changes, the order of genes in a human-bovine chimeric PIV can be changed, a PIV genome promoter replaced with its antigenome counterpart, portions of genes removed or substituted, and even entire genes deleted. Different or additional modifications in the sequence can be made to facilitate manipulations, such as the insertion of unique restriction sites in various intergenic regions or elsewhere. Nontranslated gene sequences can be removed to increase capacity for inserting foreign sequences.

Other mutations for incorporation into human-bovine chimeric PIV of the invention include mutations directed toward cis-acting signals, which can be identified, e.g., by mutational analysis of PIV minigenomes. For example, insertional and deletional analysis of the leader and trailer and flanking sequences identifies viral promoters and transcription signals and provides a series of mutations associated with varying degrees of reduction of RNA replication or transcription. Saturation mutagenesis (whereby each position in turn is modified to each of the nucleotide alternatives) of these cis-acting signals also has identified many mutations which affect RNA replication or transcription.

Any of these mutations can be inserted into a human-bovine chimeric PIV antigenome or genome as described herein. Evaluation and manipulation of trans-acting proteins and cis-acting RNA sequences using the complete antigenome cDNA is assisted by the use of PIV minigenomes as described in the above-incorporated references.

5 Additional mutations within the human-bovine chimeric PIV involve replacement of the 3' end of genome with its counterpart from antigenome, which is associated with changes in RNA replication and transcription. In one exemplary embodiment, the level of expression of specific PIV proteins, such as the protective HN and/or F antigens, can be increased by substituting the natural sequences with ones which
10 have been made synthetically and designed to be consistent with efficient translation. In this context, it has been shown that codon usage can be a major factor in the level of translation of mammalian viral proteins (Hans et al., Current Biol. 6:315-324, 1996, incorporated herein by reference). Optimization by recombinant methods of the codon usage of the mRNAs encoding the HN and F proteins of PIV, which are the major
15 protective antigens, will provide improved expression for these genes.

In another exemplary embodiment, a sequence surrounding a translational start site (preferably including a nucleotide in the -3 position) of a selected PIV gene is modified, alone or in combination with introduction of an upstream start codon, to modulate PIV gene expression by specifying up- or down-regulation of translation
20 (Kozak et al., J. Mol. Biol. 196:947-950, 1987). Alternatively, or in combination with other PIV modifications disclosed herein, gene expression of a human-bovine chimeric PIV can be modulated by altering a transcriptional GS or GE signal of any selected gene(s) of the virus. In alternative embodiments, levels of gene expression in the human-bovine chimeric PIV are modified at the level of transcription. In one aspect, the position
25 of a selected gene in the PIV gene map can be changed to a more promoter-proximal or promoter-distal position, whereby the gene will be expressed more or less efficiently, respectively. According to this aspect, modulation of expression for specific genes can be achieved yielding reductions or increases of gene expression from two-fold, more typically four-fold, up to ten-fold or more compared to wild-type levels often attended by
30 a commensurate decrease in expression levels for reciprocally, positionally substituted genes. These and other transpositioning changes yield novel human-bovine chimeric PIV having attenuated phenotypes, for example due to decreased expression of selected viral proteins involved in RNA replication, or having other desirable properties such as increased antigen expression.

Infectious human-bovine chimeric PIV clones of the invention can also be engineered according to the methods and compositions disclosed herein to enhance immunogenicity and induce a level of protection greater than that provided by infection with a wild-type PIV or a parent PIV. For example, an immunogenic epitope from a heterologous PIV strain or type, or from a non-PIV source such as RSV, can be added to a recombinant clone by appropriate nucleotide changes in the polynucleotide sequence encoding the genome or antigenome. Alternatively, mutant PIV of the invention can be engineered to add or ablate (e.g., by amino acid insertion, substitution or deletion) immunogenic proteins, protein domains, or forms of specific proteins associated with desirable or undesirable immunological reactions.

Within the methods of the invention, additional genes or genome segments may be inserted into or proximate to the human-bovine chimeric PIV genome or antigenome. These genes may be under common control with recipient genes, or may be under the control of an independent set of transcription signals. Genes of interest include the PIV genes identified above, as well as non-PIV genes. Non-PIV genes of interest include those encoding cytokines (e.g., IL-2 through IL-18, especially IL-2, IL-6 and IL-12, IL-18, etc.), gamma-interferon, and proteins rich in T helper cell epitopes. These additional proteins can be expressed either as a separate protein, or as a supernumerary copy of an existing PIV proteins, such as HN or F. This provides the ability to modify and improve the immune responses against PIV both quantitatively and qualitatively.

Deletions, insertions, substitutions and other mutations involving changes of whole viral genes or genome segments within a human-bovine chimeric PIV yield highly stable vaccine candidates, which are particularly important in the case of immunosuppressed individuals. Many of these changes will result in attenuation of resultant vaccine strains, whereas others will specify different types of desired phenotypic changes. For example, accessory (i.e., not essential for in vitro growth) genes are excellent candidates to encode proteins that specifically interfere with host immunity (see, e.g., Kato et al., 1997a, *supra*). Ablation of such genes in vaccine viruses is expected to reduce virulence and pathogenesis and/or improve immunogenicity.

In another aspect of the invention, compositions (e.g., isolated polynucleotides and vectors incorporating human-bovine chimeric PIV-encoding cDNA) are provided for producing an isolated infectious PIV. Using these compositions and methods, infectious PIV are generated from a PIV genome or antigenome, a nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), and a large (L) polymerase protein. In

related aspects of the invention, compositions and methods are provided for introducing the aforementioned structural and phenotypic changes into a recombinant PIV to yield infectious, attenuated vaccine viruses.

Introduction of the foregoing defined mutations into an infectious, human-bovine chimeric PIV clone can be achieved by a variety of well known methods. By “infectious clone” with regard to DNA is meant cDNA or its product, synthetic or otherwise, which can be transcribed into genomic or antigenomic RNA capable of serving as template to produce the genome of an infectious virus or subviral particle. Thus, defined mutations can be introduced by conventional techniques (e.g., site-directed mutagenesis) into a cDNA copy of the genome or antigenome. The use of antigenome or genome cDNA subfragments to assemble a complete antigenome or genome cDNA as described herein has the advantage that each region can be manipulated separately (smaller cDNAs are easier to manipulate than large ones) and then readily assembled into a complete cDNA. Thus, the complete antigenome or genome cDNA, or any subfragment thereof can be used as template for oligonucleotide-directed mutagenesis. This can be through the intermediate of a single-stranded phagemid form, such as using the Muta-gene® kit of Bio-Rad Laboratories (Richmond, CA) or a method using a double-stranded plasmid directly as template such as the Chameleon mutagenesis kit of Stratagene (La Jolla, CA), or by the polymerase chain reaction employing either an oligonucleotide primer or template which contains the mutation(s) of interest. A mutated subfragment can then be assembled into the complete antigenome or genome cDNA. A variety of other mutagenesis techniques are known and available for use in producing the mutations of interest in the PIV antigenome or genome cDNA. Mutations can vary from single nucleotide changes to replacement of large cDNA pieces containing one or more genes or genome regions.

Thus, in one illustrative embodiment mutations are introduced by using the Muta-gene phagemid *in vitro* mutagenesis kit available from Bio-Rad. In brief, cDNA encoding a portion of a PIV genome or antigenome is cloned into the plasmid pTZ18U, and used to transform CJ236 cells (Life Technologies). Phagemid preparations are prepared as recommended by the manufacturer. Oligonucleotides are designed for mutagenesis by introduction of an altered nucleotide at the desired position of the genome or antigenome. The plasmid containing the genetically altered genome or antigenome fragment is then amplified and the mutated piece is then reintroduced into the full-length genome or antigenome clone.

Infectious PIV of the invention are produced by intracellular or cell-free coexpression of one or more isolated polynucleotide molecules that encode a PIV genome or antigenome RNA, together with one or more polynucleotides encoding viral proteins necessary to generate a transcribing, replicating nucleocapsid. Among the viral proteins
5 useful for coexpression to yield infectious PIV are the major nucleocapsid protein (N) protein, nucleocapsid phosphoprotein (P), large (L) polymerase protein, fusion protein (F), hemagglutinin-neuraminidase glycoprotein (HN), and matrix (M) protein. Also useful in this context are products of the C, D and V ORFs of PIV.

cDNAs encoding a PIV genome or antigenome are constructed for
10 intracellular or *in vitro* coexpression with the necessary viral proteins to form infectious PIV. By "PIV antigenome" is meant an isolated positive-sense polynucleotide molecule which serves as a template for synthesis of progeny PIV genome. Preferably a cDNA is constructed which is a positive-sense version of the PIV genome corresponding to the replicative intermediate RNA, or antigenome, so as to minimize the possibility of
15 hybridizing with positive-sense transcripts of complementing sequences encoding proteins necessary to generate a transcribing, replicating nucleocapsid.

In some embodiments of the invention the genome or antigenome of a recombinant PIV (rPIV) need only contain those genes or portions thereof necessary to render the viral or subviral particles encoded thereby infectious. Further, the genes or
20 portions thereof may be provided by more than one polynucleotide molecule, i.e., a gene may be provided by complementation or the like from a separate nucleotide molecule. In other embodiments, the PIV genome or antigenome encodes all functions necessary for viral growth, replication, and infection without the participation of a helper virus or viral function provided by a plasmid or helper cell line.

By "recombinant PIV" is meant a PIV or PIV-like viral or subviral particle
25 derived directly or indirectly from a recombinant expression system or propagated from virus or subviral particles produced therefrom. The recombinant expression system will employ a recombinant expression vector which comprises an operably linked transcriptional unit comprising an assembly of at least a genetic element or elements
30 having a regulatory role in PIV gene expression, for example, a promoter, a structural or coding sequence which is transcribed into PIV RNA, and appropriate transcription initiation and termination sequences.

To produce infectious PIV from a cDNA-expressed PIV genome or antigenome, the genome or antigenome is coexpressed with those PIV N, P and L

proteins necessary to (i) produce a nucleocapsid capable of RNA replication, and (ii) render progeny nucleocapsids competent for both RNA replication and transcription. Transcription by the genome nucleocapsid provides the other PIV proteins and initiates a productive infection. Alternatively, additional PIV proteins needed for a productive infection can be supplied by coexpression.

Synthesis of PIV antigenome or genome together with the above-mentioned viral proteins can also be achieved *in vitro* (cell-free), e.g., using a combined transcription-translation reaction, followed by transfection into cells. Alternatively, antigenome or genome RNA can be synthesized *in vitro* and transfected into cells expressing PIV proteins.

In certain embodiments of the invention, complementing sequences encoding proteins necessary to generate a transcribing, replicating PIV nucleocapsid are provided by one or more helper viruses. Such helper viruses can be wild type or mutant. Preferably, the helper virus can be distinguished phenotypically from the virus encoded by the PIV cDNA. For example, it is desirable to provide monoclonal antibodies which react immunologically with the helper virus but not the virus encoded by the PIV cDNA. Such antibodies can be neutralizing antibodies. In some embodiments, the antibodies can be used in affinity chromatography to separate the helper virus from the recombinant virus. To aid the procurement of such antibodies, mutations can be introduced into the PIV cDNA to provide antigenic diversity from the helper virus, such as in the HN or F glycoprotein genes.

In alternate embodiments of the invention, the N, P, L and other desired PIV proteins are encoded by one or more non-viral expression vectors, which can be the same or separate from that which encodes the genome or antigenome. Additional proteins may be included as desired, each encoded by its own vector or by a vector encoding one or more of the N, P, L and other desired PIV proteins, or the complete genome or antigenome. Expression of the genome or antigenome and proteins from transfected plasmids can be achieved, for example, by each cDNA being under the control of a promoter for T7 RNA polymerase, which in turn is supplied by infection, transfection or transduction with an expression system for the T7 RNA polymerase, e.g., a vaccinia virus MVA strain recombinant which expresses the T7 RNA polymerase (Wyatt et al., Virology 210:202-205, 1995, incorporated herein by reference in its entirety). The viral proteins, and/or T7 RNA polymerase, can also be provided by transformed mammalian cells or by transfection of preformed mRNA or protein.

A PIV antigenome may be constructed for use in the present invention by, e.g., assembling cloned cDNA segments, representing in aggregate the complete antigenome, by polymerase chain reaction or the like (PCR; described in, e.g., U.S. Patent Nos. 4,683,195 and 4,683,202, and *PCR Protocols: A Guide to Methods and Applications*, Innis et al., eds., Academic Press, San Diego, 1990; each incorporated
5 herein by reference in its entirety) of reverse-transcribed copies of PIV mRNA or genome RNA. For example, a first construct is generated which comprises cDNAs containing the left hand end of the antigenome, spanning from an appropriate promoter (e.g., T7 RNA polymerase promoter) and assembled in an appropriate expression vector, such as a
10 plasmid, cosmid, phage, or DNA virus vector. The vector may be modified by mutagenesis and/or insertion of synthetic polylinker containing unique restriction sites designed to facilitate assembly. For ease of preparation the N, P, L and other desired PIV proteins can be assembled in one or more separate vectors. The right hand end of the antigenome plasmid may contain additional sequences as desired, such as a flanking
15 ribozyme and tandem T7 transcriptional terminators. The ribozyme can be hammerhead type (e.g., Grosfeld et al., *J. Virol.* 69:5677-5686, 1995), which would yield a 3' end containing a single nonviral nucleotide, or can be any of the other suitable ribozymes such as that of hepatitis delta virus (Perrotta et al., *Nature* 350:434-436, 1991),
incorporated herein by reference in its entirety) which would yield a 3' end free of non-
20 PIV nucleotides. The left- and right-hand ends are then joined via a common restriction site.

A variety of nucleotide insertions, deletions and rearrangements can be made in the PIV genome or antigenome during or after construction of the cDNA. For example, specific desired nucleotide sequences can be synthesized and inserted at
25 appropriate regions in the cDNA using convenient restriction enzyme sites. Alternatively, such techniques as site-specific mutagenesis, alanine scanning, PCR mutagenesis, or other such techniques well known in the art can be used to introduce mutations into the cDNA.

Alternative means to construct cDNA encoding the genome or antigenome
30 include reverse transcription-PCR using improved PCR conditions (e.g., as described in Cheng et al., *Proc. Natl. Acad. Sci. USA* 91:5695-5699, 1994, incorporated herein by reference) to reduce the number of subunit cDNA components to as few as one or two pieces. In other embodiments different promoters can be used (e.g., T3, SP6) or different

ribozymes (e.g., that of hepatitis delta virus. Different DNA vectors (e.g., cosmids) can be used for propagation to better accommodate the larger size genome or antigenome.

Isolated polynucleotides (e.g., cDNA) encoding the genome or antigenome may be inserted into appropriate host cells by transfection, electroporation, mechanical
5 insertion, transduction or the like, into cells which are capable of supporting a productive PIV infection, e.g., HEp-2, FRhL-DBS2, LLC-MK2, MRC-5, and Vero cells. Transfection of isolated polynucleotide sequences may be introduced into cultured cells by, for example, calcium phosphate-mediated transfection (Wigler et al., Cell 14:725, 1978; Corsaro and Pearson, Somatic Cell Genetics 7:603, 1981; Graham and Van der Eb,
10 Virology 52:456, 1973), electroporation (Neumann et al., EMBO J. 1:841-845, 1982), DEAE-dextran mediated transfection (Ausubel et al., ed., Current Protocols in Molecular Biology, John Wiley and Sons, Inc., New York, 1987), cationic lipid-mediated transfection (Hawley-Nelson et al., Focus 15:73-79, 1993) or a commercially available transfection reagent, e.g., LipofectACE® (Life Technologies, Gaithersburg, MD) or the
15 like (each of the foregoing references are incorporated herein by reference in its entirety).

As noted above, in some embodiments of the invention the N, P, L and other desired PIV proteins are encoded by one or more helper viruses which is phenotypically distinguishable from that which encodes the genome or antigenome. The N, P, L and other desired PIV proteins can also be encoded by one or more expression
20 vectors which can be the same or separate from that which encodes the genome or antigenome, and various combinations thereof. Additional proteins may be included as desired, encoded by its own vector or by a vector encoding one or more of the N, P, L and other desired PIV proteins, or the complete genome or antigenome.

By providing infectious clones of PIV the invention permits a wide range
25 of alterations to be recombinantly produced within the PIV genome (or antigenome), yielding defined mutations which specify desired phenotypic changes. By "infectious clone" is meant cDNA or its product, synthetic or otherwise, RNA capable of being directly incorporated into infectious virions which can be transcribed into genomic or antigenomic RNA capable of serving as a template to produce the genome of infectious
30 viral or subviral particles. As noted above, defined mutations can be introduced by a variety of conventional techniques (e.g., site-directed mutagenesis) into a cDNA copy of the genome or antigenome. The use of genomic or antigenomic cDNA subfragments to assemble a complete genome or antigenome cDNA as described herein has the advantage that each region can be manipulated separately, where small cDNA subjects provide for

better ease of manipulation than large cDNA subjects, and then readily assembled into a complete cDNA. Thus, the complete antigenome or genome cDNA, or a selected subfragment thereof, can be used as a template for oligonucleotide-directed mutagenesis. This can be through the intermediate of a single-stranded phagemid form, such as using
5 the MUTA-gen® kit of Bio-Rad Laboratories (Richmond, CA), or a method using the double-stranded plasmid directly as a template such as the Chameleon® mutagenesis kit of Strategene (La Jolla, CA), or by the polymerase chain reaction employing either an oligonucleotide primer or a template which contains the mutation(s) of interest. A mutated subfragment can then be assembled into the complete antigenome or genome
10 cDNA. A variety of other mutagenesis techniques are known and can be routinely adapted for use in producing the mutations of interest in a PIV antigenome or genome cDNA of the invention.

Thus, in one illustrative embodiment mutations are introduced by using the MUTA-gen® phagemid in vitro mutagenesis kit available from Bio-Rad Laboratories.
15 In brief, cDNA encoding an PIV genome or antigenome is cloned into the plasmid pTZ18U, and used to transform CJ236 cells (Life Technologies). Phagemid preparations are prepared as recommended by the manufacturer. Oligonucleotides are designed for mutagenesis by introduction of an altered nucleotide at the desired position of the genome or antigenome. The plasmid containing the genetically altered genome or antigenome is
20 then amplified.

Mutations can vary from single nucleotide changes to the introduction, deletion or replacement of large cDNA segments containing one or more genes or genome segments. Genome segments can correspond to structural and/or functional domains, e.g., cytoplasmic, transmembrane or ectodomains of proteins, active sites such
25 as sites that mediate binding or other biochemical interactions with different proteins, epitopic sites, e.g., sites that stimulate antibody binding and/or humoral or cell mediated immune responses, etc. Useful genome segments in this regard range from about 15-35 nucleotides in the case of genome segments encoding small functional domains of proteins, e.g., epitopic sites, to about 50, 75, 100, 200-500, and 500-1,500 or more
30 nucleotides.

The ability to introduce defined mutations into infectious PIV has many applications, including the manipulation of PIV pathogenic and immunogenic mechanisms. For example, the functions of PIV proteins, including the N, P, M, F, HN, and L proteins and C, D and V ORF products, can be manipulated by introducing

mutations which ablate or reduce the level of protein expression, or which yield mutant protein. Various genome RNA structural features, such as promoters, intergenic regions, and transcription signals, can also be routinely manipulated within the methods and compositions of the invention. The effects of trans-acting proteins and cis-acting RNA sequences can be readily determined, for example, using a complete antigenome cDNA in parallel assays employing PIV minigenomes (Dimock et al., *J. Virol.* 67:2772-2778, 1993, incorporated herein by reference in its entirety), whose rescue-dependent status is useful in characterizing those mutants that may be too inhibitory to be recovered in replication-independent infectious virus.

10 Certain substitutions, insertions, deletions or rearrangements of genes or genome segments within recombinant PIV of the invention (e.g., substitutions of a genome segment encoding a selected protein or protein region, for instance a cytoplasmic tail, transmembrane domain or ectodomain, an epitopic site or region, a binding site or region, an active site or region containing an active site, etc.) are made in structural or functional relation to an existing, "counterpart" gene or genome segment from the same or different PIV or other source. Such modifications yield novel recombinants having desired phenotypic changes compared to wild-type or parental PIV or other viral strains. For example, recombinants of this type may express a chimeric protein having a cytoplasmic tail and/or transmembrane domain of one PIV fused to an ectodomain of another PIV. Other exemplary recombinants of this type express duplicate protein regions, such as duplicate immunogenic regions.

25 As used herein, "counterpart" genes, genome segments, proteins or protein regions, are typically from heterologous sources (e.g., from different PIV genes, or representing the same (i.e., homologous or allelic) gene or genome segment in different PIV types or strains). Typical counterparts selected in this context share gross structural features, e.g., each counterpart may encode a comparable protein or protein structural domain, such as a cytoplasmic domain, transmembrane domain, ectodomain, binding site or region, epitopic site or region, etc. Counterpart domains and their encoding genome segments embrace an assemblage of species having a range of size and sequence variations defined by a common biological activity among the domain or genome segment variants.

30 Counterpart genes and genome segments, as well as other polynucleotides disclosed herein for producing recombinant PIV within the invention, often share substantial sequence identity with a selected polynucleotide "reference sequence," e.g.,

with another selected counterpart sequence. As used herein, a “reference sequence” is a defined sequence used as a basis for sequence comparison, for example, a segment of a full-length cDNA or gene, or a complete cDNA or gene sequence. Generally, a reference sequence is at least 20 nucleotides in length, frequently at least 25 nucleotides in length, and often at least 50 nucleotides in length. Since two polynucleotides may each (1) comprise a sequence (i.e., a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and (2) may further comprise a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a “comparison window” to identify and compare local regions of sequence similarity. A “comparison window”, as used herein, refers to a conceptual segment of at least 20 contiguous nucleotide positions wherein a polynucleotide sequence may be compared to a reference sequence of at least 20 contiguous nucleotides and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences.

Optimal alignment of sequences for aligning a comparison window may be conducted by the local homology algorithm of Smith & Waterman (Adv. Appl. Math. 2:482, 1981; incorporated herein by reference), by the homology alignment algorithm of Needleman & Wunsch, (J. Mol. Biol. 48:443, 1970; incorporated herein by reference), by the search for similarity method of Pearson & Lipman, (Proc. Natl. Acad. Sci. USA 85:2444, 1988; incorporated herein by reference), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Dr., Madison, WI, incorporated herein by reference), or by inspection, and the best alignment (i.e., resulting in the highest percentage of sequence similarity over the comparison window) generated by the various methods is selected. The term “sequence identity” means that two polynucleotide sequences are identical (i.e., on a nucleotide-by-nucleotide basis) over the window of comparison. The term “percentage of sequence identity” is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, U, or I) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e.,

the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The terms "substantial identity" as used herein denotes a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 85 percent sequence identity, preferably at least 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison window of at least 20 nucleotide positions, frequently over a window of at least 25-50 nucleotides, wherein the percentage of sequence identity is calculated by comparing the reference sequence to the polynucleotide sequence which may include deletions or additions which total 20 percent or less of the reference sequence over the window of comparison. The reference sequence may be a subset of a larger sequence.

In addition to these polynucleotide sequence relationships, proteins and protein regions encoded by recombinant PIV of the invention are also typically selected to have conservative relationships, i.e. to have substantial sequence identity or sequence similarity, with selected reference polypeptides. As applied to polypeptides, the term "sequence identity" means peptides share identical amino acids at corresponding positions. The term "sequence similarity" means peptides have identical or similar amino acids (i.e., conservative substitutions) at corresponding positions. The term "substantial sequence identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 80 percent sequence identity, preferably at least 90 percent sequence identity, more preferably at least 95 percent sequence identity or more (e.g., 99 percent sequence identity). The term "substantial similarity" means that two peptide sequences share corresponding percentages of sequence similarity. Preferably, residue positions which are not identical differ by conservative amino acid substitutions. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine. Abbreviations for the twenty naturally occurring amino acids used herein

follow conventional usage (Immunology - A Synthesis, 2nd ed., E.S. Golub & D.R. Gren, eds., Sinauer Associates, Sunderland, MA, 1991, incorporated herein by reference).

Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as α , α -disubstituted amino acids, N-alkyl amino acids, lactic acid, and
5 other unconventional amino acids may also be suitable components for polypeptides of the present invention. Examples of unconventional amino acids include: 4-hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, ω -N-methylarginine, and other similar amino acids and imino acids (e.g., 4-
10 hydroxyproline). Moreover, amino acids may be modified by glycosylation, phosphorylation and the like.

To select candidate vaccine viruses according to the invention, the criteria of viability, attenuation and immunogenicity are determined according to well known methods. Viruses which will be most desired in vaccines of the invention must maintain
15 viability, have a stable attenuation phenotype, exhibit replication in an immunized host (albeit at lower levels), and effectively elicit production of an immune response in a vaccinee sufficient to confer protection against serious disease caused by subsequent infection from wild-type virus. The recombinant PIV of the invention are not only viable and more appropriately attenuated than previous vaccine candidates, but are more stable
20 genetically *in vivo*--retaining the ability to stimulate a protective immune response and in some instances to expand the protection afforded by multiple modifications, e.g., induce protection against different viral strains or subgroups, or protection by a different immunologic basis, e.g., secretory versus serum immunoglobulins, cellular immunity, and the like.

25 Recombinant PIV of the invention can be tested in various well known and generally accepted *in vitro* and *in vivo* models to confirm adequate attenuation, resistance to phenotypic reversion, and immunogenicity for vaccine use. In *in vitro* assays, the modified virus (e.g., a multiply attenuated, biologically derived or recombinant PIV) is tested, e.g., level of replication, for temperature sensitivity of virus replication, i.e. ts
30 phenotype, and for the small plaque or other desired phenotype. Modified viruses are further tested in animal models of PIV infection. A variety of animal models have been described and are summarized in various references incorporated herein. PIV model systems, including rodents and non-human primates, for evaluating attenuation and

immunogenic activity of PIV vaccine candidates are widely accepted in the art, and the data obtained therefrom correlate well with PIV infection, attenuation and immunogenicity in humans.

In accordance with the foregoing description, the invention also provides
5 isolated, infectious recombinant PIV viral compositions for vaccine use. The attenuated virus which is a component of a vaccine is in an isolated and typically purified form. By isolated is meant to refer to PIV which is in other than a native environment of a wild-type virus, such as the nasopharynx of an infected individual. More generally, isolated is meant to include the attenuated virus as a component of a cell culture or other artificial
10 medium where it can be propagated and characterized in a controlled setting. For example, attenuated PIV of the invention may be produced by an infected cell culture, separated from the cell culture and added to a stabilizer.

For vaccine use, recombinant PIV produced according to the present invention can be used directly in vaccine formulations, or lyophilized, as desired, using
15 lyophilization protocols well known to the artisan. Lyophilized virus will typically be maintained at about 4°C. When ready for use the lyophilized virus is reconstituted in a stabilizing solution, e.g., saline or comprising SPG, Mg⁺⁺ and HEPES, with or without adjuvant, as further described below.

PIV vaccines of the invention contain as an active ingredient an
20 immunogenically effective amount of PIV produced as described herein. The modified virus may be introduced into a host with a physiologically acceptable carrier and/or adjuvant. Useful carriers are well known in the art, and include, e.g., water, buffered water, 0.4% saline, 0.3% glycine, hyaluronic acid and the like. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being
25 combined with a sterile solution prior to administration, as mentioned above. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan
30 monolaurate, triethanolamine oleate, and the like. Acceptable adjuvants include incomplete Freund's adjuvant, MPLTM (3-o-deacylated monophosphoryl lipid A; RIBI ImmunoChem Research, Inc., Hamilton, MT) and IL-12 (Genetics Institute, Cambridge MA), among many other suitable adjuvants well known in the art.

Upon immunization with a PIV composition as described herein, via aerosol, droplet, oral, topical or other route, the immune system of the host responds to the vaccine by producing antibodies specific for PIV virus proteins, e.g., F and HN glycoproteins. As a result of the vaccination with an immunogenically effective amount of PIV produced as described herein, the host becomes at least partially or completely immune to PIV infection, or resistant to developing moderate or severe PIV infection, particularly of the lower respiratory tract.

The host to which the vaccines are administered can be any mammal which is susceptible to infection by PIV or a closely related virus and which host is capable of generating a protective immune response to the antigens of the vaccinating strain. Accordingly, the invention provides methods for creating vaccines for a variety of human and veterinary uses.

The vaccine compositions containing the PIV of the invention are administered to a host susceptible to or otherwise at risk for PIV infection to enhance the host's own immune response capabilities. Such an amount is defined to be a "immunogenically effective dose." In this use, the precise amount of PIV to be administered within an effective dose will depend on the host's state of health and weight, the mode of administration, the nature of the formulation, etc., but will generally range from about 10^3 to about 10^7 plaque forming units (PFU) or more of virus per host, more commonly from about 10^4 to 10^6 PFU virus per host. In any event, the vaccine formulations should provide a quantity of modified PIV of the invention sufficient to effectively protect the host patient against serious or life-threatening PIV infection.

The PIV produced in accordance with the present invention can be combined with viruses of other PIV serotypes or strains to achieve protection against multiple PIV serotypes or strains. Alternatively, protection against multiple PIV serotypes or strains can be achieved by combining protective epitopes of multiple serotypes or strains engineered into one virus, as described herein. Typically when different viruses are administered they will be in admixture and administered simultaneously, but they may also be administered separately. Immunization with one strain may protect against different strains of the same or different serotype.

In some instances it may be desirable to combine the PIV vaccines of the invention with vaccines which induce protective responses to other agents, particularly other childhood viruses. In another aspect of the invention the PIV can be employed as a vector for protective antigens of other pathogens, such as respiratory syncytial virus

(RSV) or measles virus, by incorporating the sequences encoding those protective antigens into the PIV genome or antigenome which is used to produce infectious PIV, as described herein (see, e.g., U.S. Provisional Patent Application Serial No. 60/170,195, filed December 10, 1999 by Murphy et al., incorporated herein by reference).

5 In all subjects, the precise amount of recombinant PIV vaccine administered, and the timing and repetition of administration, will be determined based on the patient's state of health and weight, the mode of administration, the nature of the formulation, etc. Dosages will generally range from about 10^3 to about 10^7 plaque forming units (PFU) or more of virus per patient, more commonly from about 10^4 to 10^6 10 PFU virus per patient. In any event, the vaccine formulations should provide a quantity of attenuated PIV sufficient to effectively stimulate or induce an anti-PIV immune response, e.g., as can be determined by complement fixation, plaque neutralization, and/or enzyme-linked immunosorbent assay, among other methods. In this regard, individuals are also monitored for signs and symptoms of upper respiratory illness. As 15 with administration to rhesus monkeys, the attenuated virus of the vaccine grows in the nasopharynx of vaccinees at levels approximately 10-fold or more lower than wild-type virus, or approximately 10-fold or more lower when compared to levels of incompletely attenuated PIV.

In neonates and infants, multiple administration may be required to elicit 20 sufficient levels of immunity. Administration should begin within the first month of life, and at intervals throughout childhood, such as at two months, six months, one year and two years, as necessary to maintain sufficient levels of protection against native (wild-type) PIV infection. Similarly, adults who are particularly susceptible to repeated or serious PIV infection, such as, for example, health care workers, day care workers, family 25 members of young children, the elderly, individuals with compromised cardiopulmonary function, may require multiple immunizations to establish and/or maintain protective immune responses. Levels of induced immunity can be monitored by measuring amounts of neutralizing secretory and serum antibodies, and dosages adjusted or vaccinations repeated as necessary to maintain desired levels of protection. Further, different vaccine 30 viruses may be indicated for administration to different recipient groups. For example, an engineered PIV strain expressing a cytokine or an additional protein rich in T cell epitopes may be particularly advantageous for adults rather than for infants.

PIV vaccines produced in accordance with the present invention can be combined with viruses expressing antigens of another subgroup or strain of PIV to

achieve protection against multiple PIV subgroups or strains. Alternatively, the vaccine virus may incorporate protective epitopes of multiple PIV strains or subgroups engineered into one PIV clone, as described herein.

The PIV vaccines of the invention elicit production of an immune response that is protective against serious lower respiratory tract disease, such as pneumonia and bronchiolitis when the individual is subsequently infected with wild-type PIV. While the naturally circulating virus is still capable of causing infection, particularly in the upper respiratory tract, there is a very greatly reduced possibility of rhinitis as a result of the vaccination and possible boosting of resistance by subsequent infection by wild-type virus. Following vaccination, there are detectable levels of host-engendered serum and secretory antibodies which are capable of neutralizing homologous (of the same subgroup) wild-type virus *in vitro* and *in vivo*. In many instances the host antibodies will also neutralize wild-type virus of a different, non-vaccine subgroup.

Preferred PIV vaccine candidates of the invention exhibit a very substantial diminution of virulence when compared to wild-type virus that is circulating naturally in humans. The virus is sufficiently attenuated so that symptoms of infection will not occur in most immunized individuals. In some instances the attenuated virus may still be capable of dissemination to unvaccinated individuals. However, its virulence is sufficiently abrogated such that severe lower respiratory tract infections in the vaccinated or incidental host do not occur.

The level of attenuation of PIV vaccine candidates may be determined by, for example, quantifying the amount of virus present in the respiratory tract of an immunized host and comparing the amount to that produced by wild-type PIV or other attenuated PIV which have been evaluated as candidate vaccine strains. For example, the attenuated virus of the invention will have a greater degree of restriction of replication in the upper respiratory tract of a highly susceptible host, such as a chimpanzee, or rhesus monkey, compared to the levels of replication of wild-type virus, e.g., 10- to 1000-fold less. In order to further reduce the development of rhinorrhea, which is associated with the replication of virus in the upper respiratory tract, an ideal vaccine candidate virus should exhibit a restricted level of replication in both the upper and lower respiratory tract. However, the attenuated viruses of the invention must be sufficiently infectious and immunogenic in humans to confer protection in vaccinated individuals. Methods for determining levels of PIV in the nasopharynx of an infected host are well known in the literature.

Levels of induced immunity provided by the vaccines of the invention can also be monitored by measuring amounts of neutralizing secretory and serum antibodies. Based on these measurements, vaccine dosages can be adjusted or vaccinations repeated as necessary to maintain desired levels of protection. Further, different vaccine viruses may be advantageous for different recipient groups. For example, an engineered PIV strain expressing an additional protein rich in T cell epitopes may be particularly advantageous for adults rather than for infants.

In yet another aspect of the invention the PIV is employed as a vector for transient gene therapy of the respiratory tract. According to this embodiment the recombinant PIV genome or antigenome incorporates a sequence which is capable of encoding a gene product of interest. The gene product of interest is under control of the same or a different promoter from that which controls PIV expression. The infectious PIV produced by coexpressing the recombinant PIV genome or antigenome with the N, P, L and other desired PIV proteins, and containing a sequence encoding the gene product of interest, is administered to a patient. Administration is typically by aerosol, nebulizer, or other topical application to the respiratory tract of the patient being treated. Recombinant PIV is administered in an amount sufficient to result in the expression of therapeutic or prophylactic levels of the desired gene product. Representative gene products which may be administered within this method are preferably suitable for transient expression, including, for example, interleukin-2, interleukin-4, gamma-interferon, GM-CSF, G-CSF, erythropoietin, and other cytokines, glucocerebrosidase, phenylalanine hydroxylase, cystic fibrosis transmembrane conductance regulator (CFTR), hypoxanthine-guanine phosphoribosyl transferase, cytotoxins, tumor suppressor genes, antisense RNAs, and vaccine antigens.

The following examples are provided by way of illustration, not limitation.

EXAMPLE I

Construction of cDNAs Encoding a Chimeric HPIV3/BPIV3 Antigenome and

Recovery of Infectious Virus

The following three examples document studies to identify which of the proteins of BPIV3 contribute to its host range restriction in primates. To illustrate these methods, the N protein of the wild type HPIV3 virus was replaced with its counterpart from BPIV3. This exchange was accomplished using a reverse genetics system for recovery of infectious PIV from cDNA as described above. The studies were initiated

with the N gene of BPIV3 because this protein possesses an intermediate level of amino acid sequence difference from its HPIV3 counterpart compared to other HPIV3 and BPIV3 proteins (see Example I).

A chimeric recombinant virus was constructed in which the N ORF of the JS strain of HPIV3 was replaced by that of either the Ka or SF strain of BPIV3 (Fig. 1). These chimeric viruses possess the HN and F glycoproteins of the HPIV3 parent and will induce a high level of immunity to HPIV3 in primates. Both chimeric viruses were successfully recovered. Both grew to high titer in cell culture and both were found to be attenuated in rhesus monkeys. Thus, the N protein was identified as an exemplary protein that contributes to the host range phenotype of BPIV3. Immunization of rhesus monkeys with either the Ka or SF chimeric recombinant virus induced a high level of resistance to the replication of HPIV3 used as a wild type challenge.

The present invention, therefore, establishes the usefulness of reverse genetics methods to generate chimeric human-bovine PIV virus that combines the host range attenuation properties of BPIV3 and the immunogenicity of the HPIV3 HN and F protective antigens. Immunization of humans with such a chimeric recombinant will redress the problem of suboptimal immunogenicity of the BPIV3 vaccine previously observed in humans.

The complete consensus nucleotide sequence for each of the Ka or SF BPIV3 strains was determined from RT-PCR products generated from virion RNA. These sequences are set forth in Figures 6A-6G, and Figures 7A-7G, respectively. The full length cDNA encoding a complete 15456 nucleotide (nt) antigenomic RNA of BPIV3 Ka is set forth in Figures 6A-6G herein (see also GenBank accession #AF178654). The GenBank sequence for BPIV3 kansas strain differs from the sequence of the exemplary cDNA in two positions at nucleotide 21 and 23. Both, the published sequence and the sequence in the exemplary cDNA occur naturally in kansas strain virus population with similar frequencies. The cDNA used in the present example contains a sequence beginning at nucleotide 18, ACTGGTT (SEQ ID NO. 1), whereas the corresponding published sequence (GenBank accession #AF178654; Figures 6A-6G, SEQ ID NO. 22) reads ACTTGCT (differing nucleotides at positions 21 and 23 are underscored).

To construct consensus nucleotide sequences for the Ka and SF BPIV3 strains, virion RNA was subjected to reverse transcription using the Superscript II Preamplification System (Life Technologies, Gaithersburg, MD) and 200 ng of random hexamer primers. PCR was carried out on the first strand product using the Advantage

cDNA PCR kit (Clontech Laboratories, Palo Alto, CA). Ka and SF genomes were each amplified by PCR in 3 or 4 overlapping fragments using primers homologous to regions of RNA conserved among previously-published paramyxovirus sequences. Each primer pair was constructed to include matching restriction enzyme sites (not represented in the sequence targeted for amplification).

A separate random library was generated for each amplicon by digesting a set of PCR products with the appropriate restriction enzyme, followed by gel-purification, ligation of the products into tandem arrays and sonication. A random library was generated from this pool of sheared cDNA sequences by cloning a subset (approx. 500 bp fragments) into M13. The nucleotide sequences of cDNA inserts were determined by automated DNA sequencing using the Taq DYE Deoxy Terminator cycle sequencing kit (ABI, Foster City, CA). A continuous sequence (contig) was assembled for each of the original large RT-PCR fragments with sufficient redundancy that each nucleotide position was confirmed by a minimum 3 independent M13 clones. The 5' and 3' terminal genomic sequences of Ka and SF were converted to cDNA using the system for Rapid Amplification of cDNA Ends (Life Technologies, Gaithersburg, MD) and sequenced by automated sequencing.

These sequences are set forth in Figures 6A-6G (Ka) and Figures 7A-7G (SF), respectively. Analysis of these sequences revealed that the percent amino acid identity between HPIV3 and BPIV3 for each of the following proteins is: N (86%), P (65%), M (93%), F (83%), HN (77%), and L (91%). Thus sequence divergence was found distributed over many genes. The deduced amino acid sequence of the N genes of these two viruses is presented in GenBank #AF178654 (Ka) and #AF178655 (SF), not included. The position of the N ORF in the BPIV3 genome is indicated in the respective BenBank reports and included herein by reference. In the example below, the N ORF of the Ka or SF virus was initially selected for replacement of the corresponding gene in the HPIV3 virus because the N gene represents a gene with an intermediate level of sequence divergence among the six HPIV3 and BPIV3 proteins. In this study the N ORF, but not the 3' or 5' noncoding N gene sequences, was exchanged, which permitted us to assign an observed attenuation phenotype of cKa and cSF to the protein encoded by the N gene.

Human-bovine chimeric full-length PIV3 genomes were constructed by introducing the BPIV3 Ka or SF N coding region as a replacement for its HPIV3 counterpart into the rJS cDNA p3/7(131)2G which encodes a complete copy of HPIV3 positive-sense antigenomic RNA (see, e.g., Durbin et al., 1997a, *supra*; Hoffman et al.,

WO 01/04320

- 1997, *supra*; Skiadopoulos et al., 1998, *supra*; U.S. Patent Application Serial No. 09/083,793, filed May 22, 1998; U.S. Provisional Application No. 60/047,575, filed May 23, 1997 (corresponding to International Publication No. WO 98/53078), and U.S. Provisional Application No. 60/059,385, filed September 19, 1997; each incorporated herein by reference). BPIV3 and HPIV3 N coding regions with flanking sequences were first subcloned and further modified to permit an exchange of just the N ORF. pUC119JSN bearing the HPIV3 N gene and the plasmids with a BPIV3 N Ka or SF gene (pBSKaN and pBSSFN) were subjected to mutagenesis using the method of Kunkel (Proc. Natl. Acad. Sci. USA 82:488-492, 1985, incorporated herein by reference) to introduce NcoI and AflIII restriction enzyme recognition sites at translational start and stop sites, respectively (Fig. 1A). Following NcoI/AflIII digestion of pUC119KaN-NcoI/AflIII, the BPIV3 N coding region was introduced as an NcoI/AflIII fragment into pUC119JSN-NcoI/AflIII as a replacement for the HPIV3 N coding region (Fig. 1B). The chimeric N genes, which contain the HPIV3 3' and 5' noncoding sequences and the BPIV3 ORF, were modified by site-directed mutagenesis to restore the original HPIV3 noncoding sequence and BPIV3 coding sequence. This chimeric N gene was then introduced into the 5' half of the rJS antigenome, pLeft, in exchange for its corresponding HPIV3 sequence (Figs. 2A and 2B) using existing MluI and EcoRI sites present in the human sequence. In each case parallel reactions were carried out for the SF N ORF. The chimeric pLeft plasmid was combined with the XhoI/NgoMl fragment from pRight containing the 3' half of the rJS antigenome flanked by the delta ribozyme and the T7 terminator at its 3' end (Fig. 2). The resulting chimeric PIV3 plasmids designated pB/HPIV3NKa or pB/HPIV3NSF, contained the full-length rJS antigenome in which the N ORF encoded the BPIV3 Ka or SF N protein.
- Chimeric antigenomic HPIV3/BPIV3 cDNAs were transfected individually into HEp-2 cells grown to near-confluence in 6-well plates along with two previously-described support plasmids, pTM(P no C) and pTM(L), Lipofectace (Life Technologies, Gaithersburg, MD), and infected with a modified vaccinia virus recombinant that expresses bacteriophage T7 RNA polymerase (MVA-T7) as previously described (Durbin et al., Virology 234:74-83, 1997b). An N support plasmid used in previous work was omitted because the antigenomic plasmid expressed sufficient levels of the N protein. The cultures were maintained for 3.5 days at 32°C after which supernatants were harvested, passaged in LLC-MK2 cells and plaque-purified 3 times in

WO 01/04320

LLC-MK2 cells. The identities of the chimeric viruses incorporating a human PIV 3 background genome or antigenome and a BPIV3 N protein (designated as rHPIV3-N_B chimeric recombinants or, more specifically, as "cKa" and "cSF" chimeric viruses) recovered from the transfections were confirmed by sequencing RT-PCR products containing the regions of the N ORF start and stop codons from virion RNA isolated after amplification of triply plaque-purified virus (Fig. 3). This amplified product and the corresponding amplified HPIV3 rJS and BPIV3 Ka or SF sequences were also subjected to TaqI digestion to confirm the chimeric identity of cKa and cSF viruses (Fig. 4). TaqI digestion profiles were distinct for the 3 parental and 2 chimeric viruses, and each parental profile included TaqI fragments of unique size, allowing the contribution of sequence of rJS, Ka and SF parents to the chimeric viruses to be verified. The recovered cKa and cSF chimeric recombinants each contained the expected sequences as designed.

EXAMPLE II

Replication of HPIV3/BPIV3 Chimeric Viruses in Cell Culture

Efficient replication of live attenuated virus vaccines in tissue culture cells is a feature of human-bovine chimeric PIV of the invention that permits efficient manufacture of the recombinant vaccine materials. The multicycle replication of rJS parent, cKa, Ka parent, cSF, and SF parent in a bovine cell line (MDBK) and in a simian cell line (LLC-MK2) was determined by infecting cells with virus at a multiplicity of infection of 0.01 and harvesting samples (in triplicate) over a five day period of time (Fig. 5) as previously described (Tao et al., 1998, *supra*, incorporated herein by reference). The chimeric viruses replicated efficiently in both cell lines like their human or bovine parent viruses without significant delay in replication or a significant reduction in the titer of virus achieved. In each case, the chimeric viruses replicated to over $10^{7.0}$ TCID₅₀/ml which is well above the $10^{4.0}$ or $10^{5.0}$ dose of live attenuated human or bovine PIV vaccines currently being used in human clinical trials (Karron et al., 1996, *supra*; Karron et al., 1995a, *supra*; and Karron et al., 1995b, *supra*).

EXAMPLE III

Evaluation of Attenuation and Protective Efficacy of the HPIV3/BPIV3 Chimeric Viruses in Rhesus Monkeys

Both the SF and Ka BPIV3s are attenuated for the upper and the lower respiratory tract of the rhesus monkey (van Wyke-Coelingh et al., 1988, *supra*). This

attenuation phenotype correlates with attenuation in humans (Karron et al., 1995a, *supra*) as indicated by the fact that Ka is highly restricted in replication in the upper respiratory tract of fully susceptible seronegative infants and children. The absence of cough, croup, bronchiolitis, or pneumonia in the BPIV3-infected vaccinees suggests that the Ka BPIV3 virus is attenuated for the lower respiratory tract as well. Therefore, the rhesus monkey is widely accepted as a reasonably correlative model to evaluate attenuation of candidate PIV vaccine viruses and their efficacy against challenge with wild type PIV.

The rJS, cKa, Ka parent, cSF, and SF parent were administered intranasally and intratracheally at a dose of $10^{5.0}$ TCID₅₀ per site to rhesus monkeys. Replication was monitored using previously described procedures for obtaining samples from the upper (nasopharyngeal swab specimens) and lower (tracheal lavage specimens) respiratory tract and for titering the virus in LLC-MK2 cells (Hall et al., 1992, *supra*). The cKa and cSF recombinants were significantly attenuated for the upper respiratory tract (Table 1) exhibiting, respectively, a 63-fold or a 32-fold reduction in mean peak virus titer compared to that of the rJS HPIV3 parent. Both cKa and cSF were also attenuated for the lower respiratory tract, but this difference was only statistically significant for cSF. The low level of replication of rJS in the lower respiratory tract made it difficult to demonstrate in a statistically-significant fashion further restriction of replication due to an attenuation phenotype at this site.

Table 1. Replication of cKa and cSFis restricted relative to HPIV3 in the upper and lower respiratory tracts of rhesus monkeys.

		Virus Replication						
Immu- nizing virus ¹	No. of ani- mals	Mean titers				Mean peak titers		
		Log ₁₀ TCID ₅₀ /ml±standard error [Duncan grouping] ²				Log ₁₀ TCID ₅₀ /ml±standard error [Duncan grouping]		
		Nasopharynx		Trachea		Nasopharynx	Trachea	
		day 6	day 7	day 4	day 6			
rJS	4	5.3±0.59[A]	3.9±0.36[A]	1.7±0.45[A]	1.7±0.29[A]	5.3±0.59[A]	2.5±0.51[A]	
cKa	4	3.0±0.58[B]	2.9±0.42[AB]	1.5±0.40[A]	1.0±0.19[A]	3.5±0.54[B]	1.5±0.18[AB]	
Ka	4	2.0±0.27[B]	2.4±0.30[B]	1.3±0.26[A]	1.3±0.21[A]	2.5±0.30[B]	1.6±0.15[AB]	
cSF	4	3.3±0.40[B]	3.7±0.57[A]	1.1±0.25[A]	1.1±0.24[A]	3.8±0.46[B]	1.4±0.26[B]	
SF	4	2.8±0.48[B]	2.6±0.40[AB]	1.6±0.46[A]	1.5±0.40[A]	3.3±0.28[B]	1.8±0.41[AB]	

¹ Monkeys were inoculated intranasally and intratracheally with 10^{3.0} TCID₅₀ in 1 ml at each site.² Mean viral titers in each column were assigned to statistically similar groups (designated with a letter) using a Duncan Multiple Range test ($\alpha = 0.05$). Mean titers in each column with different letters are statistically different.

The level of replication of each chimeric virus, cKa and cSF, was not significantly different from its bovine parent in the upper or the lower respiratory tract, although the chimeric viruses each replicated somewhat better than their BPIV3 parents in the upper respiratory tract. Thus, the acquisition of the N gene of either the Ka or SF BPIV3 by rJS HPIV3 attenuated the human virus for rhesus monkeys to a level approximately equivalent to that of the BPIV parent. Since the HPIV3/BPIV3 chimeric recombinants replicated efficiently in tissue culture cells *in vitro*, it is clear that the phenotype of host range restricted replication manifested by the two bovine parental viruses was transferred to HPIV3 by the N ORF. It is possible, but unknown and unpredictable, that substitution of other BPIV3 genes, such as M, P, or L, for their HPIV3 counterpart in rJS will give similar or greater levels of attenuation as observed upon substitution of the BPIV3 N gene for the HPIV3 N gene. The observation that the level of replication of cKa and cSF is slightly greater than that of their BPIV3 parents in the upper respiratory tract suggests that additional bovine genes contribute to the host range attenuation phenotype of the BPIV3 parent virus at this site.

Uninoculated monkeys and monkeys that were previously infected with a human or bovine PIV3 parental virus, or with the cKa or cSF chimeric virus, were challenged 42 days after the initial inoculation with $10^{6.0}$ TCID₅₀ of rJS intranasally and intratracheally in a 1 ml inoculum at each site. The nasopharynx and the trachea were sampled as described previously on the days indicated in Table 2. The titer of virus present at each site was determined for each monkey on LLC-MK2 cell monolayers, and the titers presented are mean peak titers (Hall et al., 1992, *supra*). Previous infection with either chimeric virus induced a high level of resistance to replication of the rJS challenge virus in both the upper and lower respiratory tract. Monkeys previously infected with cKa manifested a 300-fold reduction of replication of wild type HPIV3 (rJS) in the upper respiratory tract and a 1000-fold reduction in the lower tract compared to uninoculated control monkeys. Monkeys previously infected with cSF manifested a 2000-fold reduction of replication of rJS in the upper respiratory tract and a 1000-fold reduction in the lower tract compared to uninoculated control monkeys. The level of reduction of replication of rJS challenge virus in monkeys previously-inoculated with cKa or cSF was comparable to that of monkeys previously infected with either the bovine or the human PIV parent. Thus, infection with either HPIV3/BPIV3 chimeric virus provided a high level of protection in the upper and lower respiratory tract of monkeys, and both chimeric viruses represent promising vaccine candidates.

Serum collected from monkeys on days 0 and 28 was tested by HAI assay using HPIV3 (JS strain) and BPIV3 (Ka strain) as antigen as previously described (Coelingh et al., *J. Infect. Dis.* 157:655-662, 1988). Although cKa-N and cSF-N were highly attenuated in the upper and lower respiratory tract of rhesus monkeys relative to rJS, each chimeric virus induced a hemagglutination-inhibiting (HAI) antibody response to HPIV3 that was 2.5 to 5-fold greater in magnitude than that induced by immunization with its respective BPIV3 patent. This likely is due to the presence of HPIV3 HN protein in the chimeric viruses. Furthermore, the HPIV3-specific HAI-responses induced by the chimeric viruses were statistically indistinguishable from that induced by immunization with rJS. An additional unexpected result demonstrated herein is that, following challenge of the monkeys with HPIV3, the level of HAI antibody in monkeys initially immunized with cKa-N or cSF-N was significantly greater than levels observed in animals immunized with rJS, Ka or SF.

EXAMPLE IVConstruction and Characterization of Chimeric HPIV3/BPIV3 Vaccine Candidates Having Heterologous Fusion And Hemagglutinin-Neuraminidase Glycoproteins

5 In the preceding example, the basis of host range restriction of replication of BPIV3 for the respiratory tract of primates was examined by the generation and characterization of a recombinant human PIV3 (rHPIV3) in which the N open reading frame (ORF) was replaced by that of its BPIV3 counterpart. The resulting chimeric virus, rHPIV3-N_B, also referred to as cKa or cSF, efficiently replicated in vitro but was
 10 restricted in replication in the upper respiratory tract of rhesus monkeys, identifying the N protein as an independent determinant of the host range restriction of BPIV3 in rhesus monkeys (Bailly et al., J. Virol. 74:3188-3195, 2000).

In the present example, the contribution of the fusion (F) and hemagglutinin-neuraminidase (HN) glycoprotein genes of bovine parainfluenza virus type
 15 3 (BPIV3) to its restricted replication in the respiratory tract of non-human primates was examined by generating and characterizing two reciprocal chimeric BPIV3/HPIV3 viruses. A chimeric HPIV3 containing heterologous BPIV3 F and HN glycoprotein genes in place of its own, and the reciprocal recombinant comprising a BPIV3 "backbone" bearing the HPIV3 F and HN genes substituted for the counterpart BPIV3 glycoprotein
 20 genes, were generated to assess the effect of glycoprotein substitution on replication of HPIV3 and BPIV3 in the upper and lower respiratory tract of rhesus monkeys. Thus, in one chimeric virus, the F and HN genes of HPIV3 were replaced with their BPIV3 counterparts, resulting in a chimeric recombinant designated rHPIV3-F_BHN_B. The reciprocal chimeric recombinant PIV3 (rBPIV3-F_HHN_H) was constructed by replacing the
 25 F and HN genes of a recombinant BPIV3 (rBPIV3) with their HPIV3 counterparts. In the latter virus, the introduction of the HPIV3 F and HN ORFs into the BPIV3 backbone combines the antigenic determinants of HPIV3 with the backbone of BPIV3 and thus provides an improved vaccine candidate compared with parental BPIV3. The F and HN genes were exchanged as pairs in view of the proposed requirement for homologous HN
 30 and F proteins for parainfluenza viruses for full functional activity (Deng et al., Virology 209:457-469, 1995; and Tanabayashi et al., J. Virol. 70:6112-6118, 1996; each incorporated herein by reference).

The foregoing chimeric viruses were readily recovered and exhibited kinetics of replication in simian LLC-MK2 cells that were comparable to those of their

parent viruses, suggesting that the heterologous glycoproteins were compatible with the PIV3 internal proteins. The distinctive features of cytopathology of BPIV3 versus HPIV3 cosegregated with their respective F and HN genes. HPIV3 bearing the BPIV3 F and HN genes was attenuated for replication in rhesus monkeys to a level similar to that of its
5 BPIV3 parent virus, indicating that the glycoprotein genes of BPIV3 are major determinants of its host range restriction of replication in rhesus monkeys. BPIV3 bearing the HPIV3 F and HN genes (rBPIV3-F_HHN_H) replicated in rhesus monkeys to a level intermediate between that of HPIV3 and BPIV3.

These results indicate that the F and HN genes make a significant
10 contribution to the overall attenuation of BPIV3. Furthermore, they demonstrate that BPIV3 sequences outside the F and HN region also contribute to the attenuation phenotype in primates. This latter finding is consistent with the demonstration in the preceding example that the nucleoprotein coding sequence of BPIV3 is a determinant of its attenuation for primates. Despite its restricted replication in the respiratory tract of
15 rhesus monkeys, rBPIV3-F_HHN_H conferred a level of protection against challenge with wild type HPIV3 that was indistinguishable from that conferred by previous infection with wild type HPIV3. From these and related findings, the usefulness of rBPIV3-F_HHN_H as a vaccine candidate against HPIV3 is readily apparent.

20 Viruses and cells

HEp-2 and simian LLC-MK2 monolayer cell cultures were maintained in MEM medium (Life Technologies, Gaithersburg, MD) supplemented with 5% fetal bovine serum (Summit Biotechnology, Ft. Collins, CO), 50ug/ml gentamicin sulfate, and 4mM glutamine (Life Technologies, Gaithersburg, MD).

25 The wild type BPIV3 strain Kansas/15626/84 (Clone 5-2-4, Lot BPI3-1) (BPIV3 Ka), the HPIV3 JS wild type, its recombinant version (rHPIV3), and the rHPIV3 virus containing the BPIV3 Ka N ORF in place of the HPIV3-N ORF (rHPIV3-N_B) are each described above (see also, Clements et al., 1991, *supra*; Karron et al., 1995a, *supra*; Bailly et al., 2000, *supra*; and Durbin et al., 1997, *supra*). PIVs were propagated at 32°C
30 in LLC-MK2 cells (ATCC CCL-7), as previously described (Hall et al., 1992, *supra*). The modified vaccinia strain Ankara (MVA) recombinant virus that expresses bacteriophage T7 RNA polymerase is described by Wyatt et al. (1995, *supra*).

Construction of antigenomic cDNAs encoding recombinant BPIV3/HPIV3 viruses.**a) Construction of cDNA to recover rBPIV3**

A full length cDNA was constructed to encode the complete 15456 nucleotide (nt) antigenomic RNA of BPIV3 Ka, as described above. The cDNA was assembled from 4 subclones derived from reverse transcription (RT) of viral RNA using the SuperScript II Pre-amplification System (Life Technologies, Gaithersburg, MD) and polymerase chain reaction (PCR) amplification with a High Fidelity PCR kit (Clontech Laboratories, Palo Alto, CA). The RT-PCR products were cloned into modified pUC19 plasmids (New England Biolabs, Beverly, MA) using the following naturally occurring internal restriction enzyme recognition sites: *Sma* I (BPIV3 Ka sequence position nt186), *Pst* I (nt 2896), *Mlu* I (nt 6192), *Sac* II (nt 10452) and *Bsp* LU11 (nt 15412). Multiple subclones of the antigenomic cDNA were sequenced using a Perkin Elmer ABI 310 sequencer with dRhodamine Terminator Cycle Sequencing (Perkin Elmer Applied Biosystems, Warrington, UK), and only those matching the consensus sequence of BPIV3 Ka were used for assembly of the full length clone. The 3' and 5' ends of BPIV3 Ka were cloned and the assembly of the full length cDNA took place in the previously described p(Right) vector (Durbin et al., 1997, *supra*), which we modified to contain a new polylinker with restriction enzyme recognition sites for *Xho* I, *Sma* I, *Mlu* I, *Sac* II, *Eco* RI, *Hind* III and *Rsr*II. The full length cDNA clone pBPIV3(184) contained the following elements in 3' to 5' order: a T7 promoter followed by 2 non-viral guanosine residues, the complete antigenomic sequence of BPIV3 Ka, a hepatitis delta virus ribozyme and a T7 polymerase transcription terminator (Bailly et al., 2000, *supra*; and Durbin et al., 1997a, *supra*).

25 b) Construction of rHPIV3-F_BHN_B and rBPIV3-F_HHN_H

Unique restriction enzyme recognition sites were introduced into the BPIV3 antigenomic cDNA and into the previously described HPIV3 antigenomic cDNA p3/7(131)2G (Durbin et al., 1997a, *supra*) to facilitate the exchange of the F and HN genes between BPIV3 and HPIV3 cDNAs. Using the transformer site-directed mutagenesis protocol from Clontech (Clontech Laboratories, Palo Alto, CA), *Sgr*AI restriction sites were introduced in the downstream non-coding region of the M gene at position 4811 of the rBPIV3 sequence and position 4835 of the rHPIV3 JS sequence (GenBank accession # Z11575). The nucleotide number given for the position of restriction enzyme recognition sites indicates the nucleotide after which the enzyme cuts,

not the first nucleotide of the restriction enzyme recognition site. The sequence was changed from TCCAACATTGCA (SEQ. ID. NO. 2) to TCCACCGGTGCA (SEQ. ID. NO. 3) in rBPIV3 and from CCGACGTATCTA (SEQ. ID. NO. 4) to CGCACCGGTGTA (SEQ. ID. NO. 5) in rHPIV3 (recognition sites underlined). *Bsi*WI restriction sites were introduced in the downstream non-coding region of the HN gene at nt 8595 of the rBPIV3 sequence and at nt 8601 of the rHPIV3 JS sequence. The sequence was changed from GATATAAAGA (SEQ. ID. NO. 6) to GACGTACGGA (SEQ. ID. NO. 7) in rBPIV3 to give pBPIVs(107) and from GACAAAAGGG (SEQ. ID. NO. 8) to GACGTACGGG (SEQ. ID. NO. 9) in rHPIV3 to give pHPIVs(106). The F and HN genes were exchanged between pBPIVs(107) and pHPIVs(106) by digestion of each with *Sgr*AI and *Bsi*WI, gel purification of the fragments, and assembly of the appropriate fragments into the two full length cDNAs. The HPIV3 backbone bearing the BPIV3 F and HN genes, designated pHPIV(215), encoded 15480 nts of viral sequence, of which nts 4835 to 8619 came from BPIV3, and it was used to derive rHPIV3-F_BHN_B (Figures 8A-8C). The BPIV3 backbone bearing the HPIV3 F and HN genes, designated pBPIV(215), encoded 15438 nts of viral sequence, of which nts 4811 to 8577 came from HPIV3, and it was used to derive rBPIV3-F_HHN_H (Figures 8A-8C).

BPIV3 support plasmids for recovery of virus from cDNA.

Support plasmids encoding the BPIV3 Ka N, P and L genes were assembled in modified pUC19 vectors and then cloned into the previously described pTM-1 vector (Durbin et al., 1997a, *supra*). In order to place the individual genes immediately downstream of the T7 promoter in the pTM vector, an *Nco* I site was introduced at the start codon of the N, P and L open reading frames (ORFs) using site-directed mutagenesis. The *Nco* I restriction site and a naturally occurring restriction site downstream of each ORF (*Spe* I for N, *Hinc*II for P and *Bsp* LU11I for L) was used for cloning into pTM. After cloning, the *Nco* I site in pTM(N) was mutagenized back to the original sequence to restore the correct amino acid assignment in the second codon. In pTM(P) and pTM(L) the amino acid sequence encoded by the ORF was not altered by the introduction of *Nco* I sites.

Transfection.

HEp-2 cells (approximately 1.5×10^6 cells per well of a six-well plate) were grown to 90% confluence and transfected with 0.2 μ g each of the BPIV3 support

plasmids pTM(N) and pTM(P), and 0.1 µg of pTM(L), along with 5 µg of the full length antigenomic cDNA and 12µl LipofectACE (Life Technologies, Gaithersburg, MD). Each transfection mixture also contained 1.5×10^7 plaque forming units (PFU) of MVA-T7, as previously described (Durbin et al., 1997, *supra*). The cultures were incubated at 32°C for 12 hrs before the medium was replaced with MEM (Life Technologies, Gaithersburg, MD) containing 10% fetal bovine serum. The supernatants were harvested after incubation at 32°C for an additional three days, and were passaged onto LLC-MK2 cell monolayers in 25cm² flasks and incubated for 5 days at 32°C. Virus present in the supernatant was plaque-purified three times prior to amplification and characterization.

10

Molecular characterization of recovered chimeric recombinants.

The presence of the heterologous F and HN genes in the bovine or human PIV3 backbone was confirmed in plaque-purified recombinant viruses by RT-PCR of viral RNA isolated from infected cells or supernatant, which was performed using a primer pair that recognizes conserved sequences in rBPIV3 and rHPIV3. This yielded similarly sized fragments (nts 4206-9035 in rBPIV3, nts 4224-9041 in rHPIV3, nts 4206-9017 in rBPIV3-F_HHN_H, and nts 4224-9059 in rHPIV3-F_BHN_B) which were then digested with *Eco* RI and analyzed by electrophoresis on a 1% agarose gel (Fig. 9). The nucleotide sequence flanking the introduced *Sgr*AI and *Bsi*WI restriction sites in each virus was confirmed by sequencing the respective RT-PCR product.

20

Replication of HPIV3/BPIV3 chimeric viruses in cell culture.

The multicycle growth kinetics of BPIV3 Ka, rHPIV3-F_BHN_B, rBPIV3-F_HHN_H, rHPIV3-N_B and rHPIV3 in LLC-MK2 cells were determined by infecting cells in triplicate at a multiplicity of infection (MOI) of 0.01 and harvesting samples at 24 hr intervals over a six day period, as previously described (Tao et al., 1998, *supra*). Samples were flash-frozen and titered in a single assay on LLC-MK2 cell monolayers in 96 well plates at 32°C, as described (Durbin et al., *Virology* 261:319-330, 1999b, incorporated herein by reference).

25

Primate Model Studies.

Rhesus monkeys seronegative for PIV3 as determined by hemagglutination-inhibition (HAI) assay (van Wyke Coelingh et al., 1988, *supra*) were

inoculated intranasally and intratracheally in groups of 2 or 4 animals with 10^5 tissue culture infectious dose₅₀ (TCID₅₀) per ml of BPIV3 Ka, rHPIV3-F_BHN_B, rBPIV3-F_HHN_H, rHPIV3-N_B or rHPIV3. Nasopharyngeal swabs were collected daily on days 1 to 11 and on day 13. Tracheal lavage samples were collected on days 2, 4, 6, 8, and 10 post-infection. Individual samples were flash-frozen and stored at -70°C until all samples were available for titration. Virus in the specimens was titered on LLC-MK2 cell monolayers in 24 and 96 well plates as previously described (Durbin et al., 1999b, *supra*). Sera collected from monkeys on days 0 and 28 was tested by HAI assay using HPIV3 JS and BPIV3 Ka as antigens, as previously described (van Wyke Coelingh et al., 1988, *supra*). On day 28 post inoculation, the monkeys were challenged intranasally and intratracheally with 10^6 TCID₅₀ per site of HPIV3 JS. Nasopharyngeal swab samples were collected on days 3, 4, 5, 6, 7 and 8, and tracheal lavage samples on days 4, 6 and 8 post challenge. Samples were titered in a single assay as described above. Serum was collected on day 28 post challenge.

15

Recovery of rBPIV3 and BPIV3/HPIV3 chimeric viruses (rHPIV3-F_BHN_B and rBPIV3-F_HHN_H) from cDNA.

A complete BPIV3 antigenomic cDNA, designated pBPIV(184), was constructed to encode the consensus sequence of BPIV3 Ka. This BPIV3 antigenomic cDNA was further modified by the introduction of unique *SgrAI* and *BsiWI* sites into the downstream noncoding region of the M and HN genes, respectively (Fig. 8C). The same restriction sites were introduced into the downstream noncoding region of the M and HN genes of a previously described complete HPIV3 antigenomic cDNA, p3/7(131)2G (Durbin et al., 1997a, *supra*). The F and HN glycoprotein genes of HPIV3 and BPIV3 were swapped by exchanging this *SgrAI*-*BsiWI* restriction fragment. A direct exchange of entire genes was anticipated to be well-tolerated because of the high level of sequence conservation between the cis-acting signals of BPIV3 and HPIV3. The HPIV3 antigenomic cDNA bearing the BPIV3 F and HN genes was designated pHPIV(215), and the BPIV3 antigenomic cDNA bearing the HPIV3 F and HN genes was designated pBPIV(215).

The antigenomic cDNAs pBPIV(184), pHPIV(215), pBPIV(215) and p3/7(131)2G were separately transfected into HEp-2 cells along with the three BPIV3 support plasmids pTM(N), pTM(P) and pTM(L), and the cells were simultaneously

infected with recombinant MVA expressing the T7 RNA polymerase. To confirm that the recovered viruses indeed were the expected rBPIV3, rHPIV3-F_BHN_B, rBPIV3-F_HHN_H and rHPIV3 viruses, intracellular RNA or RNA from supernatant from each cloned virus was analyzed by RT-PCR using a primer pair that recognized identical sequences in HPIV3 JS and BPIV3 Ka. The primer pair amplified a 4.8 kb fragment of DNA corresponding to the downstream end of the M gene, the F and HN genes, and the upstream end of the L gene (nts 4206-9035 in rBPIV3, nts 4224-9041 in rHPIV3, nts 4206-9017 in rBPIV3-F_HHN_H, and nts 4224-9059 in rHPIV3-F_BHN_B). The generation of each PCR product was dependent upon the inclusion of reverse transcriptase, indicating that each was derived from viral RNA and not from contaminating cDNA (data not shown). The PCR products were then digested with *Eco* R1, which would be predicted to yield a different, unique restriction enzyme digest pattern for each of the four viruses (Figure 9). In each case, the predicted pattern was observed, confirming the identity of the backbone and the inserted F and HN genes. In addition, nucleotide sequencing was performed on the RT-PCR products to confirm the presence of the introduced restriction sites and flanking sequences.

The cytopathic effect (CPE) caused by rBPIV3-F_HHN_H in LLC-MK2 cells was indistinguishable from that of HPIV3 JS (condensed, rounded-up cells and small syncytia) but different from BPIV3 (large multicellular syncytia), whereas the CPE caused by rHPIV3-F_BHN_B was identical to that caused by the BPIV3. This indicates that the cytopathology of the chimeric PIVs cosegregated with the parental origin of the F and HN genes.

BPIV3/HPIV3 chimeric viruses replicate efficiently in cell culture.

The growth kinetics of rHPIV3-F_BHN_B and rBPIV3-F_HHN_H were compared with that of their parental viruses by infecting LLC-MK2 monolayers at an MOI of 0.01 and monitoring the production of infectious virus. The kinetics and magnitude of replication of the two chimeric viruses were comparable to those of their HPIV3 or BPIV3 parental viruses (Figure 10). This suggested that BPIV3 and HPIV3 glycoproteins were compatible with the heterologous PIV3 internal proteins. This is an important property because it will be possible to efficiently prepare vaccine virus.

The F and HN genes of the BPIV3/HPIV3 chimeric viruses are determinants of the host range restriction of replication of BPIV3 Ka in the respiratory tract of rhesus monkeys.

5 rHPIV3-F_BHN_B and rBPIV3-F_HHN_H were evaluated for their ability to replicate in the upper and lower respiratory tract of rhesus monkeys. In particular, the effects of introduction of the BPIV3 F and HN genes into HPIV3 on attenuation of replication in rhesus monkeys was demonstrated, as described above for the BPIV3 N protein (see also, Bailly et al., 2000, *supra*). In addition, the effects of introduction of the HPIV3 F and HN genes into BPIV3 on replication in rhesus monkeys was determined. If 10 the predominant attenuating mutations of BPIV3 were in genes other than the F and HN, then one would expect little overall effect of the HPIV3-BPIV3 glycoprotein exchange on replication of BPIV3 in rhesus monkeys.

Each chimeric virus was administered intranasally and intratracheally to rhesus monkeys at a dose of 10⁵ TCID₅₀ per site. The level of replication of the chimeric 15 viruses was compared to that of the rHPIV3 and BPIV3 parental viruses and to that of rHPIV3-N_B (Table 3). Since the rHPIV3 parental virus replicated to a low to moderate level in the lower respiratory tract, meaningful comparisons between groups could only be made for replication in the upper respiratory tract. The level of replication of rHPIV3-F_BHN_B was similar to that of its BPIV3 parent and substantially lower than that of its 20 HPIV3 parent (Table 3; Figure 11, panel A). This showed that the BPIV3 glycoprotein genes contained one or more major determinants of the host range attenuation phenotype of BPIV3 for rhesus monkeys. The magnitude and pattern of replication of rHPIV3-F_BHN_B and rHPIV3-N_B were very similar, indicating that each of the two bovine genetic elements, namely the N gene versus the F and HN genes, attenuate HPIV3 to a similar 25 extent.

Table 3. The F and HN glycoprotein genes of BPIV3 contribute to its restricted replication in the respiratory tract of rhesus monkeys.

Immunizing virus ¹	Number of animals ²	Mean peak virus titer ³		Serum HAI antibody titer		Serum HAI antibody titer (mean recip. log ₂ ± S.E.) for BPIV3 on day 28 ⁷
		NP swab ⁵	(log ₁₀ TCID ₅₀ /ml ± S.E.) [Duncan Grouping] ⁴	(mean recip. log ₂ ± S.E.) for HPIV3 on day 28 ⁷		
rHPIV3	6	4.7±0.54 [A]	2.4±0.37 [A]	9.5±0.72 [A]		6.8±1.03 [B]
rBPIV3-F _H HN _H	4	3.1±0.58 [B]	1.6±0.05 [A]	6.8±0.63 [BC]		3.8±0.63 [C]
rHPIV3-N _B	6	3.0±0.60 [B]	1.4±0.19 [A]	8.2±0.48 [AB]		6.5±0.62 [B]
rHPIV3-F _B HN _B	4	2.9±0.28 [B]	2.0±0.24 [A]	4.5±0.29 [D]		9.5±0.65 [A]
BPIV3 Ka	6	2.6±0.26 [B]	1.6±0.10 [A]	5.5±0.62 [CD]		9.2±0.60 [A]

88

¹ Monkeys were inoculated intranasally and intratracheally with 10⁵ TCID₅₀ of virus in a 1 ml inoculum at each site.

² The groups with 6 animals contain 4 animals each from a previous rhesus study (Bailey et al., 2000, *supra*).

³ Mean of the peak virus titers for each animal in its group irrespective of sampling day. S.E. = standard error.

⁴ Virus titrations were performed on LLC-MK2 cells at 32°C. The limit of detectability of virus titer was 10 TCID₅₀/ml. Mean viral titers were compared using a Duncan Multiple Range test ($\alpha = 0.05$). Within each column, mean titers with different letters are statistically different. Titers indicated with two letters are not significantly different from those indicated with either letter.

⁵ Nasopharyngeal swab samples were collected on days 1 to 11 and on day 13.

⁶ Tracheal lavage samples were collected on days 2, 4, 6, 8 and 10 post-infection.

⁷ The titers on day 0 were <2.0. Day 28 was the day of challenge with wild type HPIV3.

**Two of the animals in the rHPIV3 group were infected with rHPIV3s, the virus containing two restriction enzyme recognition sites for the glycoprotein swap.

The rBPIV3-F_HHN_H chimeric virus replicated significantly less well than rHPIV3 (Table 3), and it grouped with BPIV3 in a Duncan multiple range test. However, inspection of its pattern of replication in Figure 11B suggested that rBPIV3-F_HHN_H replicated to a level intermediate between that of its HPIV3 and BPIV3 parents. The interpretation that rBPIV3-F_HHN_H replicates to a level intermediate between that of its parents is supported by Friedman's test of consistency of ranks (Sprent, P., "A Generalization Of The Sign Test," Applied Nonparametric Statistical Methods, pp. 123-126, Chapman and Hall, London, 1989, incorporated herein by reference), which indicated that the median titers of HPIV3, rBPIV3-F_HHN_H, and BPIV3 between day 3 and day 8 post infection are significantly different (d.f.2,8; p<0.05). The observation that the introduction of the HPIV3 F and HN proteins resulted in an increase in the replication of BPIV3 in rhesus monkeys indicates (i) that F and HN contain one or more determinants of host range restriction and (ii) that one or more genetic elements of BPIV3 that lie outside of the F and HN genes, e.g. the N protein, attenuate the virus for rhesus monkeys. This confirms that the genetic basis for host range restriction can involve multiple genes.

The chimeric BPIV3 bearing HPIV3 glycoprotein genes induces serum HAI antibody to HPIV3 and a high level of resistance to wt HPIV3 challenge.

rBPIV3-F_HHN_H has important features that make it a candidate live attenuated virus vaccine against HPIV3, including attenuating genes from BPIV3 and the antigenic specificity of HPIV3, i.e. the F and HN glycoproteins, which are the major protective antigens. Therefore, its immunogenicity and protective efficacy against challenge with HPIV3 were documented. Rhesus monkeys were immunized by infection with BPIV3 Ka, rHPIV3-F_BHN_B, rBPIV3-F_HHN_H, rHPIV3-N_B, or rHPIV3. They were challenged 28 days later with HPIV3 JS wild type virus. Serum samples were taken prior to the initial infection on day 0 and prior to the challenge. BPIV3 and rHPIV3-F_BHN_B induced serum HAI antibodies that reacted more efficiently with BPIV3 than HPIV3, whereas the converse was the case for HPIV3 and rBPIV3-F_HHN_H. Thus, the origin of the glycoprotein genes in each virus determined whether the HAI antibody response was directed predominantly against HPIV3 or against BPIV3. The replication of challenge HPIV3 virus was significantly reduced in the upper and lower respiratory tract of previously immunized monkeys (Table 4). Although the level of protective efficacy against HPIV3 was not significantly different among the different viruses,

viruses bearing HPIV3 F and HN were consistently more protective in the upper respiratory tract than were viruses bearing BPIV3 F and HN. This is in accordance with the higher level of HPIV3-specific serum HAI antibodies induced by viruses bearing HPIV3 F and HN.

5

Table 4. Immunization of rhesus monkeys with BPIV3/HPIV3 chimeric recombinant virus induces resistance to challenge with wild type HPIV3

Immunizing virus ¹	Number of animals ²	Mean peak virus titer ³		Serum HAI antibody		Serum HAI antibody titer (mean recip. log ₂ ±S.E.) for HPIV3 28 days after challenge
		(log ₁₀ TCID ₅₀ /ml ± S.E.)	[Duncan Grouping] ⁴	titer (mean recip. log ₂ ±S.E.) for HPIV3 on the day of challenge		
		Nasopharyngeal swab ⁵		Tracheal lavage ⁶		
none	4	4.5±0.33 [A]	4.5±0.19 [A]	<2		12.0±0.58 [A]
rHPIV3	6	2.3±0.14 [B]	1.2±0.20 [B]	9.5±0.72 [A]		11.7±0.21 [A]
rBPIV3-F _H HN _H	4	2.5±0.25 [B]	1.0±0.48 [B]	6.8±0.63 [BC]		10.5±0.29 [AB]
rHPIV3-N _B	6	2.3±0.41 [B]	1.4±0.08 [B]	8.2±0.48 [AB]		11.5±0.22 [A]
rHPIV3-F _B HN _B	4	3.0±0.14 [B]	1.0±0.0 [B]	4.5±0.29 [D]		9.5±0.87 [B]
BPIV3 Ka	6	2.9±0.26 [B]	1.3±0.20 [B]	5.5±0.62 [CD]		9.3±0.76 [B]

¹ Each previously immunized monkey and non-immunized controls were challenged with 10⁶ TCID₅₀ of HPIV3 JS in a 1 ml inoculum at each site 28 days after immunization.

² The groups with 6 animals contain 4 animals each from a previous rhesus study (Bailey et al., 2000, *supra*).

³ Mean of the peak virus titers for each animal in its group irrespective of sampling day.

⁴ Virus titrations were performed on LLC-MK2 cells. The limit of detectability of virus titer was 10 TCID₅₀/ml. Mean viral titers were compared using a Duncan Multiple Range test ($\alpha=0.05$). Within each column, mean titers with different letters are statistically different. Titers indicated with two letters are not significantly different from those indicated with either letter. The group of unimmunized animals were not included in the Duncan analysis at the day of challenge.

⁵ Nasopharyngeal swab samples were collected on days 3 to 8 post challenge.

⁶ Tracheal lavage samples were collected on days 4, 6 and 8 post challenge.

**Two animals in the rHPIV3 group were infected with rHPIV3s.

Based on the foregoing examples, the invention provides for importation of BPIV genes into a virulent HPIV backbone and visa versa to yield novel, human-bovine chimeric PIV vaccine candidates. In exemplary chimeric recombinants disclosed in the present example, rBPIV3-F_HHN_H and its rHPIV3-F_BHN_B counterpart, replicated in vitro as well as the respective parental viruses. It was also confirmed that the F and HN exchange between the BPIV3 and HPIV3 is compatible since the considerably more divergent HPIV1 F and HN proteins were highly functional in a HPIV3 background (Tao et al., J. Virol. 72:2955-2961, 1998), which was evinced by the undiminished capacity of the chimeric viruses for replication in vitro. rBPIV3-F_HHN_H replicated in the upper respiratory tract of rhesus monkeys to a level intermediate between that of its HPIV3 and BPIV3 parents indicating that the BPIV3 F and HN genes make an independent contribution to the overall attenuation of BPIV3 for primates. The overall attenuation of BPIV3 virus thus is the sum of two or more genetic elements, one of which is the set of F and HN genes and one of the others is indicated to be N.

Although BPIV3 itself is being evaluated as a vaccine virus for HPIV3 (Karron et al., Pediatr. Infect. Dis. J. 15:650-654, 1996; and Karron et al., J. Infect. Dis. 171:1107-1114, 1995), it is only 25% related antigenically to HPIV3 (Coelingh et al., J. Infect. Dis. 157:655-662, 1988). Thus, the immunogenicity of BPIV3 against HPIV3 will be improved if it is modified according to the present invention to express the protective F and HN antigens of HPIV3. rBPIV3-F_HHN_H represents such a virus, and, in the present example, immunization of rhesus monkeys with rBPIV3-F_HHN_H induced a higher level of antibody to HPIV3 than did immunization with BPIV3. Furthermore, rBPIV3-F_HHN_H conferred a level of protection against replication of HPIV3 challenge in the upper and lower respiratory tract that was statistically indistinguishable from that conferred by a previous infection with rHPIV3. Similarly, rHPIV3-N_B, which is attenuated by the BPIV3 N protein but possesses HPIV3 protective antigens, also induced a high level of resistance to HPIV3 challenge. Despite replicating to similar levels in rhesus monkeys, rHPIV3-N_B induced higher levels of antibodies to HPIV3 than rBPIV3-F_HHN_H.

rBPIV3-F_HHN_H replicates to higher levels in rhesus monkeys than BPIV3, although it is significantly attenuated compared to HPIV3. Since the level of replication of BPIV3 in humans is low (Karron et al., J. Infect. Dis. 171:1107-1114, 1995), this increase is expected to be well tolerated among vaccinees. Alternatively, additional methods to attenuate human-bovine chimeric viruses of the invention are disclosed herein

WO 01/04320

to ensure that the vaccine viruses replicate only to moderate levels, for example in human infants, to prevent unacceptable respiratory tract illness among vaccinees. Within other aspects of the invention, the slight increase in replication of rBPIV3-F_HHN_H in primates offers an opportunity to use rBPIV3-F_HHN_H as a vector for heterologous viral antigens such as glycoproteins of other PIVs (e.g., HPIV1 and HPIV2), the RSV F and G glycoproteins, and the measles HA glycoprotein, which can be incorporated as added or substituted gene(s) or genome segment(s) into the attenuated HPIV3 vaccine candidate. In various alternative embodiments disclosed herein, the slight increase in replication of rBPIV3-F_HHN_H in monkeys over that of BPIV3 can be offset by the addition of foreign viral protective antigens, e.g., RSV glycoproteins, whose addition provides a selected level of attenuation. The data presented here further defined the basis for the host range restriction of BPIV3 for primates and identify rBPIV3-F_HHN_H as a potential vaccine candidate against HPIV3 and as a vector for heterologous viral antigens.

Microorganism Deposit Information

The following materials have been deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209, under the conditions of the Budapest Treaty and designated as follows:

	<u>Plasmid</u>	<u>Accession No.</u>	<u>Deposit Date</u>
	p3/7(131)	(ATCC 97990)	April 18, 1997
	p3/7(131)2G	(ATCC 97989)	April 18, 1997
	p218(131)	(ATCC 97991)	April 18, 1997

Although the foregoing invention has been described in detail by way of example for purposes of clarity of understanding, it will be apparent to the artisan that certain changes and modifications are comprehended by the disclosure and may be practiced without undue experimentation within the scope of the appended claims, which are presented by way of illustration not limitation.

WHAT IS CLAIMED IS:

1. An isolated infectious human-bovine chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a partial or complete PIV background genome or antigenome of a human PIV (HPIV) or bovine PIV (BPIV) combined with one or more
5 heterogeneous gene(s) or genome segment(s) of a different PIV to form a human-bovine chimeric PIV genome or antigenome.
2. The chimeric PIV of claim 1, wherein said one or more heterogeneous gene(s) or genome segment(s) encodes one or more PIV N, P, C, D, V, M, F, HN and/or L protein(s) or fragment(s) thereof.
- 10 3. The chimeric PIV of claim 1, wherein said one or more heterogeneous gene(s) or genome segment(s) encodes a complete open reading frame (ORF) of one or more PIV N, P, C, D, V, M, F, HN and/or L protein(s).
4. The chimeric PIV of claim 1, wherein said one or more heterogeneous gene(s) or genome segment(s) includes a heterogeneous regulatory element
15 comprising an extragenic 3' leader or 5' trailer region, a gene-start signal, gene-end signal, RNA editing site, encapsidation signal, intergenic region, or 3' or 5' non-coding region.
5. The chimeric PIV of claim 1, wherein said background genome or antigenome incorporates a heterogeneous genome segment integrated with the background
20 genome or antigenome to form a chimeric gene.
6. The chimeric PIV of claim 5, wherein said chimeric genome or antigenome encodes a chimeric glycoprotein.
7. The chimeric PIV of claim 1, wherein a heterogeneous gene or genome segment is substituted for a counterpart gene or genome segment in a partial PIV
25 background genome or antigenome.
8. The chimeric PIV of claim 1, wherein a heterogeneous gene or genome segment is added adjacent to or within a noncoding region of the partial or complete PIV background genome or antigenome.

9. The chimeric PIV of claim 1, wherein a heterologous gene or genome segment is added or substituted at a position corresponding to a wild-type gene order position of a counterpart gene or genome segment within the partial or complete PIV background genome or antigenome.

5 10. The chimeric PIV of claim 1, wherein a heterologous gene or genome segment is added or substituted at a position that is more promoter-proximal or promoter-distal compared to a wild-type gene order position of a counterpart gene or genome segment within the partial or complete PIV background genome or antigenome.

10 11. The chimeric PIV of claim 1, wherein the chimeric genome or antigenome comprises a partial or complete BPIV background genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a human PIV.

15 12. The chimeric PIV of claim 11, wherein one or more HPIV glycoprotein genes selected from HN and F, or one or more genome segments encoding a cytoplasmic domain, transmembrane domain, ectodomain or immunogenic epitope thereof, is/are substituted for one or more counterpart genes or genome segments within the BPIV background genome or antigenome.

20 13. The chimeric PIV of claim 11, wherein one or more HPIV glycoprotein genes selected from HN and F is/are substituted to replace one or more counterpart glycoprotein genes in the BPIV background genome or antigenome.

14. The chimeric PIV of claim 13, wherein both HPIV glycoprotein genes HN and F are substituted to replace counterpart HN and F glycoprotein genes in the BPIV background genome or antigenome.

15. The chimeric PIV of claim 13, which is rBPIV3-F_HHN_H.

25 16. The chimeric PIV of claim 11, wherein the human-bovine chimeric PIV genome or antigenome encodes a chimeric glycoprotein having a HPIV glycoprotein ectodomain, antigenic determinant or immunogenic epitope.

17. The chimeric PIV of claim 16, wherein the heterologous genome segment encodes a glycoprotein ectodomain.
18. The chimeric PIV of claim 11, wherein one or more HPIV glycoprotein genes HN and F, or a genome segment encoding a cytoplasmic domain,
5 transmembrane domain, ectodomain or immunogenic epitope thereof is added to or incorporated within a BPIV background genome or antigenome.
19. The chimeric PIV of claim 11, wherein the chimeric genome or antigenome is further modified by addition or substitution of one or more additional heterologous gene(s) or genome segment(s) from a human PIV within the partial or
10 complete bovine background genome or antigenome to increase genetic stability or alter attenuation, reactogenicity or growth in culture of the chimeric virus.
20. The chimeric PIV of claim 1, wherein the chimeric genome or antigenome comprises a partial or complete human PIV background genome or antigenome combined with one or more heterologous genes or genome segments from a
15 bovine PIV.
21. The chimeric PIV of claim 20, wherein the heterologous gene or genome segment encodes a bovine PIV3 N protein.
22. The chimeric PIV of claim 20, wherein a bovine PIV3 N open reading frame (ORF) is substituted for a human PIV3 N ORF in the chimeric genome or
20 antigenome.
23. The chimeric PIV of claim 22, which is rHPIV3-N_B.
24. The chimeric PIV of claim 1, wherein the genome or antigenome is further modified by introduction of one or more attenuating mutations identified in a biologically derived mutant PIV or other mutant nonsegmented negative stranded RNA
25 virus.
25. The chimeric PIV of claim 24, wherein the genome or antigenome incorporates at least one and up to a full complement of attenuating mutations present within PIV3 JS cp45.

26. The chimeric PIV of claim 24, wherein the genome or antigenome incorporates at least one and up to a full complement of attenuating mutations specifying an amino acid substitution in the L protein at a position corresponding to Tyr₉₄₂, Leu₉₉₂, or Thr₁₅₅₈ of JS; in the N protein at a position corresponding to residues Val₉₆ or Ser₃₈₉ of JS, in the C protein at a position corresponding to Ile₉₆ of JS, in the M protein at a position corresponding to Pro₁₉₉ of JS in the F protein at a position corresponding to residues Ile₄₂₀ or Ala₄₅₀ of JS, in the HN protein at a position corresponding to residue Val₃₈₄ of JS, a nucleotide substitution a 3' leader sequence of the chimeric virus at a position corresponding to nucleotide 23, 24, 28, or 45 of JS cp45, and/or a mutation in an N gene start sequence at a position corresponding to nucleotide 62 of JS cp45.

27. The chimeric PIV of claim 24, wherein the genome or antigenome incorporates attenuating mutations from different biologically derived mutant PIVs.

28. The chimeric PIV of claim 24, wherein the genome or antigenome incorporates an attenuating mutation at an amino acid position corresponding to an amino acid position of an attenuating mutation identified in a heterologous, mutant negative stranded RNA virus.

29. The chimeric PIV of claim 24, wherein the genome or antigenome includes at least one attenuating mutation stabilized by multiple nucleotide changes in a codon specifying the mutation.

30. The chimeric PIV of claim 1, wherein the genome or antigenome comprises an additional nucleotide modification specifying a phenotypic change selected from a change in growth characteristics, attenuation, temperature-sensitivity, cold-adaptation, plaque size, host-range restriction, or a change in immunogenicity.

31. The chimeric PIV of claim 30, wherein the additional nucleotide modification alters one or more of the PIV N, P, C, D, V, M, F, HN and/or L genes and/or a 3' leader, 5' trailer RNA editing site, encapsidation signal, and/or an intergenic region.

32. The chimeric PIV of claim 30, wherein one or more genes of the chimeric virus is deleted in whole or in part or expression of the genes is reduced or ablated by a mutation in an RNA editing site, by a frameshift mutation, by a mutation that

alters an amino acid specified by an initiation codon, or by introduction of one or more stop codons in an open reading frame (ORF) of the gene.

33. The chimeric PIV of claim 30, wherein a modification is introduced in the chimeric genome or antigenome comprising a partial or complete
5 deletion of one or more C, D and/or V ORF(s) or one or more nucleotide change(s) that reduces or ablates expression of said one or more C, D and/or V ORF(s).

34. The chimeric PIV of claim 30, wherein the chimeric genome or antigenome is modified to encode a non-PIV molecule selected from a cytokine, a T-lymphocyte helper epitope, a restriction site marker, or a protein of a microbial pathogen
10 capable of eliciting a protective immune response in a mammalian host.

35. The chimeric PIV of claim 1, wherein the bovine-human chimeric genome or antigenome comprises a partial or complete PIV vector genome or antigenome combined with one or more heterologous genes or genome segments encoding one or more antigenic determinants of one or more heterologous pathogens.

15 36. The chimeric PIV of claim 35, wherein said one or more heterologous pathogens is a heterologous PIV and said heterologous gene(s) or genome segment(s) encode(s) one or more PIV N, P, C, D, V, M, F, HN and/or L protein(s) or fragment(s) thereof.

37. The chimeric PIV of claim 35, wherein the vector genome or
20 antigenome is a partial or complete HPIV genome or antigenome and the heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are of one or more heterologous PIV(s).

38. The chimeric PIV of claim 37, wherein said one or more heterologous PIV(s) is/are selected from HPIV1, HPIV2, or HPIV3.

25 39. The chimeric PIV of claim 37, wherein the vector genome or antigenome is a partial or complete HPIV genome or antigenome and the heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are of one or more heterologous HPIV(s).

40. The chimeric PIV of claim 39, wherein the vector genome or antigenome is a partial or complete HPIV3 genome or antigenome and the heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are of one or more heterologous HPIV(s).

5 41. The chimeric PIV of claim 40, wherein the chimeric genome or antigenome incorporates one or more gene(s) or genome segment(s) of a BPIV that specifies attenuation.

42. The chimeric PIV of claim 35, wherein one or more HPIV1 or HPIV2 gene(s) or genome segment(s) encoding one or more HN and/or F glycoprotein(s) or antigenic domain(s), fragment(s) or epitope(s) thereof is/are added to or incorporated
10 within the partial or complete HPIV3 vector genome or antigenome.

43. The chimeric PIV of claim 35, wherein both HPIV1 genes encoding HN and F glycoproteins are substituted for counterpart HPIV3 HN and F genes to form a chimeric HPIV3-1 vector genome or antigenome which is further modified by
15 addition or incorporation of one or more gene(s) or gene segment(s) encoding one or more antigenic determinant(s) of HPIV2 and one or more heterologous gene(s) or genome segment(s) of a BPIV that specifies attenuation.

44. The chimeric PIV of claim 43, wherein a transcription unit comprising an open reading frame (ORF) of an HPIV2 HN or F gene is added to or
20 incorporated within the chimeric HPIV3-1 vector genome or antigenome.

45. The chimeric PIV of claim 35, wherein the vector genome or antigenome is a partial or complete BPIV genome or antigenome and the heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are of one or more HPIV(s).

25 46. The chimeric PIV of claim 45, wherein said one or more antigenic determinant(s) is/are selected from HPIV1, HPIV2 or HPIV3 HN and F glycoproteins and antigenic domains, fragments and epitopes thereof.

47. The chimeric PIV of claim 45, wherein one or more gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of HPIV2 is/are added to or substituted within the partial or complete BPIV vector genome or antigenome.

48. The chimeric PIV of claim 45, wherein a plurality of heterologous
5 genes or genome segments encoding antigenic determinants of multiple HPIVs are added to or incorporated within the partial or complete BPIV vector genome or antigenome.

49. The chimeric PIV of claim 35, wherein the vector genome or antigenome is a partial or complete HPIV genome or antigenome and the heterologous pathogen is selected from measles virus, subgroup A and subgroup B respiratory
10 syncytial viruses, mumps virus, human papilloma viruses, type 1 and type 2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses.

50. The chimeric PIV of claim 35, wherein the vector genome or antigenome is a partial or complete BPIV genome or antigenome and the heterologous
15 pathogen is selected from measles virus, subgroup A and subgroup B respiratory syncytial viruses, mumps virus, human papilloma viruses, type 1 and type 2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses.

51. The chimeric PIV of claim 50, wherein said one or more
20 heterologous antigenic determinant(s) is/are selected from measles virus HA and F proteins, subgroup A or subgroup B respiratory syncytial virus F, G, SH and M2 proteins, mumps virus HN and F proteins, human papilloma virus L1 protein, type 1 or type 2 human immunodeficiency virus gp160 protein, herpes simplex virus and cytomegalovirus gB, gC, gD, gE, gG, gH, gI, gJ, gK, gL, and gM proteins, rabies virus G protein, Epstein
25 Barr Virus gp350 protein; filovirus G protein, bunyavirus G protein, Flavivirus E and NS1 proteins, and alphavirus E protein, and antigenic domains, fragments and epitopes thereof.

52. The chimeric PIV of claim 51, wherein the heterologous pathogen is measles virus and the heterologous antigenic determinant(s) is/are selected from the
30 measles virus HA and F proteins and antigenic domains, fragments and epitopes thereof.

53. The chimeric PIV of claim 52, wherein a transcription unit comprising an open reading frame (ORF) of a measles virus HA gene is added to or incorporated within a HPIV3 vector genome or antigenome.

54. The chimeric PIV of claim 51, which incorporates a gene or
5 genome segment from respiratory syncytial virus (RSV).

55. The chimeric PIV of claim 54, wherein the gene or genome segment encodes a RSV F and/or G glycoprotein or immunogenic domain(s) or epitope(s) thereof.

56. The chimeric PIV of claim 1 which is a virus.

10 57. The chimeric PIV of claim 1 which is a subviral particle.

58. A method for stimulating the immune system of an individual to induce protection against PIV which comprises administering to the individual an immunologically sufficient amount of the chimeric PIV of claim 1 combined with a physiologically acceptable carrier.

15 59. The method of claim 58, wherein the chimeric PIV is administered in a dose of 10^3 to 10^7 PFU.

60. The method of claim 58, wherein the chimeric PIV is administered to the upper respiratory tract.

20 61. The method of claim 58, wherein the chimeric PIV is administered by spray, droplet or aerosol.

62. The method of claim 58, wherein the chimeric PIV and a second attenuated PIV are administered simultaneously as a mixture.

25 63. An immunogenic composition to elicit an immune response against PIV comprising an immunogenically sufficient amount of the chimeric PIV of claim 1 in a physiologically acceptable carrier.

64. The immunogenic composition of claim 63, formulated in a dose of 10^3 to 10^7 PFU.

65. The immunogenic composition of claim 63, formulated for administration to the upper respiratory tract by spray, droplet or aerosol.

66. The immunogenic composition of claim 63, wherein the chimeric PIV elicits an immune response against one or more virus(es) selected from HPIV 1,
5 HPIV2 and HPIV3.

67. The immunogenic composition of claim 66, wherein the chimeric PIV elicits an immune response against HPIV3 and another virus selected from HPIV 1, HPIV2 and HPIV3.

68. An isolated polynucleotide molecule comprising a chimeric PIV
10 genome or antigenome which includes a partial or complete PIV background genome or antigenome of a human or bovine PIV combined with a heterologous gene or genome segment of a different PIV to form a human-bovine chimeric PIV genome or antigenome.

69. The isolated polynucleotide of claim 68, wherein the chimeric genome or antigenome comprises a partial or complete BPIV background genome or
15 antigenome combined with one or more heterologous gene(s) or genome segment(s) from a human PIV.

70. The isolated polynucleotide of claim 69, wherein one or more HPIV glycoprotein genes selected from HN and F, or one or more genome segments encoding a cytoplasmic domain, transmembrane domain, ectodomain or immunogenic
20 epitope thereof, is/are substituted for one or more counterpart genes or genome segments within the BPIV background genome or antigenome.

71. The isolated polynucleotide of claim 69, wherein one or more HPIV glycoprotein genes selected from HN and F is/are substituted to replace one or more counterpart glycoprotein genes in the BPIV background genome or antigenome.

25 72. The isolated polynucleotide of claim 71, wherein both HPIV glycoprotein genes HN and F are substituted to replace counterpart HN and F glycoprotein genes in the BPIV background genome or antigenome.

73. The isolated polynucleotide of claim 71, which comprises a plasmid pBPIV(215).

74. The isolated polynucleotide molecule of claim 68, wherein the chimeric genome or antigenome comprises a partial or complete human PIV background genome or antigenome combined with one or more heterologous genes or genome segments from a bovine PIV.

75. The isolated polynucleotide of claim 74, wherein one or more BPIV glycoprotein genes selected from HN and F is/are substituted to replace one or more counterpart glycoprotein genes in the HPIV background genome or antigenome.

76. The isolated polynucleotide of claim 75, wherein both BPIV glycoprotein genes HN and F are substituted to replace counterpart HN and F glycoprotein genes in the HPIV background genome or antigenome.

77. The isolated polynucleotide of claim 76, which comprises a plasmid pHPIV(215).

78. The isolated polynucleotide molecule of claim 74, wherein a bovine N gene is substituted to replace a counterpart N gene in a human PIV background genome or antigenome.

79. The isolated polynucleotide molecule of claim 68, wherein the chimeric genome or antigenome is further modified by one or more attenuating mutations.

80. The isolated polynucleotide molecule of claim 68, further comprising a nucleotide modification specifying a phenotypic change selected from a change in growth characteristics, attenuation, temperature-sensitivity, cold-adaptation, plaque size, host-range restriction, or a change in immunogenicity.

81. A method for producing an infectious attenuated chimeric PIV particle from one or more isolated polynucleotide molecules encoding said PIV, comprising:


expressing in a cell or cell-free lysate an expression vector comprising an isolated polynucleotide comprising a partial or complete PIV background genome or antigenome of a human or bovine PIV combined with a heterologous gene or genome segment of a different PIV to form a human-bovine chimeric PIV genome or antigenome,
5 and PIV N, P, and L proteins.

82. The method of claim 81, wherein the chimeric PIV genome or antigenome and the N, P, and L proteins are expressed by two or more different expression vectors.


83. An expression vector comprising an operably linked transcriptional
10 promoter, a polynucleotide sequence which includes a partial or complete PIV background genome or antigenome of a human or bovine PIV combined with one or more heterologous gene(s) or genome segment(s) of a different PIV to form a human-bovine chimeric PIV genome or antigenome, and a transcriptional terminator.

Figure 1. Cloning of BPIV3 strain Ka or strain SF coding region into HPIV3 context

A. Mutagenesis to create restriction sites at start and stop codons of N



B.



C. Mutagenesis to restore start and stop codon context

Legend





Figure 2. Cloning of BPIV3 N coding region into HPV3 antigenomic cDNA




Figure 3. Nucleotide sequences of HPIV3, BPIV3 and chimeric viruses around the start (A) and stop (B) codons of the N gene

A. rJS GGAACTCTATAATTCAAAAAATGTTGAGCCTATTTGATAC
 cKa GGAACTCTATAATTCAAAAAATGTTGAGTCTATTCGACAC
 cSF GGAACTCTATAATTCAAAAAATGTTGAGTCTATTCGACAC
 Ka GAAATCCTAAGACTGTAATCATGTTGAGTCTATTCGACAC
 SF GAAATCCTAAGACTGTAATCATGTTGAGTCTATTCGACAC

B. rJS TTAACGCATTTGGAAGCAACTAATCGAATCAACATTTTAA
 cKa TCAGTGCATTCGGAAGCAACTAGTTCGAATCAACATTTTAA
 cSF TCAGTGCATTCGGAAGCAACTAGTTCGAATCAACATTTTAA
 Ka TCAGTGCATTCGGAAGCAACTAGTTCACAAAGAGATGACCA
 SF TCAGTGCATTCGGAAGCAACTAGTTCACAAAGAGATGACCA





Figure 4. Confirmation of identity of potential BPIV3/HPIV3 primers by *TaqI* digestion

Figure 5. Multicycle growth curves in MDBK (A) or LLC-MK2 (B) cells



WO 01/04320

6/25

ACCAAACAAG	AGAAGAGACT	TGCTTGGGAA	TATTAATTCA	AATAAAAATT	50
AACCTTAGGAT	TAAAGAACTT	TACCGAAAGG	TAAGGGGAAA	GAAATCCTAA	100
GACTGTAATC	ATGTTGAGTC	TATTCGACAC	ATTGAGTGCG	CGTAGGCAGG	150
AGAACATAAC	GAAATCAGCT	GGTGGGGCTG	TTATTTCCCGG	GCAAAAAAAC	200
ACTGTGTCTA	TATTTGCTCT	TGGACCATCA	ATAACAGATG	ACAATGATAA	250
AATGACATTG	GCTCTTCTCT	TTTTGTCTCA	TTCTTTAGAC	AATGAAAAGC	300
AGCATGCGCA	AAGAGCTGGA	TTTTTAGTTT	CTCTGTTATC	AATGGCTTAT	350
GCCAACCCAG	AATTATATTT	AACATCAAAT	GGTAGTAATG	CAGATGTAA	400
ATATGTTATC	TACATGATAG	AGAAAGACCC	AGGAAGACAG	AAATATGGTG	450
GGTTTGTCTG	CAAGACTAGA	GAGATGGTTT	ATGAAAAGAC	AACTGATTGG	500
ATGTTGCGGA	GTGATCTTGA	GTATGATCAA	GACAATATGT	TGCAAAATGG	550
TAGAAGCACT	TCTACAATCG	AGGATCTTGT	TCATACTTTT	GGATATCCAT	600
CGTGTCTTGG	AGCCCTTATA	ATCCAAGTTT	GGATAATACT	TGTTAAGGCT	650
ATAACCACTA	TATCAGGATT	GAGGAAAGGA	TTCTTTACTC	GGTTAGAAGC	700
ATTTCGACAA	GATGGAACAG	TTAAATCCAG	TCTAGTGTTG	AGCGGTGATG	750
CAGTAGAACA	AATTGGATCA	ATTATGAGGT	CCCAACAGAG	CTTGGTAACA	800
CTCATGGTTG	AAACACTGAT	AACAATGAAC	ACAGGCAGGA	ATGATCTGAC	850
AACAATAGAA	AAGAATATAC	AGATTGTAGG	AAACTACATC	AGAGATGCAG	900
GTCTTGCTTC	ATTTTTCAAC	ACAATCAGAT	ATGGCATTGA	GAAGATGATG	950
GCAGCTCTAA	CTCTGTCTAC	CCTTAGACCG	GATATCAACA	GAAGCAAGGC	1000
ACTGATCGAG	TTATATCTAT	CAAAGGGGCC	ACGTGCTCCT	TTTATATGCA	1050
TTTTGAGAGA	TCCCGTGCAT	GGTGAGTTTG	CACCAGGCAA	CTATCCTGCC	1100
CTCTGGAGTT	ATGCGATGGG	TGTAGCAGTT	GTACAAAACA	AGGCCATGCA	1150
ACAGTATGTA	ACAGGAAGGT	CTTATCTGGA	TATTGAAATG	TTCCAAGTTG	1200
GTCAAGCAGT	GGCACGTGAT	GCCGAGTCGC	AGATGAGTTC	AATATTAGAG	1250
GATGAAGTGG	GGGTCACACA	AGAAGCCAAG	CAAAGCTTGA	AGAAACACAT	1300
GAAGAACATC	AGCAGTTCAG	ATACAACCTT	TCATAAGCCT	ACAGGGGGAT	1350
CAGCCATAGA	AATGGCGATA	GATGAAGAAG	CAGGGCAGCC	TGAATCCAGA	1400
GGAGATCAGG	ATCAAGGAGA	TGAGCCTCGG	TCATCCATAG	TTCTTATGCA	1450
ATGGGCAGAC	GAAACCGGGA	ATGACAATCA	AACTGAATCA	ACTACAGAAA	1500
TTGACAGCAT	CAAAACTGAA	CAAAAGAAACA	TCAGAGACAG	GCTGAACAAA	1550
AGACTCAACG	AGAAAAGGAA	ACAGAGTGAC	CCGAGATCAA	CTGACATCAC	1600
AAACAACACA	AATCAAAGTG	AAATAGATGA	TTTGTTTCACT	GCATTGCGAA	1650
GCAACTAGTC	ACAAAGAGAT	GACCACTATC	ACCAGCAACA	AGTAAGAAAA	1700
ACTTAGGATT	AATGGAAATT	ATCCAATCCA	GAGACGGAAG	GACAAATCCA	1750
GAATCCAACC	ACAACTCAAT	CAACCAAGA	TTCATGGAAG	ACAATGTTCA	1800
AAACAATCAA	ATCATGGATT	CTTGGGAAGA	GGGATCAGGA	GATAAATCAT	1850
CTGACATCTC	ATCGGCCCTC	GACATCATTG	AATTCATACT	CAGCACCAGC	1900
TCCCAAGAGA	ACACGGCAGA	CAGCAATGAA	ATCAACACAG	GAACCACAAG	1950
ACTTAGCAGG	ACAATCTACC	AACCTGAATC	CAAAACAACA	GAAACAAGCA	2000
AGGAAAATAG	TGGACCAGCT	AACAAAAATC	GACAGTTTGG	GGCATCACAC	2050
GAACGTGCCA	CAGAGACAAA	AGATAGAAAT	GTAAATCAGG	AGACTGTACA	2100
GGGAGGATAT	AGGAGAGGAA	GCAGCCAGGA	TAGTAGAACT	GAGACTATGG	2150
TCACTCGAAG	AATCTCCAGA	AGCAGCCAG	ATCCTAACAA	TGGAACCCAA	2200
ATCCAGGAAG	ATATTGATTA	CAATGAAGTT	GGAGAGATGG	ATAAGGACTC	2250
TACTAAGAGG	GAAATGCGAC	AATTTAAAGA	TGTTCCAGTC	AAGGTATCAG	2300
GAAGTGATGC	CATTCCTCCA	ACAAAACAAG	ATGGAGACGG	TGATGATGGA	2350

FIG. 6A

WO 01/04320

7/25

AGAGGCCTGG	AATCTATCAG	TACATTTGAT	TCAGGATATA	CCAGTATAGT	2400
GACTGCCGCA	ACACTAGATG	ACGAAGAAGA	ACTCCTTATG	AAGAACAACA	2450
GGCCAAGAAA	GTATCAATCA	ACACCCCAGA	ACAGTGACAA	GGAATTAAA	2500
AAAGGGGTTG	GAAGGCCAAA	AGACACAGAC	AAACAATCAT	CAATATTGGA	2550
CTACGAACTC	AACCTCAAAG	GATCGAAGAA	GAGCCAGAAA	ATCCTCAAAG	2600
CCAGCACGAA	TACAGGAGAA	CCAACAAGAC	CACAGAATGG	ATCCCAGGGG	2650
AAGAGAATCA	CATCCTGGAA	CATCCTCAAC	AGCGAGAGCG	GCAATCGAAC	2700
AGAATCAACA	AACCAAACCC	ATCAGACATC	AACCTCGGGA	CAGAACCACA	2750
CAATGGGACC	AAGCAGAACA	ACCTCCGAAC	CAAGGATCAA	GACACAAAAG	2800
ACGGATGGAA	AGGAAAGAGA	GGACACAGAA	GAGAGCACTC	GATTTACAGA	2850
AAGGGCGATT	ACATTATTAC	AGAATCTTGG	TGTAATCCAA	TCTGCAGCAA	2900
AATTAGACCT	ATACCAAGAC	AAGAGAGTTG	TGTGTGTGGC	GAATGTCCTA	2950
AACAATGCAG	ATACTGCATC	AAAGATAGAC	TTCCTAGCAG	GTTTGATGAT	3000
AGGAGTGTC	ATGGATCATG	ATACCAAATT	AAATCAGATT	CAGAACGAGA	3050
TATTAAGTTT	GAAAACCTGAT	CTTAAAAAGA	TGGATGAATC	ACATAGAAGA	3100
CTAATTGAGA	ATCAAAAAGA	ACAATTATCA	CTGATCACAT	CATTAATCTC	3150
AAATCTTAAA	ATTATGACAG	AGAGAGGAGG	GAAGAAGGAC	CAACCAGAAC	3200
CTAGCGGGAG	GACATCCATG	ATCAAGACAA	AAGCAAAAGA	AGAGAAAATA	3250
AAGAAAGTCA	GGTTTGACCC	TCTTATGGAA	ACACAGGGCA	TCGAGAAAAA	3300
CATCCCTGAC	CTCTATAGAT	CAATAGAGAA	AACACCAGAA	AACGACACAC	3350
AGATCAAATC	AGAAATAAAC	AGATTGAATG	ATGAATCCAA	TGCCACTAGA	3400
TTAGTACCTA	GAAGAATAAG	CAGTACAATG	AGATCATTAA	TAATAATCAT	3450
TAACAACAGC	AATTTATCAT	CAAAAGCAAA	GCAATCATA	ATCAACGAAC	3500
TCAAGCTCTG	CAAGAGTGAC	GAGGAAGTGT	CTGAGTTGAT	GGACATGTTT	3550
AATGAGGATG	TCAGCTCCCA	GTAAACCGCC	AACCAAGGGT	CAACACCAAG	3600
AAAACCAATA	GCACAAAACA	GCCAATCAGA	GACCACCCCA	ATACACCAAA	3650
CCAATCAACA	CATAACAAAG	ATCTCCAGAT	CATAGATGAT	TAAGAAAAAC	3700
TTAGGATGAA	AGGACTAATC	AATCCTCCGA	AACAATGAGC	ATCACCAACT	3750
CCACAATCTA	CACATTCCCA	GAATCCTCTT	TCTCCGAGAA	TGGCAACATA	3800
GAGCCGTTAC	CACTCAAGGT	CAATGAACAG	AGAAAGGCCA	TACCTCATAT	3850
TAGGGTTGTC	AAGATAGGAG	ATCCGCCCAA	ACATGGATCC	AGATATCTGG	3900
ATGTCTTTTT	ACTGGGCTTC	TTTGAGATGG	AAAGGTCAAA	AGACAGGTAT	3950
GGGAGCATAA	GTGATCTAGA	TGATGATCCA	AGTTACAAGG	TTTGTGGCTC	4000
TGGATCATTG	CCACTTGGGT	TGGCTAGATA	CACCGGAAAT	GATCAGGAAC	4050
TCCTACAGGC	TGCAACCAAG	CTCGATATAG	AAGTAAGAAG	AACTGTAAAG	4100
GCTACGGAGA	TGATAGTTTA	CACTGTACAA	AACATCAAAC	CTGAACTATA	4150
TCCATGGTCC	AGTAGATTAA	GAAAAGGGAT	GTTATTTGAC	GCTAATAAGG	4200
TTGCACTTGC	TCCTCAATGT	CTTCCACTAG	ATAGAGGGAT	AAAATTCAGG	4250
GTGATATTTG	TGAACTGCAC	AGCAATTGGA	TCAATAACTC	TATTCAAAT	4300
CCCTAAGTCC	ATGGCATTGT	TATCATTGCC	TAATACAATA	TCAATAAATC	4350
TACAAGTACA	TATCAAAAACA	GGAGTTCAGA	CAGATTCCAA	AGGAGTAGTT	4400
CAGATTCTAG	ATGAAAAAGG	TGAAAAATCA	CTAAATTTCA	TGGTTCATCT	4450
CGGGTTGATC	AAAAGGAAGA	TGGGCAGAAT	GTAATCAGTT	GAATATTGTA	4500
AGCAGAAGAT	CGAGAAGATG	AGATTATTAT	TCTCATTGGG	ATTAGTTGGA	4550
GGGATCAGCT	TCCACGTCAA	CGCAACTGGC	TCTATATCAA	AGACATTAGC	4600
AAGTCAATTA	GCATTCAAAA	GAGAAATCTG	CTATCCCCTA	ATGGATCTGA	4650
ATCCACACTT	AAATTCAGTT	ATATGGGCAT	CATCAGTTGA	AATTACAAGG	4700

FIG. 6B

GTAGATGCAG	TTCTCCAGCC	TTCATTACCT	GGCGAATTCA	GATACTACCC	4750
AAACATCATA	GCAAAAGGGG	TCGGGAAAAT	CAGACAGTAA	AATCAACAAC	4800
CCTGATATCC	AACATTGCAA	ATCAGGCTAC	CCACAGGAGA	AAAATCAAAA	4850
ACTTAGGATC	AAAGGGATCA	CCACGAACCC	CGGAAAACAG	CCAAACAAAC	4900
CAACACACAA	ATCACAGACA	AAAAGGAGAA	GGCACTGCAA	AGACCGAGAA	4950
AAAACAGAAC	GCACACAACC	AAGCAGAGAA	AAGCCAAAGC	CCGCCATTCA	5000
CAAACACACC	AACAATCCTG	CAAACAAGCA	CCAAAACAGA	GGTCAAAAGA	5050
CAAAGAGCAC	CAGATATGAC	CATCACAACC	ACAATCATAG	CCATATTACT	5100
AATACCCCCA	TCATTTTGTC	AAATAGACAT	AACAAAACCTG	CAACGTGTAG	5150
GTGTGTTAGT	CAACAATCCT	AAAGGCATGA	AGATTTCACA	AAATTTTCGAA	5200
ACGAGATACC	TGATATTAAG	TTTGATACCC	AAAATAGAGA	ATTCACACTC	5250
ATGTGGGGAT	CAACAGATAA	ACCAATACAA	GAAGTTATTG	GATAGATTGA	5300
TAATTCCTCT	ATATGATGGA	TTAAAAATTAC	AAAAAGATGT	AATAGTAGTA	5350
AGTCATGAAA	CCCACAACAA	TACTAATCTT	AGGACAAAAC	GATTCTTTGG	5400
AGAGATAATT	GGGACAATTG	CGATAGGGAT	AGCCACTTCA	GCACAAATCA	5450
CCGCAGCAGT	CGCTCTTGTC	GAAGCTAAAC	AGGCAAAGTC	AGACATAGAA	5500
AAACTCAAAG	AGGCTATAAG	AGACACAAAC	AAGGCAGTAC	AATCGATTCA	5550
AAGTTCTGTA	GGTAACCTAA	TTGTTGCAGT	TAAATCAGTT	CAAGACTATG	5600
TCAACAATGA	AATTATACCT	TCAATCACAA	GATTAGGCTG	TGAAGCAGCA	5650
GGGTTACAAT	TGGGAATTGC	ATTGACACAA	CATTACTCAG	AATTAACAAA	5700
TATATTTGGT	GATAATATAG	GAACACTGAA	AGAAAAAGGG	ATAAAATTAC	5750
AAGGGATAGC	ATCATTATAT	CACACAAACA	TAACGGAAAT	ATTTACTACT	5800
TCAACAGTTG	ACCAATATGA	TATTTATGAC	CTATTATTCA	CTGAGTCAAT	5850
CAAGATGAGA	GTGATAGATG	TTGATTTGAG	TGATTACTCA	ATTACTCTTC	5900
AAGTTAGACT	TCCTTTATTA	ACTAACTAT	CAAATACTCA	AATTTATAAA	5950
GTAGATTCTA	TATCATACAA	CATCCAGGGC	AAAGAGTGGT	ATATTCCTCT	6000
TCCCAATCAC	ATCATGACAA	AAGGGGCTTT	TCTAGGTGGT	GCTGATATTA	6050
AAGAATGCAT	AGAGGCATTG	AGCAGTTATA	TATGTCCTTC	TGATCCAGGT	6100
TACATATTAA	ATCACGAGAT	AGAGAATTGT	TTATCAGGGA	ACATAACACA	6150
GTGTCCTAAG	ACTGTTGTGA	CATCAGATGT	GGTACCACGA	TACGCGTTTG	6200
TGAATGGTGG	ATTAATTGCA	AACTGCATAA	CAACTACATG	TACATGCAAT	6250
GGAATTGACA	ATAGAATTAA	TCAATCACCT	GATCAAGGAA	TTAAGATCAT	6300
AACACATAAA	GAATGCCAGG	TAATAGGTAT	AAACGGAATG	TTATTCAATA	6350
CTAATAGAGA	AGGGACATTA	GCAACTTATA	CATTTGATGA	CATCATATTA	6400
AATAACTCTG	TTGCACTTAA	TCCAATTGAT	ATATCTATGG	AACTCAACAA	6450
GGCAAACTA	GAATTAGAAG	AATCGAAGGA	ATGGATAAAG	AAATCAAATC	6500
AAAAGTTAGA	TTCCGTTGGA	AGTTGGTATC	AATCTAGTGC	AACAATCACC	6550
ATAATCATAG	TGATGATAAT	AATTCTAGTT	ATAATCAATA	TAACAATTAT	6600
TGTAGTCATA	ATCAAATTCC	ATAGAATTCA	GGGGAAAGAT	CAAAACGACA	6650
AAAACAGTGA	GCCGTATATA	CTGACAAATA	GACAATAAGA	CTATACACGA	6700
TCAAATATAA	AAAGTACAAA	AACTTAGGA	ACAAAGTTGT	TCAACACAGC	6750
AGCACCGAAT	AGACCAAAAAG	GCAGCGCAGA	GGCGACACCA	AACTCAAAAA	6800
TGGAATATTG	GAAACACACA	AACAGCATAA	ATAACACCAA	CAATGAAACC	6850
GAAACAGCCA	GAGGCAAAACA	TAGTAGCAAG	GTTACAAATA	TCATAATGTA	6900
CACCTTCTGG	ACAATAACAT	TAACAATATT	ATCAGTCATT	TTTATAATGA	6950
TATTGACAAA	CTTAATTCAA	GAGAACAATC	ATAATAAATT	AATGTTGCAG	7000
GAAATAAGAA	AAGAATTCGC	GGCAATAGAC	ACCAAGATTC	AGAGGACTTC	7050

FIG. 6C

WO 01/04320

9/25

GGATGACATT	GGAACCTCAA	TACAGTCAGG	AATAAATACA	AGACTTCTCA	7100
CAATTCAGAG	TCATGTTCAA	AACTATATCC	CACTATCATT	AACACAACAA	7150
ATGTCAGATC	TCAGAAAATT	TATCAATGAT	CTAACAAATA	AAAGAGAACA	7200
TCAAGAAGTG	CCAATACAGA	GAATGACTCA	TGATAGAGGT	ATAGAACCCC	7250
TAAATCCAAA	CAAGTTCTGG	AGGTGTACAT	CTGGTAACCC	ATCTCTAACA	7300
AGTAGTCCTA	AGATAAGGTT	AATACCAGGA	CCAGGTTTAT	TAGCAACATC	7350
TACTACAGTA	AATGGCTGTA	TTAGAATTCC	ATCGTTAGTA	ATCAATCATC	7400
TAATCTATGC	TTACACCTCT	AATCTTATTA	CCCAGGGCTG	TCAAGATATA	7450
GGGAAATCTT	ACCAAGTACT	ACAAATAGGG	ATAATTACTA	TAAATTCGGA	7500
CCTAGTACCT	GATTTAAACC	CCAGAGTCAC	ACATACATTT	AATATTGATG	7550
ATAATAGAAG	ATCTTGCTCT	CTGGCACTAT	TGAATACAGA	TGTTTATCAG	7600
TTATGCTCAA	CACCAAAAGT	TGATGAAAGA	TCCGATTATG	CATCAACAGG	7650
TATTGAGGAT	ATTGTACTTG	ACATTGTCAC	TAATAATGGA	TTAATTATAA	7700
CAACAAGGTT	TACAAATAAT	AATATAACTT	TTGATAAACC	GTATGCAGCA	7750
TTGTATCCAT	CAGTGGGACC	AGGAATCTAT	TATAAGGATA	AAGTTATATT	7800
TCTCGGATAT	GGAGGTCTAG	AGCATGAAGA	AAACGGAGAC	GTAATATGTA	7850
ATACAACCTGG	TTGTCCTGGC	AAAACACAGA	GAGACTGTAA	TCAGGCTTCT	7900
TATAGCCCAT	GGTTCTCAAA	TAGGAGAATG	GTAAACTCTA	TTATTGTTGT	7950
TGATAAAGGC	ATAGATGCAA	CTTTTAGCTT	GAGGGTGTGG	ACTATTCCAA	8000
TGAGCCAAAA	TTATTGGGGA	TCAGAAGGAA	GATTACTTTT	ATTAGGTGAC	8050
AGAATATACA	TATATACTAG	ATCCACAAGT	TGGCACAGTA	AATTACAGTT	8100
AGGGGTAATT	GATATTTCTG	ATTATACTAA	TATAAGAATA	AATTGGACTT	8150
GGCATAATGT	ACTATCACGG	CCAGGGAATG	ATGAATGTCC	ATGGGGTCAT	8200
TCATGCCCAG	ACGGATGTAT	AACAGGAGTT	TACACTGATG	CATATCCGCT	8250
AAACCCATCG	GGGAGTGTTG	TATCATCAGT	AATTCTTGAT	TCACAAAAGT	8300
CTAGAGAAAA	CCCAATCATT	ACTTACTCAA	CAGCTACAAA	TAGAATAAAT	8350
GAATTAGCTA	TATATAACAG	AACACTTCCA	GCTGCATATA	CAACAACAAA	8400
TTGTATCACA	CATTATGATA	AAGGGTATTG	TTTTCATATA	GTAGAAATAA	8450
ATCACAGAAG	TTTGAATACG	TTTCAACCTA	TGTTATTCAA	AACAGAAGTT	8500
CCAAAAAACT	GCAGCTAAAT	TGATCATCGC	ATATCGGATG	CAAGATGACA	8550
TTAAAAGAGA	CCACCAGACA	GACAACACAG	GAGACGATGC	AAGATATAAA	8600
GAAATAATAA	AAAACCTTAGG	AGAAAAGTGT	GCAAGAAAAA	TGGACACCGA	8650
GTCCACACAGC	GGCACAACAT	CTGACATTCT	GTACCCTGAA	TGTCACCTCA	8700
ATTCTCCTAT	AGTTAAAGGA	AAGATAGCAC	AACTGCATAC	AATAATGAGT	8750
TTGCCTCAGC	CCTACGATAT	GGATGATGAT	TCAATACTGA	TTATTACTAG	8800
ACAAAAAATT	AAACTCAATA	AATTAGATAA	AAGACAACGG	TCAATTAGGA	8850
AATTAAGATC	AGTCTTAATG	GAAAGAGTAA	GTGATCTAGG	TAAATATACC	8900
TTTATCAGAT	ATCCAGAGAT	GTCTAGTGAA	ATGTTCCAAT	TATGTATACC	8950
CGGAATTAAT	AATAAAATAA	ATGAATTGCT	AAGTAAAGCA	AGTAAAACAT	9000
ATAATCAAAT	GACTGATGGA	TTAAGAGATC	TATGGGTTAC	TATACTATCG	9050
AAGTTAGCAT	CGAAAAATGA	TGGAAGTAAT	TATGATATCA	ATGAAGATAT	9100
TAGCAATATA	TCAAATGTTC	ACATGACTTA	TCAATCAGAC	AAATGGTATA	9150
ATCCATTCAA	GACATGGTTT	ACTATTAAGT	ATGACATGAG	AAGATTACAA	9200
AAAGCCAAAA	ATGAGATTAC	ATTCAATAGG	CATAAAGATT	ATAATCTATT	9250
AGAAGACCAA	AAGAATATAT	TGCTGATACA	TCCAGAACTC	GTCTTAATAT	9300
TAGATAAACA	AAATTACAAT	GGGTATATAA	TGACTCCTGA	ATTGGTACTA	9350
ATGTATTGTG	ATGTAGTTGA	AGGGAGGTGG	AATATAAGTT	CATGTGCAAA	9400

FIG. 6D

WO 01/04320	10/25	
ATTGGATCCT AAGTTACAAT CAATGTATTA TAAGGGTAAC AATTTATGGG	9450	
AAATAATAGA TGGACTATTC TCGACCTTAG GAGAAAGAAC ATTTGACATA	9500	
ATATCACTAT TAGAACCACT TGCATTATCG CTCATTCAAA CTTATGACCC	9550	
GGTTAAACAG CTCAGGGGGG CTTTTTAA TCACGTGTTA TCAGAAATGG	9600	
AATTAATATT TGCAGCTGAG TGTACAACAG AGGAAATACC TAATGTGGAT	9650	
TATATAGATA AAATTTTAGA TGTGTTCAAA GAATCAACAA TAGATGAAAT	9700	
AGCAGAAATT TTCTCTTTCT TCCGAACCTT TGGACACCCT CCATTAGAGG	9750	
CGAGTATAGC AGCAGAGAAA GTTAGAAAGT ATATGTATAC TGAGAAATGC	9800	
TTGAAATTTG ATACTATCAA TAAATGTCAT GCTATTTTTT GTACAATAAT	9850	
TATAAATGGA TATAGAGAAA GACATGGTGG TCAATGGCCT CCAGTTACAT	9900	
TACCTGTCCA TGCACATGAA TTTATCATAA ATGCATACGG ATCAAATTCT	9950	
GCCATATCAT ATGAGAATGC TGTAGATTAT TATAAGAGCT TCATAGGAAT	10000	
AAAATTTGAC AAGTTTATAG AGCCTCAATT GGATGAAGAC TTAACATTT	10050	
ATATGAAAGA TAAAGCATTA TCCCCAAGA AATCAAACTG GGACACAGTC	10100	
TATCCAGCTT CAAACCTGTT ATACCGCACT AATGTGTCTC ATGATTCAG	10150	
AAGATTGGTT GAAGTATTTA TAGCAGATAG TAAATTTGAT CCCCACCAAG	10200	
TATTAGATTA CGTAGAATCA GGATATTGGC TGGATGATCC TGAATTTAAT	10250	
ATCTCATATA GTTTAAAGA GAAAGAAATA AAACAAGAAG GTAGACTTTT	10300	
TGCAAAAATG ACATACAAGA TGAGGGCTAC ACAAGTATTA TCAGAAACAT	10350	
TATTGGCGAA TAATATAGGG AAATCTTCC AAGAGAATGG GATGGTTAAA	10400	
GGAGAAATTG AATTACTCAA GAGACTAACA ACAATATCTA TGTCTGGAGT	10450	
TCCGCGGTAT AATGAGGTAT ACAATAATTC AAAAGTCAC ACAGAAGAAC	10500	
TTCAAGCTTA TAATGCAATT AGCAGTTCCA ATTTATCTTC TAATCAGAAG	10550	
TCAAAGAAGT TTGAATTTAA ATCTACAGAT ATATACAATG ATGGATACGA	10600	
AACCGTAAGC TGCTTCTTAA CGACAGATCT TAAAAAATAT TGTTTAAATT	10650	
GGAGGTATGA ATCAACAGCT TTATTCCGTG ATACTTGTA TCAGATATTT	10700	
GGGTAAAGG AATTATTTAA TTGGCTGCAC CCTCGCCTTG AAAAGAGTAC	10750	
AATATATGTT GGAGATCCTT ATTGCCCGCC ATCAGATATT GAACATTTAC	10800	
CACTTGATGA CCATCCTGAT TCAGGATTTT ATGTTTCATA TCCTAAAGGA	10850	
GGAATAGAAG GGTTTTGCCA AAAGTTATGG AACTCATAT CTATCAGTGC	10900	
AATACATTTA GCAGCTGTCA AAATCGGTGT AAGAGTTACT GCAATGGTTC	10950	
AAGGGGATAA TCAAGCCATA GCTGTTACCA CAAGAGTACC TAATAATTAT	11000	
GATTATAAAG TTAAGAAAGA GATTGTTTAT AAAGATGTGG TAAGATTTTT	11050	
TGATTCCTTG AGAGAGGTGA TGGATGATCT GGGTCATGAG CTCAAATAA	11100	
ATGAACTAT AATAAGTAGT AAAATGTTTA TATATAGCAA AAGGATATAC	11150	
TATGACGGAA GAATCCTTCC TCAGGCATTA AAAGCATTGT CTAGATGTGT	11200	
TTTTTGGTCT GAAACAATCA TAGATGAGAC AAGATCAGCA TCCTCAAATC	11250	
TGGCTACATC GTTTGCAAAG GCCATTGAGA ATGGCTACTC ACCTGTATTG	11300	
GGATATGTAT GCTCAATCTT CAAAAATATC CAACAGTTGT ATATAGCGCT	11350	
TGGAATGAAT ATAAACCCAA CTATAACCCA AAATATTAAA GATCAATATT	11400	
TCAGGAATAT TCATTGGATG CAATATGCCT CCTTAATCCC TGCTAGTGTC	11450	
GGAGGATTTA ATTATATGGC CATGTCAAGG TGTTTTGTCA GAAACATTGG	11500	
AGATCCTACA GTCGCTGCGT TAGCCGATAT TAAAAGATTT ATAAAAGCAA	11550	
ATTTGTTAGA TCGAGGTGTC CTTTACAGAA TTATGAATCA AGAACCAGGC	11600	
GAGTCTTCTT TTTTAGACTG GGCCTCAGAT CCCTATTTCAT GTAACCTACC	11650	
ACAATCTCAA AATATAACCA CCATGATAAA GAATATAACT GCAAGAAATG	11700	
TACTACAGGA CTCACCAAAC CCATTACTAT CTGGATTATT TACAAGTACA	11750	

FIG. 6E

11/25

ATGATAGAAG	AGGATGAGGA	ATTAGCTGAG	TTCCTAATGG	ACAGGAGAAT	11800
AATCCTCCCA	AGAGTTGCAC	ATGACATTTT	AGATAATTCT	CTTACTGGAA	11850
TTAGGAATGC	TATAGCTGGT	ATGTTGGATA	CAACAAAATC	ACTAATTCGA	11900
GTAGGGATAA	GCAGAGGAGG	ATTAACCTAT	AACTTATTAA	GAAAGATAAG	11950
CAACTATGAT	CTTGTACAAT	ATGAGACACT	TAGTAAAAC	TTAAGACTAA	12000
TAGTCAGTGA	CAAGATTAAG	TATGAAGATA	TGTGCTCAGT	AGACCTAGCC	12050
ATATCATTAA	GACAAAAAAT	GTGGATGCAT	TTATCAGGAG	GAAGAATGAT	12100
AAATGGACTT	GAAACTCCAG	ATCCTTTAGA	GTTACTGTCT	GGAGTAATAA	12150
TAACAGGATC	TGAACATTGT	AGGATATGTT	ATTCAACTGA	AGGTGAAAGC	12200
CCATATACAT	GGATGTATTT	ACCAGGCAAT	CTTAATATAG	GATCAGCTGA	12250
GACAGGAATA	GCATCATTAA	GGGTCCCTTA	CTTTGGATCA	GTTACAGATG	12300
AGAGATCTGA	AGCACAATTA	GGGTATATCA	AAAATCTAAG	CAAACCAGCT	12350
AAGGCTGCTA	TAAGAATAGC	AATGATATAT	ACTTGGGCAT	TTGGGAATGA	12400
CGAAATATCT	TGGATGGAAG	CATCACAGAT	TGCACAAACA	CGTGCAAAC	12450
TTACATTGGA	TAGCTTAAAG	ATTTTGACAC	CAGTGACAAC	ATCAACAAAT	12500
CTATCACACA	GGTTAAAAAG	TACTGCTACT	CAGATGAAAT	TTTCTAGTAC	12550
ATCACTTATT	AGAGTAAGCA	GGTTCATCAC	AATATCTAAT	GATAATATGT	12600
CTATTAAAGA	AGCAAATGAA	ACTAAAGATA	CAAATCTTAT	TTATCAACAG	12650
GTAATGTAA	CAGGATTAAG	TGTATTTGAA	TATCTATTTA	GGTTAGAGGA	12700
GAGTACAGGA	CATAACCCTA	TGGTCATGCA	TCTACATATA	GAGGATGGAT	12750
GTTGTATAAA	AGAGAGTTAC	AATGATGAGC	ATATCAATCC	GGAGTCTACA	12800
TTAGAGTTAA	TCAAATACCC	TGAGAGTAAT	GAATTTATAT	ATGATAAGGA	12850
CCCTTTAAAG	GATATAGATC	TATCAAAATT	AATGGTTATA	AGAGATCATT	12900
CTTATACAAT	TGACATGAAT	TACTGGGATG	ACACAGATAT	TGTACATGCA	12950
ATATCAATAT	GTACTGCAGT	TACAATAGCA	GATACAATGT	CGCAGCTAGA	13000
TCGGGATAAT	CTTAAGGAGC	TGGTTGTGAT	TGCAAATGAT	GATGATATTA	13050
ACAGTCTGAT	AACTGAATTT	CTGACCCTAG	ATATACTAGT	GTTTCTCAAA	13100
ACATTGAGG	GGTTACTCGT	GAATCAATTT	GCATATACCC	TTTATGGATT	13150
GAAAATAGAA	GGAAGGGATC	CCATTGCGA	TTATATAATG	AGAACATTAA	13200
AAGACACCTC	ACATTTCAGTA	CTTAAAGTAT	TATCTAATGC	ACTATCTCAT	13250
CCAAAAGTGT	TTAAGAGATT	TTGGGATTGT	GGAGTTTTGA	ATCCTATTTA	13300
TGGTCCTAAT	ACTGCTAGTC	AAGATCAAGT	TAAGCTTGCT	CTCTCGATTT	13350
GCGAGTACTC	CTTGGATCTA	TTTATGAGAG	AATGGTTGAA	TGGAGCATCA	13400
CTTGAGATCT	ATATCTGTGA	TAGTGACATG	GAAATAGCAA	ATGACAGAAG	13450
ACAAGCATTT	CTCTCAAGAC	ATCTTGCCCT	TGTGTGTTGT	TTAGCAGAGA	13500
TAGCATCTTT	TGGACCAAAT	TTATTAAATC	TAACATATCT	AGAGAGACTT	13550
GATGAATTAA	AACAATACTT	AGATCTGAAC	ATCAAAGAAG	ATCCTACTCT	13600
TAAATATGTG	CAAGTATCAG	GACTGTTAAT	TAAATCATT	CCCTCAACTG	13650
TTACGTATGT	AAGGAAAAC	GCGATTAAAGT	ATCTGAGGAT	TCGTGGTATT	13700
AATCCGCCTG	AAACGATTGA	AGATTGGGAT	CCCATAGAAG	ATGAGAATAT	13750
CTTAGACAAT	ATTGTTAAAA	CTGTAAATGA	CAATTGCAGT	GATAATCAAA	13800
AGAGAAATAA	AAGTAGTTAT	TTCTGGGGAT	TAGCTCTAAA	GAATTATCAA	13850
GTCGTGAAAA	TAAGATCCAT	AACGAGTGAT	TCTGAAGTTA	ATGAAGCTTC	13900
GAATGTTACT	ACACATGGAA	TGACACTTCC	TCAGGGAGGA	AGTTATCTAT	13950
CACATCAGCT	GAGGTTATTT	GGAGTAAACA	GTACAAGTTG	TCTTAAAGCT	14000
CTTGAATTAT	CACAAATCTT	AATGAGGGAA	GTTAAAAAAG	ATAAAGATAG	14050
ACTCTTTTTA	GGAGAAGGAG	CAGGAGCTAT	GTTAGCATGT	TATGATGCTA	14100

FIG. 6F

12/25

CACTCGGTCC	TGCAATAAAT	TATTATAATT	CTGGTTTAA	TATTACAGAT	14150
GTAATTGGTC	AACGGGAATT	AAAAATCTTC	CCATCAGAAG	TATCATTAGT	14200
AGGTAAAAAA	CTAGGAAATG	TAACACAGAT	TCTTAATCGG	GTGAGGGTGT	14250
TATTTAATGG	GAATCCCAAT	TCAACATGGA	TAGGAAATAT	GGAATGTGAG	14300
AGTTTAATAT	GGAGTGAATT	AAATGATAAG	TCAATTGGTT	TAGTACATTG	14350
TGACATGGAG	GGAGCGATAG	GCAAATCAGA	AGAAACTGTT	CTACATGAAC	14400
ATTATAGTAT	TATTAGGATT	ACATATTTAA	TCGGGGATGA	TGATGTTGTC	14450
CTAGTATCAA	AAATTATACC	AACTATTACT	CCGAATTGGT	CTAAAATACT	14500
CTATCTATAC	AAGTTGTATT	GGAAGGATGT	AAGTGTAGTG	TCCCTTAAAA	14550
CATCCAATCC	TGCCTCAACA	GAGCTTTATT	TAATTTCAAA	AGATGCTTAC	14600
TGTACTGTAA	TGGAACCCAG	TAATCTTGTT	TTATCAAAAC	TTAAAAGGAT	14650
ATCATCAATA	GAAGAAAATA	ATCTATTAAA	GTGGATAATC	TTATCAAAAA	14700
GGAAGAATAA	CGAGTGGTTA	CAGCATGAAA	TCAAAGAAGG	AGAAAGGGAT	14750
TATGGGATAA	TGAGGCCATA	TCATACAGCA	CTGCAAATTT	TTGGATTCCA	14800
AATTAACCTA	AATCACTTAG	CTAGAGAATT	TTTATCAACT	CCTGATTTAA	14850
CCAACATTAA	TAATATAATT	CAAAGTTTTA	CAAGAACAAT	TAAAGATGTT	14900
ATGTTCTGAAT	GGGTCAATAT	CACTCATGAC	AATAAAAAGAC	ATAAATTAGG	14950
AGGAAGATAT	AATCTATTCC	CGCTTAAAAA	TAAGGGGAAA	TTAAGATTAT	15000
TATCACGAAG	ATTAGTACTA	AGCTGGATAT	CATTATCCTT	ATCAACCAGA	15050
TTACTGACGG	GCCGTTTCC	AGATGAAAAA	TTTGAAAATA	GGGCACAGAC	15100
CGGATATGTA	TCATTGGCTG	ATATTGATTT	AGAATCCTTA	AAGTTATTAT	15150
CAAGAAATAT	TGTCAAAAAT	TACAAAGAAC	ACATAGGATT	AATATCATAC	15200
TGGTTTTTGA	CCAAAGAGGT	CAAATACTA	ATGAAGCTTA	TAGGAGGAGT	15250
CAAATACTA	GGAATTCCTA	AACAGTACAA	AGAGTTAGAG	GATCGATCAT	15300
CTCAGGGTTA	TGAATATGAT	AATGAATTTG	ATATTGATTA	ATACATAAAA	15350
ACATAAAATA	AAACACCTAT	TCCTCACCCA	TTCACCTCCA	ACAAAATGAA	15400
AAGTAAGAAA	AACATGTAAT	ATATATATAC	CAAACAGAGT	TTTTCTCTTG	15450
TTTGGT					15456

FIG. 6G

ACCAAACAAG	AGAAGAGACT	TGCTTGGGAA	TATTAATTCA	AATAAAAATT	50
AACTTAGGAT	TAAAGAACTT	TACCGAAAGG	TAAGGGGAAA	GAAATCCTAA	100
GACTGTAATC	ATGTTGAGTC	TATTCGACAC	ATTCAGTGCG	CGTAGGCAGG	150
AGAACATAAC	AAAATCAGCT	GGTGGGGCTG	TTATTCCCGG	GCAAAAAAAC	200
ACTGTGTCTA	TATTTGCTCT	TGGACCATCA	ATAACAGATG	ACAATGACAA	250
AATGACATTG	GCTCTTCTCT	TTTTGTCTCA	TTCTTTAGAC	AATGAAAAGC	300
AGCATGCGCA	AAGAGCTGGA	TTTTTAGTTT	CTCTGTTATC	AATGGCTTAT	350
GCCAACCCAG	AATTATATTT	AACATCAAAT	GGTAGTAATG	CAGATGTTAA	400
ATATGTCATC	TACATGATAG	AGAAAGACCC	AGGAAGACAG	AAATATGGTG	450
GGTTTGTCTG	CAAGACTAGA	GAGATGGTTT	ATGAAAAGAC	AACTGACTGG	500
ATGTTTGGGA	GTGATCTTGA	GTATGATCAA	GACAATATGT	TGCAAAATGG	550
TAGAAGCACT	TCTACAATCG	AGGATCTTGT	TCATACTTTT	GGATATCCAT	600
CGTGTCTTGG	AGCCCTTATA	ATCCAGGTTT	GGATAATACT	TGTTAAGGCT	650
ATAACCAGTA	TATCAGGATT	GAGGAAAGGA	TTCTTTACTC	GGTTAGAAGC	700
ATTTGCACAA	GATGGAACAG	TTAAATCCAG	TCTAGTGTTG	AGCGGTGATG	750
CAGTAGAACA	AATTGGATCA	ATTATGAGCT	CCCAACAGAG	CTTGGTAACA	800
CTCATGGTTG	AAACACTGAT	AACAATGAAC	ACAGGCAGGA	ATGACCTGAC	850
AACAATAGAA	AAGAATATAC	AGATTGTAGG	AAACTACATC	AGAGATGCAG	900
GTCTTGCTTC	ATTTTCAAC	ACAATCAGAT	ATGGCATTGA	GACTIONATG	950
GCAGCTCTAA	CTCTGTCTAC	CCTTAGACCG	GACATCAACA	GACTIONAGGC	1000
ACTGATAGAG	CTATATCTAT	CAAAGGGGCC	ACGTGCTCCT	TTTATATGCA	1050
TTTTGAGAGA	TCCTGTGCAT	GGTGAGTTG	CACCAGGCAA	CTATCCTGCC	1100
CTCTGGAGTT	ATGCGATGGG	TGTAGCAGTT	GTACAAAACA	AGGCCATGCA	1150
ACAGTATGTA	ACAGGAAGGT	CCTATCTGGA	TATTGAAATG	TTCCAACCTG	1200
GTCAAGCAGT	GGCACGTGAC	GCCGAGTCGC	AGATGAGTTC	AATATTAGAG	1250
GATGAACCTG	GGGTCACACA	AGAAGCCAAG	CAAAGCTTGA	AGAAACACAT	1300
GAAGAACATC	AGCAGTTCAG	ATACAACCTT	CTATAAGCCT	ACAGGGGGAT	1350
CAGCCATAGA	AATGGCAATA	GATGAGGAAG	CAGAGCAGCC	CGAATCCAGA	1400
GGAGACCAAG	ACCAAGGAGA	TGAACCTCGG	TCATCCATAG	TTCTTATGTC	1450
ATGGGCAGAC	GAAACCGGGA	ATGACAACCA	AACTGAATCA	ACCACAGAAA	1500
TTGACAGCAT	CAAAACTGAA	CAAAGAAACA	TCAGAGACAG	GCTGAACAAA	1550
AGACTCAACG	AGAAAAGGAA	ACAGAGTAAC	CCGGGATCAA	CTGACATCAC	1600
AAACAACACA	AATCAAACCTG	AAATAGATGA	TTTATTTCAGT	GCATTCCGGA	1650
GCAACTAGTC	ACAAAGAGAT	GACCACCATC	ATCAGCAACA	AGTAAGAAAA	1700
ACTTAGGATT	AATGGAAATT	ATCCAATCCG	GAGACGGAAG	GACAAATCCA	1750
GAATCCAACC	ACAACCTCAAT	CAACCAAAGA	TTCATGGAAG	ACAATGTTCA	1800
AAACAATCAA	ATCATGGATT	CTTGGGAAGA	GGGATCAGGA	GATAAATCAT	1850
CTGACATCTC	ATCGGCCCTC	GACATCATTG	AATTCATACT	CAACACCGAC	1900
TCCCAAGAGA	ACACGGCAGA	CAGCAATGAA	ATCAACACAG	GAGCCACAAG	1950
ACTTAGCACG	ACAATCTACC	AACTTGAGTC	CAAAACAACA	GAAACAAGCA	2000
AGGAAAATAG	TGGACCAGCT	AACAAAAATC	GACAGTTTGG	GGCATCACAC	2050
GAACGTGCCA	CAGAGACAAA	AGATAGAAAT	GTTAATCAGA	AGACTGTACA	2100
GGGAGGATAT	AGGAGAGGAA	GCAGCCCAGA	TAGTAGAACT	GAGACTATGG	2150
TCACTCGAGG	AATCTCCAGA	AGCAGCCCAG	ATCCTAACAA	TGGAACCCAA	2200
ATCCAGGAAG	ATATTGATTA	CAATGAAGTT	GGAGAGATGG	ATAAGGACTC	2250
TACTAAGAGG	GAAATGCGAC	AATTTAAAGA	TGTTCCAGTC	AAGGTATCAG	2300
GAAGTGATGC	CATTCCTCCA	ACAAAACAAG	ATGGAGACGG	TGATGATGGA	2350

FIG. 7A

AGAGGCCTGG	AATCTATCAG	TACATCTGAT	TCAGGATATA	CCAGTATAGT	2400
GACTGCCGCA	ACACTAGATG	ACGAAGAAGA	ACTCCTTATG	AAGAACAACA	2450
GGCCAAGAAA	GTATCAATCA	ACACCCCAGA	ACAGTGACAA	GGGAATTAAA	2500
AAAGGGAGTG	GAAGGCCAAA	AGACACAGAC	AAACAATCAC	CAATATTGGA	2550
CTACGAAGTC	AACTCCAAAG	GATCGAAGAA	GAGCCAGAAA	ATCCTCAAAG	2600
CCAGCACGAA	TACAGGAGAA	CCAACAAGAT	CACAGAGTGG	ATCCCAGGGG	2650
AAGAGAATCA	CATCCTGGAA	CATCCTCAAC	AGCGAGAGCG	GCAATCGAGC	2700
AGAATCAACA	AACCAAACCC	ATCAGACATC	AATCTCGGGA	CAGAACCACA	2750
CAATGGGACC	AAGCAGAACA	ACCTCAGAAC	CAAGGACCAA	GACACAAAAG	2800
ACGGATGGAA	AGGAAAGAGA	GGACACAGAA	GAGAGCACTC	GATTTACAGA	2850
AAGGGCGATT	ACATTATTAC	AGAATCTTGG	TGTAATCCAA	TCTGCAGCAA	2900
AATTAGACCT	ATACCAAGAC	AAGAGAGTTG	TGTGTGTGGC	GAATGTCCTA	2950
AACAATGCAG	ATACTGCATC	AAAGATAGAC	TTCCTAGCAG	GTTTGATGAT	3000
AGGAGTGTCA	ATGGATCATG	ATGTCAAATT	AAATCAGATT	CAGAACGAGA	3050
TATTAAGTTT	AAAAACTGAT	CTTAAGAAGA	TGGATGAATC	ACATAGAAGA	3100
CTAATTGAGA	ATCAAAAAGA	ACAATTATCA	CTGATCACAT	CATTAATCTC	3150
AAATCTTAAA	ATCATGACAG	AGAGAGGAGG	GAAGAAGGAC	CAACCAGAAC	3200
CTAGCGGGAG	GACATCCATG	ATCAAGACAA	AGGCAAAAGA	AGAGAGAATA	3250
AAGAAAGTCA	GGTTTGACCC	TCTTATGGAA	ACACAGGGCA	TCGAGAAAAA	3300
CATCCCTGAC	CTCTACAGAT	CAATAGAGAA	AACACCAGAA	AACGACACAC	3350
AGATCAAATC	AGAAATAAAC	AGATTGAATG	ATGAATCCAA	TGCCACTAGA	3400
TTAGTACCTA	GAAGAATAAG	CAGTACAATG	AGATCACTAA	TAATAATCAT	3450
CAACAACAGC	AATTTATCAT	CAAAAGCAAA	GCAATCATAC	ATCAACGAAC	3500
TCAAGCTCTG	CAAGAGTGAT	GAGGAAGTGT	CTGAGTTGAT	GGACATGTTC	3550
AATGAGGATG	TCAGCTCCCA	GTAAACCGCC	AACCAAGGGT	CAACACCAAG	3600
AAAACCAACA	GCACAAAACA	GCCAATAAGA	GACCATCCCA	ACACACCGAA	3650
CCAATCAACA	CATAACAAAG	ATCTTTAGAT	CATAGATGAC	TAAGAAAAAC	3700
TTAGGATGAA	AGGACTGATC	AATCCTCCAA	AACAATGAGC	ATCACCAGCT	3750
CCACAATCTA	CACATTCCCA	GAATCCTCTT	TCTCCGAGAA	TGGCAACATA	3800
GAGCCGTTAC	CACTCAAGGT	CAATGAACAG	AGAAAGGCCA	TACCTCATAT	3850
TAGGGTTGTC	AAGATAGGAG	ATCCGCCCAA	ACATGGATCC	AGATATCTGG	3900
ATGTCTTTTT	ACTGGGCTTC	TTTGAAATGG	AAAGGTCAAA	AGACAGGTAT	3950
GGGAGCATAA	GTGATCTAGA	TGATGATCCA	AGTTACAAGG	TTTGTGGCTC	4000
TGGATCATTG	CCACTTGGGT	TGGCTAGATA	CACTGGAAAT	GATCAGGAAC	4050
TCCTACAGGC	TGCAACCAAG	CTCGATATAG	AAGTAAGAAG	AACTGTAAAG	4100
GCTACGGAGA	TGATAGTTTA	CACTGTGCAA	AACATCAAAC	CTGAACTATA	4150
TCCATGGTCC	AGTAGATTAA	GAAAAGGGAT	GTTATTTGAC	GCTAACAAGG	4200
TTGCACTTGC	TCCTCAATGT	CTTCCACTAG	ATAGAGGGAT	AAAATTTCAGG	4250
GTGATATTTG	TGAÄCTGCAC	AGCAATTGGA	TCAATAACTC	TATTCAAAAT	4300
CCCCAAGTCC	ATGGCATTGT	TATCATTGCC	TAATACAATA	TCAATAAATC	4350
TACAAGTACA	TATCAAAACA	GGAATTCAGA	CAGATTCCAA	AGGAGTAGTT	4400
CAGATTCTAG	ATGAAAAAGG	TGAAAAATCA	CTAAATTTCA	TGGTTCATCT	4450
CGGGTTGATC	AAAAGGAAGA	TGGGTAGAAT	GTACTCAGTT	GAATATTGTA	4500
AGCAGAAGAT	TGAGAAGATG	AGATTATTAT	TCTCATTGGG	ATTAGTTGGA	4550
GGGATCAGCT	TCCACGTCAA	CGCAACTGGC	TCTATATCAA	AGACATTAGC	4600
AAGTCAATTA	GCATTTAAAA	GAGAAATCTG	CTATCCCCTA	ATGGATCTGA	4650
ATCCCACTT	AAATTTAGTT	ATATGGGCAT	CATCAGTTGA	AATTACAAGA	4700

FIG. 7B

GTAGATGCAA	TTCTCCAGCC	TTCATTACCT	GGCGAATTCA	GATACTACCC	4750
AAACATCATA	GCAAAAGGGG	TCGGGAAAAT	CAGACAGTAA	AACCAACAAC	4800
CCTGACATCC	AACACTGCAA	ATCAGGCTAC	CCACAGGAGA	AAAATCAAAA	4850
ACTTAGGATC	AAAGGGATCA	CCACAAACCC	CGGGAAACAG	CCAAACCAAC	4900
CAACACACAA	ATCACAGACA	AAAAGGAAAA	GGCACTGCAA	AGACCGAGAA	4950
CAAGCAGAAC	GCACACAACC	AAGCAGAGGA	AAGCCAAAGC	CCGCCATTCA	5000
CAAACACACC	AACAATCCTA	CAAACAAGCA	CCAAAATAGA	GGTCAAAAGA	5050
CAAAGAGCAT	CAGATATGAC	CATCACAACC	ATAATCATAG	CCATACTACT	5100
AATACCCCTA	TCATTCTGTC	AAATAGACAT	AACAAAACCTG	CAACGTGTAG	5150
GTGTATTAGT	CAACAATCCC	AAAGGCATGA	AAATTTTACA	AAATTTTGAA	5200
ACGAGATACC	TGATATTAAG	TCTGATACCC	AAAATAGAGA	ATTCACACTC	5250
ATGTGGGGAT	CAACAGATAA	ACCAATACAA	GAAGTTATTG	GATAGATTGA	5300
TAATTCCTCT	ATATGATGGA	TTAAAATTAC	AAAAAGATGT	AATAGTAGTA	5350
AGTCATGAAA	CCCATAATAA	TACTAATCTT	AGGACAAAAC	GATTCTTTGG	5400
AGAGATAATT	GGGACAATTG	CGATAGGGAT	AGCCACCTCA	GCGCAAATCA	5450
CCGCAGCAGT	CGCTCTTGTC	GAAGCTAAAC	AGGCAAGGTC	AGACATAGAA	5500
AAACTCAAAG	AAGCTATAAG	AGACACAAAC	AAGGCAGTAC	AATCGATTCA	5550
AAGTTCTGTA	GGTAACCTAA	TTGTTGCAGT	TAAATCAGTT	CAAGACTATG	5600
TCAACAATGA	AATTGTACCT	TCAATCACAA	GATTAGGCTG	TGAAGCAGCA	5650
GGGTTACAAT	TGGGAATTGC	ACTGACACAA	CATTACTCAG	AATTAACAAA	5700
TATATTTGGT	GATAATATAG	GAACACTGAA	AGAAAAAGGG	ATAAAATTAC	5750
AGGGGATAGC	ATCGTTATAT	CATACAAACA	TAACAGAAAT	ATTTACTACT	5800
TCAACAGTTG	ACCAATATGA	TATTTATGAC	CTATTATTCA	CTGAATCAAT	5850
CAAGATGAGA	GTGATAGATG	TTGATTTGAG	TGATTACTCA	ATTACTCTTC	5900
AAGTTAGACT	TCCTTTATTA	ACTAAACTAT	CAAATACTCA	GATTTATAAA	5950
GTAGATTCTA	TATCATACAA	CATCCAGGGC	AAAGAGTGGT	ATATTCCTCT	6000
TCCCAATCAC	ATCATGACAA	AAGGGGCTTT	TCTAGGTGGT	GCTGATATTA	6050
AAGAATGCAT	AGAGGCATTC	AGCAGTTATA	TATGTCCTTC	TGATCCAGGT	6100
TATATATTAA	ATCACGAGAT	AGAGAATTGT	TTATCAGGGA	ACATAACACA	6150
GTGTCCTAAG	ACTGTTGTTA	CATCAGATGT	GGTACCACGA	TACGCGTTTG	6200
TGAATGGTGG	ATTAATTGCA	AACTGCATAA	CAACTACATG	TACATGCAAT	6250
GGAATTGACA	ATAGAATTAA	TCAATCACCT	GATCAAGGAA	TTAAGATCAT	6300
AACACATAAA	GAATGCCAGG	TAATAGGTAT	AAACGGAATG	TTATTCAATA	6350
CTAATAGAGA	AGGGACATTA	GCAACTTATA	CATTTGATGA	CATTATATTA	6400
AATAACTCTG	TTGCACTTAA	TCCAATTGAT	ATATCTATGG	AACTTAACAA	6450
GGCAAAACTA	GAATTAGAAG	AATCGAAGGA	ATGGATAAAG	AAATCAAATC	6500
AAAAGTTAGA	TTCCGTTGGA	AGTTGGTATC	AATCTAGTGC	AACAATCACC	6550
ATAATCATAG	TGATGATAAT	AATTCTATTT	ATAATCAATA	TAACAATTAT	6600
TGTAGTCATA	ATCAAATTCT	ATAGAATTAA	GGGGGAAAAT	CAAAACGACA	6650
AAAACAGTGA	GCCGTATATA	CTGACAAATA	GACAATAAGA	CTATACACGA	6700
TCAAATATAG	AAAGTACAAA	AAACTTAGGA	ACAAAGTTGT	TCAACACAGC	6750
AGCAGCGAAC	AGACCCAAAG	GCAGCGCAGA	GGCGACACCG	AACCCAAAAA	6800
TGGAATATTG	GAAACACACA	AACAGCACAA	AAAACACCAA	CAATGAAACC	6850
GAAACAACCA	GAGGCAAACA	CAGTAGCAAG	GTTACAAATA	TCATAATGTA	6900
CACCTTCTGG	ACAATAACAT	CAACAATATT	ATTAGTCATT	TTTATAATGA	6950
TATTGACAAA	CTTAATTCAA	GAGAACAATC	ATAATAAATT	AATGTTGCAG	7000
GAAATAAGAA	AAGAATTTCG	GGCAATAGAC	ACCAAGATTC	AGAGGACCTC	7050

FIG. 7C

GGATGACATT	GGAACCTCAA	TACAGTCAGG	AATAAATACA	AGACTTCTCA	7100
CAATTCAGAG	TCATGTTCAA	AACTATATCC	CACTATCACT	AACACAACAA	7150
ATGTCAGATC	TCAGAAAATT	TATCAATGAT	CTAACAAATA	AAAGAGAACA	7200
TCAAGAAGTG	CCAATACAGA	GAATGACTCA	TGATAGAGGT	ATAGAACCCC	7250
TAAATCCAGA	CAAGTTCTGG	AGGTGTACAT	CTGGTAACCC	ATCTCTAACA	7300
AGTAGTCCTA	AGATAAGGTT	AATACCAGGG	CCAGGTTTAT	TAGCAACATC	7350
TACTACAGTA	AATGGCTGTA	TTAGAATCCC	ATCGTTAGCA	ATCAATCATT	7400
TAATCTACGC	TTACACCTCT	AATCTTATCA	CCCAGGGCTG	TCAAAATATA	7450
GGGAAATCTT	ACCAAGTACT	ACAAATAGGG	ATAATTAATA	TAAATTTCGA	7500
CCTAGTACCT	GATTTAAATC	CCAGAGTCAC	ACATACATTT	AATATTGATG	7550
ATAATAGGAA	ATCTTGCTCT	CTGGCACTAT	TGAATACAGA	TGTTTATCAG	7600
TTATGCTCAA	CACCAAAAGT	TGATGAGAGA	TCCGATTATG	CATCAACAGG	7650
TATTGAGGAT	ATTGTACTTG	ACATTGTCAC	TAATAATGGA	TTAATTATAA	7700
CAACAAGGTT	TACAAATAAT	AATATAACTT	TTGATAAACC	GTATGCAGCA	7750
TTGTATCCAT	CAGTAGGACC	AGGAATCTAT	TATAAGGGTA	AAGTTATATT	7800
TCTCGGATAT	GGAGGTCTAG	AGCATGAAGA	AAACGGAGAC	GTAATATGTA	7850
ATACAACTGG	TTGTCTTGGC	AAAACACAGA	GAGACTGTAA	TCAGGCTTCT	7900
TATAGCCCAT	GGTTCTCAAA	TAGGAGAATG	GTAAACTCTA	TTATTGTTGT	7950
TGATAAAGGC	ATAGATGCAA	CTTTTAGCTT	GAGGGTGTGG	ACTATTCCAA	8000
TGAGCCAAAA	TTATTGGGGA	TCAGAAGGAA	GATTACTTTT	ATTAGGTGAC	8050
AGAATATACA	TATATACTAG	ATCCACAAGT	TGGCACAGTA	AATTACAGTT	8100
AGGGGTAATT	GATATTTCTG	ATTATAATAA	TATAAGAATA	AATTGGACTT	8150
GGCATAATGT	ACTATCACGG	CCAGGAAATG	ATGAATGTCC	ATGGGGTCAT	8200
TCATGCCCAG	ACGGATGTAT	AACAGGAGTT	TACACTGATG	CATATCCGCT	8250
AAACCCATCG	GGGAGTGTTG	TATCATCAGT	AATTCTTGAC	TCACAAAAGT	8300
CTAGAGAAAA	CCCAATCATT	ACCTACTCAA	CAGCTACAAA	TAGAATAAAT	8350
GAATTAGCTA	TATATAACAG	AACACTTCCA	GCTGCATATA	CAACAACAAA	8400
TTGTATCACA	CATTATGATA	AAGGGTATTG	TTTTCATATA	GTAGAAATAA	8450
ATCACAGAAG	TTTGAATACG	TTTCAACCTA	TGTTATTCAA	AACAGAAGTT	8500
CCAAAAAACT	GCAGCTAAAT	TGATCATCGC	ATATCGGATG	CCAGATGACA	8550
TTAAAAAGAGA	CCACCAGACA	GACAACACAG	GAGATGATGC	AAGATATAAA	8600
GGAATAATAA	AAAACCTTAGG	AGAAAAGTGT	GCAAGAAAAA	TGGACACTGA	8650
ATCCCACAGC	GGCACAACAT	CTGACATTCT	GTACCCTGAA	TGTCACCTCA	8700
ATTCTCCTAT	AGTTAAAGGA	AAAATAGCAC	AACTGCATAC	AATAATGAGT	8750
TTGCCCCAAC	CCTACGATAT	GGATGATGAT	TCAATACTGA	TTATTACTAG	8800
ACAAAAAATC	AAACTCAATA	AATTAGATAA	AAGACAACGG	TCAATTAGGA	8850
AATTAAGATC	AGTCTTAATG	GAAAGAGTAA	ATGATCTTGG	TAAATACACC	8900
TTTATCAGAT	ATCCAGAAAT	GTCTAGTGAA	ATGTTCCAAT	TATGTATACC	8950
CGGAATTAAT	AATAAAATAA	ATGAATTGCT	AAGTAAAGCA	AGTAAACAT	9000
ATAATCAAAT	GACTGATGGA	TTAAGAGATC	TATGGGTTAC	TGTACTATCG	9050
AAGTTAGCAT	CGAAAAATGA	TGGAAGTAAT	TATGATATCA	ATGAAGATAT	9100
TAGCAATATA	TCAAATGTTC	ACATGACTTA	CCAATCAGAC	AAATGGTATA	9150
ATCCATTCAA	GACATGGTTT	ACTATTAAGT	ATGACATGAG	GAGATTACAA	9200
AAAGCCAAAA	ATGAGATTAC	ATTCAATAGG	CATAAAGATT	ATAATCTATT	9250
AGAAGACCAA	AAGAATATAT	TGCTGATACA	TCCAGAACTC	GTCTTAATAT	9300
TAGATAAACA	AAATTACAAT	GGGTATATAA	TGACTCCTGA	ATTGGTACTA	9350
ATGTATTGTG	ATGTAGTTGA	AGGGAGGTGG	AATATAAGTT	CATGTGCAAA	9400

FIG. 7D

ATTGGATCCT	AAATTACAAT	CAATGTATTA	TAAAGGTAAC	AATTTATGGG	9450
AAATAATAGA	TGGACTATTC	CTGACCTTAG	GAGAAAGAAC	ATTTGACATA	9500
ATATCACTAT	TAGAACCGCT	TGCATTATCG	CTCATTCAAA	CTCATGACCC	9550
GGTTAAACAG	CTCAGAGGGG	CTTTTTTAAA	TCACGTGTTA	TCAGAAATGG	9600
AATCAATATT	CGCAGCTGAG	TGTACAACAG	AGGAAATACC	TAATGTGGAT	9650
TATATAGATA	AAATTTTAGA	TGTATTCAAA	GAATCAACAA	TAGATGAAAT	9700
AGCAGAAATT	TTCTCTTTCT	TCCGAACCTT	TGGACACCCT	CCATTAGAGG	9750
CGAGTATAGC	AGCAGAGAAA	GTTAGAAAGT	ATATGTACAC	TGAGAAATGT	9800
TTGAAATTTG	ATACTATCAA	TAAATGTCAT	GCTATTTTTT	GTACAATAAT	9850
TATAAATGGA	TATAGAGAAA	GACATGGTGG	TCAATGGCCT	CCAGTTACAT	9900
TACCTATTCA	TGCACATGAA	TTTATCATAA	ATGCGTACGG	ATCAAATTCT	9950
GCCATATCAT	ATGAAAATGC	TGTAGATTAT	TATAAGAGCT	TCATAGGAAT	10000
AAAATTTGAC	AAGTTTATAG	AGCCTCAATT	GGATGAAGAC	TTAACTATTT	10050
ATATGAAAGA	TAAAGCATTA	TCCCCAAAGA	AATCTAACTG	GGACACAGTC	10100
TATCCAGCTT	CAAACCTGTT	ATACCGCACT	AATGTGTCTC	ATGATTACAG	10150
AAGATTGGTT	GAAGTATTTA	TAGCAGATAG	TAAATTTGAT	CCCCACCAAG	10200
TATTAGATTA	CGTAGAATCA	GGATATTGGC	TAGATGATCC	TGAATTTAAT	10250
ATCTCATATA	GTTTAAAAGA	GAAAGAAATA	AAACAAGAAG	GTAGACTTTT	10300
TGCAAAAATG	ACATACAAGA	TGAGAGCTAC	ACAAGTATTA	TCAGAAACAT	10350
TATTGGCGAA	TAATATAGGG	AAATTCTTCC	AAGAGAATGG	GATGGTTAAA	10400
GGAGAAATTG	AATTACTCAA	GAGACTGACA	ACAATATCTA	TGTCTGGGGT	10450
TCCGCGGTAT	AATGAGGTAT	ACAATAATTC	AAAAAGTCAC	ACAGAGGAAC	10500
TTCAAGCTTA	TAATGCAATT	AGCAGTTCCA	ATTTATCTTC	TAATCAGAAG	10550
TCAAAGAAGT	TTGAATTTAA	ATCAACAGAT	ATATACAATG	ATGGATACGA	10600
AACCGTAAGC	TGCTTCTTAA	CGACAGATCT	TAAAAAATAT	TGTTTTAAAT	10650
GGAGGTATGA	ATCAACAGCT	TTATTCCGTG	ATACTTGTA	TCAGATATTT	10700
GGGTTAAAGG	AATTATTTAA	TTGGCTGCAC	CCTCGCCTTG	AAAAGAGTAC	10750
AATATATGTT	GGAGATCCTT	ATTGCCCGCC	ATCAGATATT	GAACATTTAC	10800
CACTTGATGA	CCATCCTGAT	TCAGGATTTT	ATGTTCATAA	TCCTAAAGGA	10850
GGAATAGAAG	GGTTTTGCCA	AAAGTTATGG	ACACTCATAT	CTATCAGTGC	10900
CATACATTTA	GCAGCTGTCA	AAATCGGTGT	AAGAGTTACT	GCAATGGTTC	10950
AAGGGGATAA	TCAAGCCATA	GCTGTTACCA	CCAGAGTACC	TAATAATTAT	11000
GATTATAAGG	TTAAGAAAGA	GATTGTTTAT	AAAGATGTGG	TAAGATTTTT	11050
TGATTCTTTG	AGAGAGGTTA	TGGATGATCT	GGGTCATGAG	CTCAAACATA	11100
ATGAAACTAT	AATAAGTAGT	AAAATGTTTA	TATATAGCAA	AAGGATATAC	11150
TATGACGGAA	GAATCCTTCC	TCAGGCGTTA	AAAGCATTGT	CTAGATGTGT	11200
TTTTTGGTCT	GAAACAATCA	TAGATGAGAC	AAGATCAGCA	TCCTCAAATC	11250
TGGCGACATC	GTTTGCAAAG	GCCATTGAGA	ATGGCTACTC	ACCTGTATTG	11300
GGATATGTAT	GCTCAATCTT	CAAAAATATC	CAACAGTTGT	ATATAGCACT	11350
TGGAATGAAT	ATAAATCCAA	CTATAACCCA	AAATATTAAA	GATCAATATT	11400
TCAGGAATAT	TCATTGGATG	CAATATGCAT	CTCTAATCCC	TGCTAGTGTC	11450
GGAGGATTTA	ATTATATGGC	CATGTCAAGG	TGTTTTGTCA	GAAACATTGG	11500
AGATCCTACA	GTCGCTGCAT	TAGCTGATAT	TAAAAGATTT	ATAAAAGCAA	11550
ATTTGTTAGA	TCGAGGTGTC	CTTTACAGAA	TTATGAATCA	GGAACCAGGC	11600
GAGTCCTCCT	TTTTAGACTG	GGCTTCAGAC	CCCTATTCAT	GTAACCTACC	11650
ACAATCTCAA	AATATAACCA	CCATGATAAA	GAATATAACT	GCAAGAAATG	11700
TACTACAGGA	CTCACCAAAC	CCATTACTAT	CTGGATTATT	TACAAGTACA	11750

FIG. 7E

ATGATAGAAG	AGGATGAGGA	ATTAGCTGAG	TTCCTAATGG	ACAGGAGAAT	11800
AATTCTCCCA	AGGGTTGCGC	ATGACATTTT	AGATAATTCT	CTTACTGGAA	11850
TTAGGAATGC	TATAGCTGGT	ATGTTGGATA	CAACAAAATC	ACTAATTCGA	11900
GTAGGGATAA	ACAGAGGAGG	ATTAACCTAT	AACTTATTAA	GAAAGATAAG	11950
CAACTATGAT	CTTGTACAAT	ATGAGACACT	TAGTAAAAC	TTAAGACTAA	12000
TAGTCAGTGA	CAAGATTAAG	TATGAAGATA	TGTGCTCAGT	AGACCTAGCC	12050
ATATCATTA	GACAAAAAAT	GTGGATGCAT	TTATCAGGAG	GAAGAATGAT	12100
AAATGGACTT	GAAACTCCAG	ATCCTTTAGA	GTTACTGTCT	GGAGTAATAA	12150
TAACAGGATC	TGAGCATTGT	AGGATATGTT	ATTCAACTGA	AGGTGAAAGC	12200
CCATATACAT	GGATGTATTT	ACCAGGCAAT	CTTAATATAG	GATCAGCTGA	12250
AACAGGAATA	GCATCATTA	GGGTCCCTTA	CTTTGGATCA	GTTACGGATG	12300
AGAGATCTGA	AGCACAAATG	GGGTATATCA	AAAATCTAAG	CAAACCAGCT	12350
AAGGCTGCTA	TAAGAATAGC	AATGATATAT	ACTTGGGCAT	TTGGGAATGA	12400
CGAAATATCT	TGGATGGAAG	CATCACAGAT	TGCACAAACA	CGTGCGAACT	12450
TTACATTAGA	TAGCTTAAAG	ATTTTGACAC	CAGTGACAAC	ATCAACAAAT	12500
CTATCACATA	GGTTAAAAGA	TACTGCTACT	CAGATGAAAT	TTTCTAGTAC	12550
ATCACTTATT	AGAGTAAGCA	GGTTCATCAC	AATATCTAAT	GATAATATGT	12600
CTATTAAAGA	GGCAAATGAA	ACTAAAGATA	CAAATCTTAT	TTATCAACAG	12650
GTAATGTAA	CAGGGTTAAG	TGTATTTGAA	TATCTATTTA	GGTTAGAGGA	12700
GAGTACAGGA	CATAACCTTA	TGGTCATGCA	TCTACATATA	GAGGATGGAT	12750
GTTGTATCAA	AGAGAGTTAC	AATGATGAGC	ATATCAATCC	GGAGTCTACA	12800
TTAGAGTTAA	TTAAATACCC	TGAGAGTAAT	GAATTTATAT	ATGATAAGGA	12850
CCCTTTAAAG	GATATAGATC	TATCAAAATT	AATGGTTATA	AGAGATCATT	12900
CTTATACAAT	TGACATGAAT	TACTGGGACG	ACACAGATAT	TGTACATGCA	12950
ATATCAATAT	GTAATGAGT	TACAATAGCA	GATACAATGT	CGCAGCTAGA	13000
TCGGGATAAT	CTTAAGGAGC	TGGTTGTAAT	TGCAAATGAT	GATGATATTA	13050
ACAGTCTGAT	AACTGAATTT	CTGACCCTAG	ATATACTAGT	GTTTCTCAAA	13100
ACATTTGGAG	GGTTACTCGT	GAATCAATTT	GCATATACCC	TTTATGGATT	13150
GAAAATAGAA	GGAAGGGATC	CCATTTGGGA	TTATATAATG	AGAACATTAA	13200
AAGACACCTC	ACATTCAGTA	CTTAAAGTAT	TATCTAATGC	ACTATCTCAT	13250
CCAAAAGTGT	TTAAGAGATT	TTGGGATTGT	GGAGTTTTGA	ATCCTATTTA	13300
TGGTCCTAAT	ACTGCTAGTC	AGGACCAAGT	TAAGCTTGCT	CTCTCAATTT	13350
GCGAGTACTC	CTTGGATCTA	TTTATGAGAG	AATGGCTGAA	TGGAGCATCA	13400
CTTGAGATCT	ATATCTGTGA	TAGTGACATG	GAAATAGCAA	ATGATAGAAG	13450
ACAAGCATTT	CTCTCAAGAC	ACCTTGCCCT	TGTGTGTTGT	TTAGCAGAGA	13500
TAGCATCTTT	TGGACCAAAT	TTATTAAATC	TAACATATCT	AGAGAGACTT	13550
GACGAATTAA	AACAATACTT	GGATCTGAAC	ATCAAAGAAG	ATCCTACTCT	13600
TAAATATGTG	CAAGTATCAG	GACTGTTAAT	TAAATCATTC	CCCTCAACTG	13650
TTACGTATGT	GAGGAAAAC	GCGATTAAGT	ATCTGAGGAT	TCGTGGCATT	13700
AATCCGCCTG	AAACGATTGA	AGATTGGGAT	CCCATAGAAG	ATGAGAATAT	13750
CTTAGACAAT	ATTGTTAAAA	CTGTAAATGA	CAATTGCAGT	GATAATCAAA	13800
AGAGAAATAA	AAGTAGTTAT	TTCTGGGGAT	TAGCTCTAAA	GAATTATCAA	13850
GTCGTAAAAA	TAAGATCCAT	AACGAGTGAT	TCTGAAGTTA	ATGAAGCTTC	13900
GAATGTTACT	ACACATGGAA	TGACACTTCC	TCAGGGAGGA	AGTTATCTAT	13950
CACATCAGCT	GAGGTTATTT	GGAGTAAACA	GTACAAGTTG	TCTGAAAGCT	14000
CTTGAATTGT	CACAAATTTT	AATGAGGGAA	GTTAAAAAAG	ATAAAGATAG	14050
ACTCTTTTTA	GGAGAAGGAG	CAGGAGCTAT	GTTAGCATGT	TATGATGCTA	14100

FIG. 7F

CACTCGGTCC	TGCAATAAAT	TATTACAATT	CTGGTTTAAA	TATTACAGAT	14150
GTAATTGGTC	AACGGGAATT	AAAAATCTTC	CCATCAGAAG	TATCATTAGT	14200
AGGTAAAAAA	CTAGGAAATG	TAACACAGAT	TCTTAATCGG	GTGAGGGTGT	14250
TATTTAATGG	GAATCCCAAT	TCAACATGGA	TAGGAAATAT	GGAATGTGAG	14300
AGTTTAATAT	GGAGTGAATT	AAATGATAAG	TCAATTGGTT	TAGTACATTG	14350
TGACATGGAG	GGAGCAATAG	GCAAATCAGA	AGAAACTGTT	TTACATGAAC	14400
ATTATAGTAT	TATTAGGATT	ACATATTTAA	TTGGGGATGA	TGATGTTGTT	14450
CTAGTATCAA	AAATTATACC	AACTATTACT	CCGAATTGGT	CTAAAATACT	14500
CTATCTATAC	AGGTTGTATT	GGAAGGATGT	GAGTGTAGTG	TCCCTTAAAA	14550
CATCCAATCC	TGCCTCAACA	GAGCTTTATT	TAATTTCAA	GGATGCTTAC	14600
TGTACTGTAA	TGGAACCCAG	TAATCTTGTT	TTATCAAAAC	TTAAAAGGAT	14650
ATCATCAGTA	GAAGAAAATA	ATCTATTAAA	ATGGATAATC	TTATCAAAAA	14700
GGAAGAACAA	CGAATGGTTA	CAGCATGAAA	TCAAAGAAGG	AGAAAGGGAT	14750
TATGGGATAA	TGAGGCCATA	TCATACAGCA	CTGCAAATTT	TTGGATTCCA	14800
AATTAECTTA	AATCACTTAG	CTAAAGAATT	TTTATCAACT	CCTGATTTAA	14850
CCAACATTAA	TAATATAATT	CAAAGTTTTA	CAAGAACAAT	TAAAGATGTT	14900
ATGTTCGAAT	GGGTCAATAT	CACTCATGAC	AATAAAAGAC	ATAAATTAGG	14950
AGGAAGATAT	AATCTATTCC	CGCTTAAAA	TAAGGGGAAG	TTAAGATTAC	15000
TATCACGAAG	ATTAGTACTA	AGCTGGATAT	CATTATCTTT	ATCAACCAGA	15050
TTACTGACAG	GCCGTTTCCC	AGATGAAAA	TTTGAAAATA	GGGCACAGAC	15100
CGGATATGTA	TCATTGGCTG	ATACTGATTT	AGAATCTTTA	AAGTTATTAT	15150
CAAGAAATAT	TGTCAAAAAGT	TACAAAGAAC	ACATAGGATT	AATATCATAC	15200
TGGTTTTTTAA	CCAAAGAGGT	CAAATACTA	ATGAAACTTA	TAGGGGGAGT	15250
CAAATACTA	GGAATTCCCA	AACAGTACAA	AGAGTTAGAG	GATCGATCAT	15300
TTCAGGGTTA	TGAATATGAT	AATGAATTTG	ATATTGATTA	ATACATAAAA	15350
ACAAAAAATA	AAACACCTAA	TCCTCTCCCA	TTCACCTCCA	ACAAAATGAA	15400
AAGTAAGAAA	AACATATAAT	ATACATATAC	CAAACAGAGT	TTTTCTCTTG	15450
TTTGGT					15456

FIG. 7G

Figure 8A

rHPiV3						
	3	N	P _{cov}	M	F	L
					HN	
						5
rHPiV3-FgHNB						
	3	N	P _{cov}	M	F	L
					HN	
						5
rBPiV3-FgHNB						
	3	N	P _{cov}	M	F	L
					HN	
						5
BPiV3 Ka						
	3	N	P _{cov}	M	F	L
					HN	
						5

Figure 8B

Assembly of an antigenomic cDNA for BPIV3 Ka






Figure 8C

Generation of full length cDNA clones encoding HPIV3/BPIV3 antigenic chimeric viruses

1. Generation of HPIV3 and BPIV3 full length clones



2. Mutagenesis to create uniqueSgrAI and BsiWI restriction sites



3. Cloning of the F and HN genes into the heterologous full length cDNA






Figure 9







Figure 10

Figure 11



SEQUENCE LISTING

<110> THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as

<120> ATTENUATED HUMAN-BOVINE CHIMERIC PARAINFLUENZA VIRUS
(PIV) VACCINES

<130> 15280-399100PC

<140>

<141>

<150> 60/143,134

<151> 1999-07-09

<160> 31

<170> PatentIn Ver. 2.1

<210> 1

<211> 7

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sequence
flanking sites of sequence polymorphism in BPIV3
Ka

<400> 1

actggtt

7

<210> 2

<211> 12

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sequence
flanking site for introduction of Sgr A1 site for
rBPIV3 Ka

<400> 2

tccaacattg ca

12

<210> 3

<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking site for introduction of Sgr A1 site for
rBPIV3 s

<400> 3
tccaccggtg ca

12

<210> 4
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking site for introduction of Sgr A1 site for
rHPIV3 JS

<400> 4
tccaccggtg ca

12

<210> 5
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking site for introduction of Sgr A1 site for
rHPIV3 s

<400> 5
cgcaccggtg ta

12

<210> 6
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence

flanking site for introduction of Bsi W1 site for
rBPIV3 Ka

<400> 6
gatataaaga 10

<210> 7
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking site for introduction of Bsi W1 site for
rBPIV3 s

<400> 7
gacgtacgga 10

<210> 8
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking site for introduction of Bsi W1 site for
rHPIV3 JS

<400> 8
gacaaaaggg 10

<210> 9
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking site for introduction of Bsi W1 site for
rHPIV3 s

<400> 9
gacgtacggg 10

<210> 10
<211> 11
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking N gene start codon

<400> 10
caaaaatggt g 11

<210> 11
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking N gene stop codon

<400> 11
gcaactaatc ga 12

<210> 12
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking introduced restriction site at N gene
start codon

<400> 12
taaccatggt ga 12

<210> 13
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence

WO 01/04320

flanking introduced restriction site at N gene
stop codon

<400> 13
gcacttaagc ac

12

<210> 14
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking mutation to restore context for N gene
start codon

<400> 14
caaaaatggt ga

12

<210> 15
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking mutation to restore context for N gene
stop codon

<400> 15
gcaactagtc ga

12

<210> 16
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking N gene start codon in rJS

<400> 16
ggaactctat aatttcaaaa atgttgagcc tatttgatac

40

<210> 17
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking N gene start codon in cKa and cSF

<400> 17
ggaactctat aatttcaaaa atgttgagtc tattcgacac 40

<210> 18
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking N gene start codon in Ka and SF

<400> 18
gaaatcctaa gactgtaatc atgttgagtc tattcgacac 40

<210> 19
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking N gene stop codon in rJS

<400> 19
ttaacgcatt tggaagcaac taatcgaatc aacattttaa 40

<210> 20
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking N gene stop codon in cKa and cSF

<400> 20

tcagtgcatt cggaagcaac tagtcgaatc aacatttttaa

40

<210> 21

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sequence
flanking N gene stop codon in Ka and SF

<400> 21

tcagtgcatt cggaagcaac tagtcacaaa gagatgacca

40

<210> 22

<211> 15456

<212> DNA

<213> Bovine Parainfluenza Virus 3 (Ka strain)

<400> 22

accaaacaag agaagagact ggtttgggaa tattaattca aataaaaatt aacttaggat 60
 taaagaactt taccgaaagg taaggggaaa gaaatcctaa gactgtaatc atgttgagtc 120
 tattcgacac attcagtgcg cgtaggcagg agaacataac gaaatcagct ggtggggctg 180
 ttattcccgg gcaaaaaaac actgtgtcta ttttgtctca ttcttttagac aatgaaaagc 240
 acaatgataa aatgacattg gctcttctct ttttgtctca ttcttttagac aatgaaaagc 300
 agcatgcgca aagagctgga tttttagttt ctctgttata aatggcttat gccaacccag 360
 aattatattt aacatcaaataa ggtagtaata cagatgttaa atatgttata tacatgatag 420
 agaaagaccc aggaagacag aaatatgggt ggtttgtcgt caagactaga gagatgggtt 480
 atgaaaagac aactgattgg atgttcggga gtgatcttga gtatgatcaa gacaatatgt 540
 tgcaaaatgg tagaagcact tctacaatcg aggatcttgt tcatactttt ggatatccat 600
 cgtgtcttgg agcccttata atccaagttt ggataatact tgtaaggctt ataaccagta 660
 tatcaggatt gaggaagga ttctttactc ggtagaagc atttcgacaa gatggaacag 720
 ttaaattccag tctagtgttg agcgtgatg cagtagaaca aattggatca attatgaggt 780
 cccaacagag cttggtaaca ctcatgggtg aaacactgat aacaatgaac acaggcagga 840
 atgatctgac aacaatagaa aagaatatac agattgtagg aaactacatc agagatgcag 900
 gtcttgcttc atttttcaac acaatcagat atggcattga gactagaatg gcagctctaa 960
 ctctgtctac ccttagaccg gatatacaaa gactcaaggc actgatcgag ttatatctat 1020
 caaagggggc acgtgctcct tttatatgca ttttgagaga tcccgatcat ggtgagtttg 1080
 caccaggcaa ctatcctgcc ctctggagtt atgcgatggg tgtagcagtt gtacaaaaca 1140
 aggccatgca acagtatgta acaggaaggt cttatctgga tattgaaatg ttccaacttg 1200
 gtcaagcagt ggcacgtgat gccgagtcgc agatgagttc aatattagag gatgaactgg 1260
 gggtcacaca agaagccaag caaagcttga agaacacat gaagaacatc agcagttcag 1320
 atacaacctt tcataagcct acagggggat cagccataga aatggcgata gatgaagaag 1380
 cagggcagcc tgaatccaga ggagatcagg atcaaggaga tgagcctcgg tcatccatag 1440
 ttccttatgc atgggcagac gaaaccggga atgacaatca aactgaatca actacagaaa 1500

ttgacagcat	caaaaactgaa	caaagaaaca	tcagagacag	gctgaacaaa	agactcaacg	1560
agaaaaggaa	acagagtgc	ccgagatcaa	ctgacatcac	aaacaacaca	aatcaaaactg	1620
aaatagatga	ttgtttcagt	gcatttcggaa	gcaactagtc	acaaagagat	gaccactatc	1680
accagcaaca	agtaagaaaa	acttaggatt	aatggaaatt	atccaatcca	gagacggaag	1740
gacaaatcca	gaatccaacc	acaactcaat	caaccaaaga	ttcatggaag	acaatgttca	1800
aaacaatcaa	atcatggatt	cttgggaaga	gggatcagga	gataaatcat	ctgacatctc	1860
atcggccctc	gacatcattg	aattcatact	cagcaccgac	tcccaagaga	acacggcaga	1920
cagcaatgaa	atcaacacag	gaaccacaag	acttagcacg	acaatctacc	aacctgaatc	1980
caaaaacaaca	gaaacaagca	aggaaaatag	tggaccagct	aacaaaaatc	gacagtttgg	2040
ggcatcacac	gaacgtgcc	cagagacaaa	agatagaaat	gttaatcagg	agactgtaca	2100
gggaggatat	aggagaggaa	gcagcccaga	tagtagaact	gagactatgg	tactcgaag	2160
aatctccaga	agcagcccag	atcctaacaa	tggaaaccaa	atccaggaag	atattgatta	2220
caatgaagtt	ggagagatgg	ataaggactc	tactaagagg	gaaatgcgac	aatttaaaga	2280
tgttccagtc	aaggtatcag	gaagtgatgc	cattcctcca	acaaaacaag	atggagacgg	2340
tgatgatgga	agaggcctgg	aatctatcag	tacatttgat	tcaggatata	ccagtatagt	2400
gactgcccga	acactagatg	acgaagaaga	actccttatg	aagaacaaca	ggccaagaaa	2460
gtatcaatca	acaccccaga	acagtgcaca	gggaattaaa	aaaggggttg	gaaggccaaa	2520
agacacagac	aaacaatcat	caatattgga	ctacgaactc	aacttcaaag	gatcgaagaa	2580
gagccagaaa	atcctcaaag	ccagcacgaa	tacaggagaa	ccaacaagac	cacagaatgg	2640
atcccagggg	aagagaatca	catcctggaa	catcctcaac	agcgagagcg	gcaatcgaac	2700
agaatcaaca	aaccaaaccc	atcagacatc	aacctcgga	cagaaccaca	caatgggacc	2760
aagcagaaca	acctccgaac	caaggatcaa	gacacaaaag	acggatggaa	aggaaagaga	2820
ggacacagaa	gagagcactc	gatttacaga	aagggcgatt	acattattac	agaatcttgg	2880
tgtaatccaa	tctgcagcaa	aattagacct	ataccaagac	aagagagttg	tgtgtgtggc	2940
gaatgtccta	aacaatgcag	atactgcatc	aaagatagac	ttcctagcag	gtttgatgat	3000
aggagtgtca	atggatcatg	ataccaaatt	aaatcagatt	cagaacgaga	tattaagttt	3060
gaaaactgat	cttaaaaaga	tggatgaatc	acatagaaga	ctaattgaga	atcaaaaaga	3120
acaattatca	ctgatcacat	cattaatctc	aaatcttaaa	attatgacag	agagaggagg	3180
gaagaaggac	caaccagaac	ctagcgggag	gacatccatg	atcaagacaa	aagcaaaaaga	3240
agagaaaata	aagaaagtca	ggtttgaccc	tcttatggaa	acacagggca	tcgagaaaaa	3300
catccctgac	ctctatagat	caatagagaa	aacaccagaa	aacgacacac	agatcaaatc	3360
agaaaataaac	agattgaatg	atgaatccaa	tgccactaga	ttagtaccta	gaagaataag	3420
cagtacaatg	agatcattaa	taataatcat	taacaacagc	aattttatcat	caaaagcaaa	3480
gcaatcatac	atcaacgaac	tcaagctctg	caagagtgc	gaggaagtgt	ctgagttgat	3540
ggacatgttc	aatgaggatg	tcagctccca	gtaaaccgcc	aaccaagggt	caacaccaag	3600
aaaaccaata	gcacaaaaca	gccaatcaga	gaccaccca	atacaccaaa	ccaatcaaca	3660
cataacaaaag	atctccagat	catagatgat	taagaaaaac	ttaggatgaa	aggactaatc	3720
aatcctccga	aacaatgagc	atcaccaact	ccacaatcta	cacattcca	gaatcctctt	3780
tctccgagaa	tggcaacata	gagccgttac	cactcaaggt	caatgaacag	agaaaggcca	3840
tacctcatat	tagggttgtc	aagataggag	atccgcccaa	acatggatcc	agatatctgg	3900
atgtcttttt	actgggcttc	tttgagatgg	aaaggtcaaa	agacagggtat	gggagcataa	3960
gtgatctaga	tgatgatcca	agttacaagg	tttgtggctc	tggatcattg	ccacttgggt	4020
tggctagata	caccggaaat	gatcaggaac	tctacaggc	tgcaaccaag	ctcgatatag	4080
aagtaagaag	aactgtaaag	gctacggaga	tgatagttta	cactgtacaa	aacatcaaac	4140
ctgaactata	tccatggtcc	agtagattaa	gaaaagggat	gttatttgac	gctaataaag	4200
ttgcacttgc	tcttcaatgt	cttccactag	atagagggat	aaaattcagg	gtgatatttg	4260
tgaactgcac	agcaattgga	tcaataaactc	tattcaaaat	ccctaagtcc	atggcattgt	4320
tatcattgcc	taatacaata	tcaataaatc	tacaagtaca	tatcaaaaaca	ggagttcaga	4380

```

cagattccaa aggagtagtt cagattctag atgaaaaagg tgaaaaatca ctaaatttca 4440
tggttcattc cgggttgatc aaaaggaaga tgggcagaat gtactcagtt gaattattgta 4500
agcagaagat cgagaagatg agattattat tctcattggg attagttgga gggatcagct 4560
tccacgtcaa cgcaactggc tctatatcaa agacattagc aagtcaatta gcattcaaaa 4620
gagaaatctg ctatccccta atggatctga atccacactt aaattcagtt atatgggcat 4680
catcagttga aattacaagg gtagatgcag ttctccagcc ttcattacct ggcgaattca 4740
gatactaccc aaacatcata gcaaaagggg tggggaaaat cagacagtaa aatcaacaac 4800
cctgatatcc aacattgcaa atcaggctac ccacaggaga aaaatcaaaa acttaggatc 4860
aaagggatca ccacgaaccc cggaaaacag ccaaacaaac caacacacaa atcacagaca 4920
aaaaggagaa ggcactgcaa agaccgagaa aaaacagaac gcacacaacc aagcagagaa 4980
aagccaaagc ccgccattca caaacacacc aacaatcctg caaacaagca ccaaaacaga 5040
ggtcaaaaga caaagagcac cagatatgac catcacaacc acaatcatag ccatattact 5100
aataccccca tcattttgtc aaatagacat aacaaaactg caacgtgtag gtgtgttagt 5160
caacaatcct aaaggcatga agatttcaca aaatttcgaa acgagatacc tgatattaag 5220
tttgataccc aaaatagaga attcacactc atgtggggat caacagataa accaatataa 5280
gaagttattg gatagattga taattcctct atatgatgga ttaaaattac aaaaagatgt 5340
aatagtagta agtcatgaaa ccacacaaca tactaatctt aggacaaaac gattctttgg 5400
agagataatt gggacaattg cgatagggat agccacttca gcacaaatca ccgcagcagt 5460
cgctcttgtc gaagctaaac aggcmaaagtc agacatagaa aaactcaaag aggcataaag 5520
agacacaaac aaggcagtac aatcgattca aagttctgta ggtaacctaa ttgttgagc 5580
taaatacagtt caagactatg tcaacaatga aattatacct tcaatcacia gattaggctg 5640
tgaagcagca gggttacaat tgggaattgc attgacacia cattactcag aattaacaaa 5700
tatatttggt gataatatag gaacactgaa agaaaaaggg ataaaattac aagggatagc 5760
atcattatat cacacaaaca taacggaaat atttactact tcaacagttg accaatatga 5820
tatattatgac ctattattca ctgagtcaat caagatgaga gtgatagatg ttgatttgag 5880
tgattactca attactcttc aagttagact tcctttatta actaaactat caaatactca 5940
aatattataa gtagattcta tatcatacaa catccagggc aaagagtggg atattcctct 6000
tcccaatcac atcatgacaa aaggggcttt tctaggtggg gctgatatta aagaatgcat 6060
agaggcattc agcagttata tatgtccttc tgatccaggt tacatatata atcacgagat 6120
agagaattgt ttatcaggga acataacaca gtgtcctaag actgttgta catcagatgt 6180
ggtagcacga tacgcgtttg tgaatgggtg attaatgca aactgcataa caactacatg 6240
tacatgcaat ggaattgaca atagaattaa tcaatcacct gatcaaggaa ttaagatcat 6300
aacacataaa gaatgccag taataggtat aaacggaatg ttattcaata ctaatagaga 6360
agggacatta gcaacttata cttttgatga catcatatta aataactctg ttgcacttaa 6420
tccaattgat atatctatgg aactcaacaa ggcaaaacta gaattagaag aatcgaagga 6480
atggataaag aaatcaaatac aaaagttaga ttccgttgga agttggatc aatctagtgc 6540
aacaatcacc ataatacatg tgatgataat aattctagtt ataataata taacaattat 6600
tgtagtcata atcaaattcc atagaattca ggggaaagat caaaacgaca aaaacagtga 6660
gccgtatata ctgacaaata gacaataaga ctatacacga tcaaatataa aaagtacaaa 6720
aaacttagga acaaagttgt tcaacacagc agcaccgaat agaccaaag gcagcgagga 6780
ggcgacacca aactcaaaaa tggaatattg gaaacacaca aacagcataa ataacaccaa 6840
caatgaaacc gaaacagcca gaggcaaaca tagtagcaag gttacaaata tcataatgta 6900
caccttctgg acaataacat taacaatatt atcagtcatt tttataatga tattgacaaa 6960
cttaattcaa gagaacaatc ataataaatt aatgttgag gaaataagaa aagaattcgc 7020
ggcaatagac accaagattc agaggacttc gtaggacatt ggaacctcaa tacagtcagg 7080
aataaatata agacttctca caattcagag tcatgttcaa aactatatcc cactatcatt 7140
aacacaacaa atgtcagatc tcagaaaatt tatcaatgat ctaacaaata aaagagaaca 7200
tcaagaagtg ccaatacaga gaatgactca tgatagaggt atagaacccc taaatccaaa 7260

```

```

caagttcttg aggtgtacat ctggttaaccc atctctaaca agtagtccta agataaggtt 7320
aataccagga ccaggtttat tagcaacatc tactacagta aatggctgta ttagaattcc 7380
atcggttagta atcaatcatc taatctatgc ttacacctct aatcttatta cccagggctg 7440
tcaagatata gggaaatctt accaagtact acaaataggg ataattacta taaattcggg 7500
cctagtacct gatttaaacc ccagagtcac acatacattt aatattgatg ataatagaag 7560
atcttgctct ctggcactat tgaatacaga tgtttatcag ttatgctcaa caccaaaagt 7620
tgatgaaaga tccgattatg catcaacagg tattgaggat attgtacttg acattgtcac 7680
taataatgga ttaattataa caacaagggt tacaataat aatataactt ttgataaacc 7740
gtatgcagca ttgtatccat cagtgggacc aggaatctat tataaggata aagttatatt 7800
tctcggatat ggaggtctag agcatgaaga aaacggagac gtaatatgta atacaactgg 7860
ttgtcctggc aaaacacaga gagactgtaa tcaggcttct tatagcccat ggttctcaaa 7920
taggagaatg gtaaactcta ttattgttgt tgataaaggc atagatgcaa cttttagctt 7980
gagggtgtgg actattccaa tgagccaaaa ttattgggga tcagaaggaa gattactttt 8040
attagtgac agaatacata tatatactag atccacaagt tggcacagta aattacagtt 8100
aggggtaatt gatatttctg attatactaa tataagaata aattggactt ggcataatgt 8160
actatcacgg ccagggaatg atgaatgtcc atggggtcac tcatgcccag acggatgtat 8220
aacaggagtt tactctgat catatccgct aaacccatcg gggagtgttg tatcatcagt 8280
aattcttgat tcacaaaagt ctagagaaaa cccaatcatt acttactcaa cagctacaaa 8340
tagaataaat gaattagcta tatataacag aacacttcca gctgcatata caacaacaaa 8400
ttgtatcaca cattatgata aagggtattg ttttcatata gtagaaataa atcacagaag 8460
tttgaatacg tttcaaccta tgttattcaa aacagaagtt ccaaaaaact gcagctaaat 8520
tgatcatcgc atatcggatg caagatgaca ttaaaagaga ccaccagaca gacaacacag 8580
gagacgatgc aagatatataa gaaataataa aaaacttagg agaaaagtgt gcaagaaaaa 8640
tggaacaccg gtcccacagc ggcacaacat ctgacattct gtaccctgaa tgtcacctca 8700
attctcctat agttaagga aagatagcac aactgcatac aataatgagt ttgctcagc 8760
cctacgatat ggatgatgat tcaatactga ttattactag acaaaaaatt aaactcaata 8820
aattagataa aagacaacgg tcaattagga aattaagatc agtcttaatg gaaagagtaa 8880
gtgatctagg taaatatacc tttatcagat atccagagat gtctagttaa atgttccaat 8940
tatgtatacc cggaaattaat aataaaataa atgaattgct aagtaaagca agtaaaacat 9000
ataatcaaat gactgatgga ttaagagatc tatgggttac tatactatcg aagttagcat 9060
cgaaaaatga tggaagtaat tatgatatca atgaagatat tagcaatata tcaaatgttc 9120
acatgactta tcaatcagac aaatgggtata atccattcaa gacatggttt actattaagt 9180
atgacatgag aagattacaa aaagccaaaa atgagattac attcaatagg cataaagatt 9240
ataatctatt agaagaccaa aagaatatat tgctgataca tccagaactc gtcttaatat 9300
tagataaaca aaattacaat gggatatataa tgactcctga attggtacta atgtattgtg 9360
atgtagttga agggaggtgg aatataagtt catgtgcaa attggatcct aagttacaat 9420
caatgtatta taagggtaac aatttatggg aaataataga tggactattc tcgaccttag 9480
gagaaagaac atttgacata atatcactat tagaaccact tgcattatcg ctcatcaca 9540
cttatgacct ggttaaaccag ctacaggggg ctttttttaa tcacgtgtta tcagaaatgg 9600
aattaatatt tgcagctgag tgtacaacag aggaaatacc taatgtggat tatatagata 9660
aaattttaga tgtgttcaaa gaatcaacaa tagatgaaat agcagaaatt ttctctttct 9720
tccgaacttt tggacaccct ccattagagg cgagtatagc agcagagaaa gttagaaagt 9780
atatgtatac tgagaaatgc ttgaaatttg atactatcaa taaatgtcat gctatttttt 9840
gtacaataat tataaatgga tatagagaaa gacatgggtg tcaatggcct ccagttacat 9900
tacctgtcca tgcacatgaa tttatcataa atgcatacgg atcaaattct gccatatcat 9960
atgagaatgc ttagattat tataagagct tcataggaat aaaatttgac aagtttatag 10020
agcctcaatt ggatgaagac ttaactatct atatgaaaga taaagcatta tcccaaaga 10080
aatcaaactg ggacacagtc tatccagctt caaacctgtt ataccgcact aatgtgtctc 10140

```



```

atgattcacg aagattgggt gaagtattta tagcagatag taaatttgat cccaccaaag 10200
tattagatta cgtagaatca ggatattggc tggatgatcc tgaatttaat atctcatata 10260
gtttaaaaga gaaagaaata aaacaagaag gtagactttt tgcaaaaatg acatacaaga 10320
tgagggttac acaagtatta tcagaaacat tattggcgaa taatataggg aaattcttcc 10380
aagagaatgg gatggttaaa ggagaaattg aattactcaa gagactaaca acaatatcta 10440
tgtctggagt tccgcggtat aatgaggtat acaataattc aaaaagtcac acagaagaac 10500
ttcaagctta taatgcaatt agcagttcca atttatcttc taatcagaag tcaaagaagt 10560
ttgaatttaa atctacagat atatacaatg atggatacga aaccgtaagc tgcttcttaa 10620
cgacagatct taaaaaatat tgtttaaatt ggaggtatga atcaacagct ttattcggtg 10680
atacttgtaa tcagatattt gggttaaagg aattatttaa ttggctgcac cctgccttg 10740
aaaagagtac aatatatgtt ggagatcctt attgcccgcc atcagatatt gaacatttac 10800
cacttgatga ccacctgat tcaggatttt atgttcataa tcctaaagga ggaatagaag 10860
ggttttgccca aaagtattgg acactcatat ctatcagtgc aatacattta gcagctgtca 10920
aaatcggtgt aagagttact gcaatggttc aaggggataa tcaagccata gctgttacca 10980
caagagtacc taataattat gattataaag ttaagaaaga gattgtttat aaagatgtgg 11040
taagattttt tgattccttg agagaggtga tggatgatct gggcatgag ctcaaactaa 11100
atgaaactat aataagtagt aaaatgttta tatatagcaa aaggatatac tatgacggaa 11160
gaatccttcc tcaggcatta aaagcattgt ctagatgtgt tttttggtct gaaacaatca 11220
tagatgagac aagatcagca tcctcaaatc tggctacatc gtttgcaag gccattgaga 11280
atggctactc acctgtattg ggatatgtat gctcaatctt caaaaatata caacagttgt 11340
atatagcgct tggaatgaat ataaacccaa ctataaccca aaatattaaa gatcaatatt 11400
tcaggaatat tcattggatg caatatgcct ccttaatccc tgctagtgtc ggaggattta 11460
attatatggc catgtcaagg tgttttgtca gaaacattgg agatcctaca gtcgctgcgt 11520
tagccgatat taaaagattt ataaaagcaa atttgttaga tcgaggtgtc ctttacagaa 11580
ttatgaatca agaaccaggc gagtcttctt ttttagactg ggcctcagat ccctattcat 11640
gtaacttacc acaatctcaa aatataacca ccatgataaa gaatataact gcaagaaatg 11700
tactacagga ctacccaaac ccattactat ctggattatt tacaagtaca atgatagaag 11760
aggatgagga attagctgag ttcctaatgg acaggagaat aatcctccca agagttgcac 11820
atgacatttt agataattct cttactggaa ttaggaatgc tatagctggg atgttgata 11880
caacaaaatc actaattcga gtagggataa gcagaggagg attaacctat aactatttaa 11940
gaaagataag caactatgat cttgtacaat atgagacact tagtaaaact ttaagactaa 12000
tagtcagtga caagattaag tatgaagata tgtgctcagt agacctagcc atatcattaa 12060
gacaaaaaat gtggatgcat ttatcaggag gaagaatgat aaatggactt gaaactccag 12120
atcctttaga gttactgtct ggagtaataa taacaggatc tgaacattgt aggatatgtt 12180
attcaactga aggtgaaagc ccatatacat ggatgtattt accaggcaat cttaatatag 12240
gatcagctga gacaggaata gcatcattaa gggccctta ctttgatca gttacagatg 12300
agagatctga agcacaatta ggtatatca aaaatctaag caaaccagct aaggctgcta 12360
taagaatagc aatgatatat acttgggcat ttgggaatga cgaatatct tggttggaag 12420
catcacagat tgcacaaaca cgtgcaaaact ttacattgga tagcttaaag attttgacac 12480
cagtgacaac atcaacaaat ctatcacaca ggttaaaaga tactgtact cagatgaaat 12540
tttctagtac atcacttatt agagtaagca ggttcacac aatatctaata gataatatgt 12600
ctattaaaga agcaaatgaa actaaagata caaatcttat ttatcaacag gtaatgttaa 12660
caggattaag tgtatttgaa tatctattta ggttagagga gactacagga cataacccta 12720
tggtcatgca tctacatata gaggatggat gttgtataaa agagagttac aatgatgagc 12780
atatcaatcc ggagtctaca ttagagttaa tcaaataccc tgagagtaat gaatttatat 12840
atgataagga ccctttaaag gatatagatc tatcaaaatt aatggttata agagatcatt 12900
cttatacaat tgacatgaat tactgggatg acacagatat tgtacatgca atatcaatat 12960
gtactgcagt tacaatagca gatacaatgt cgcagctaga tcgggataat cttaaggagc 13020

```

```

tggttgtgat tgcaaatgat gatgatatta acagtctgat aactgaattt ctgaccctag 13080
atatactagt gtttctcaaa acatttggag ggttactcgt gaatcaattt gcatataccc 13140
tttatggatt gaaaatagaa ggaagggatc ccatthggga ttatataatg agaacattaa 13200
aagacacctc acattcagta cttaaagtat tatctaattg actatctcat ccaaaagtgt 13260
ttaagagatt ttgggattgt ggagttttga atcctattta tggtcctaact actgctagtc 13320
aagatcaagt taagcttgct ctctcgattt gcgagtactc cttggatcta tttatgagag 13380
aatggttgaa tggagcatca cttgagatct atatctgtga tagtgacatg gaaatagcaa 13440
atgacagaag acaagcattt ctctcaagac atcttgcctt tgtgtgttgt ttagcagaga 13500
tagcatcttt tggaccaaact ttattaaatc taacatatct agagagactt gatgaattaa 13560
aacaatactt agatctgaac atcaaagaag atcctactct taaatatgtg caagtatcag 13620
gactgttaat taaatcattc ccctcaactg ttacgtatgt aaggaaaact gcgattaagt 13680
atctgaggat tctgtgtatt aatccgcctg aaacgattga agattgggat cccatagaag 13740
atgagaatat cttagacaat attgttaaaa ctgtaaatga caattgcagt gataatcaaa 13800
agagaataaa aagtagttat ttctggggat tagctctaaa gaattatcaa gtcgtgaaaa 13860
taagatccat aacgagtgat tctgaagtta atgaagcttc gaatgttact acacatggaa 13920
tgacacttcc tcaggaggga agttatctat cacatcagct gaggttattt ggagtaaaac 13980
gtacaagttg tcttaaagct cttgaattat cacaatctt aatgaggga gttaaaaaag 14040
ataaagatag actcttttta ggagaaggag caggagctat gttagcatgt tatgatgcta 14100
cactcggctc tgcaataaat tattataatt ctggtttaaa tattacagat gtaattggtc 14160
aacgggaatt aaaaatcttc ccatacagaag tatcattagt aggtaaaaaa ctaggaaatg 14220
taacacagat tcttaatcgg gtgagggtgt tatttaattg gaatccaat tcaacatgga 14280
taggaaatat ggaatgtgag agtttaatat ggagtgaatt aaatgataag tcaattggtt 14340
tagtacattg tgacatggag ggagcgatag gcaaatcaga agaaactgtt ctacatgaac 14400
attatagtat tattaggatt acatatthaa tcggggatga tgatgttgtc ctagtatcaa 14460
aaattatacc aactattact ccgaattggt ctaaaatact ctatctatac aagttgtatt 14520
ggaaggatgt aagtgtagt tcccttaaaa catccaatcc tgcctcaaca gagctttatt 14580
taatttcaaa agatgcttac tgtactgtaa tggaaaccag taatcttggt ttatcaaaac 14640
ttaaaaggat atcatcaata gaagaaaata atctattaaa gtggataatc ttatcaaaaa 14700
ggaagaataa cgagtgggta cagcatgaaa tcaaagaagg agaaagggat tatgggataa 14760
tgaggccata tcatacagca ctgcaaattt ttggattcca aattaactta aatcacttag 14820
ctagagaatt tttatcaact cctgattthaa ccaacattaa taatataatt caaagtthta 14880
caagaacaat taaagatggt atgttcgaat gggtcfaat cactcatgac aataaaagac 14940
ataaattagg aggaagatat aatctattcc cgcttaaaaa taaggggaaa ttaagattat 15000
tatcacgaag attagtacta agctggatat cattatcctt atcaaccaga ttactgacgg 15060
gccgttttcc agatgaaaaa ttgaaaaata gggcacagac cggatatgta tcattggctg 15120
atattgattt agaatcctta aagttattat caagaaatat tgtcaaaaat taaaaaagac 15180
acataggatt aatatcatac tggtttttga ccaaagaggt caaaatacta atgaagctta 15240
taggaggagt caaactacta ggaattccta aacagtacaa agagttagag gatcgatcat 15300
ctcagggtta tgaatatgat aatgaatttg atattgatta atacataaaa acataaaata 15360
aaacacctat tcctcaccac ttcaattcca acaaaatgaa aagtaagaaa aacatgtaat 15420
atatatatac caaacagagt ttttctcttg ttttgtt 15456

```

<210> 23

<211> 15456

<212> DNA

<213> Bovine Parainfluenza Virus 3 (SF strain)

<400> 23

```

accaaacaag agaagagact tgcttgggaa tattaattca aataaaaatt aacttaggat 60
taaagaactt taccgaaagg taaggggaaa gaaatcctaa gactgtaatc atgttgagtc 120
tattcgacac attcagtgcg cgtaggcagg agaacataac aaaatcagct ggtggggctg 180
ttattcccgg gcaaaaaaac actgtgtcta ttttgtctct tggaccatca ataacagatg 240
acaatgacaa aatgacattg gctcttctct ttttgtctca ttctttagac aatgaaaagc 300
agcatgcgca aagagctgga tttttagttt ctctgttatc aatggcttat gccaacccag 360
aattatatatt aacatcaaat ggtagtaatg cagatgttaa atatgtcatc tacatgatag 420
agaaagaccc aggaagacag aaatatggtg ggtttgtcgt caagactaga gagatggttt 480
atgaaaagac aactgactgg atgtttggga gtgatcttga gtatgatcaa gacaatatgt 540
tgcaaaatgg tagaagcact tctacaatcg aggatcttgt tcatactttt ggatatccat 600
cgtgtcttgg agcccttata atccaggttt ggataatact tgtaaaggct ataaccagta 660
tatcaggatt gaggaaggga ttctttactc ggttagaagc atttcgacaa gatggaacag 720
ttaaatccag tctagtgttg agcgtgatg cagtagaaca aattggatca attatgaggt 780
cccaacagag cttgtaaca ctcatggttg aaacactgat aacaatgaac acaggcagga 840
atgacctgac aacaatagaa aagaatatac agattgtagg aaactacatc agagatgcag 900
gtcttgcttc attttcaac acaatcagat atggcattga gactagaatg gcagctctaa 960
ctctgtctac ccttagaccg gacatcaaca gactcaaggc actgatagag ctatatctat 1020
caaagggggc acgtgctcct tttatatgca ttttgagaga tcctgtgcat ggtgagtttg 1080
caccaggcaa ctatcctgcc ctctggagtt atgcgatggg ttagcagtt gtacaaaaca 1140
aggccatgca acagtatgta acaggaaggt cctatctgga tattgaaatg ttccaactgg 1200
gtcaagcagt ggcacgtgac gccgagtcgc agatgagttc aatattagag gatgaactgg 1260
gggtcacaca agaagccaag caaagcttga agaaacacat gaagaacatc agcagttcag 1320
atacaacctt ctataagcct acagggggat cagccataga aatggcaata gatgaggaag 1380
cagagcagcc cgaatccaga ggagaccaag accaaggaga tgaacctcgg tcatccatag 1440
ttccttatgc atgggcagac gaaaccggga atgacaacca aactgaatca accacagaaa 1500
ttgacagcat caaaactgaa caaagaaaca tcagagacag gctgaacaaa agactcaacg 1560
agaaaaggaa acagagtaac ccgggatcaa ctgacatcac aaacaacaca aatcaaactg 1620
aaatagatga tttattcagt gcattcggaa gcaactagtc acaaagagat gaccaccatc 1680
atcagcaaca agtaagaaaa acttaggatt aatggaaatt atccaatccg gagacggaag 1740
gacaaatcca gaatccaacc acaactcaat caaccaaaaga ttcatggaag acaatgttca 1800
aaacaatcaa atcatggatt cttgggaaga gggatcagga gataaatcat ctgacatctc 1860
atcggccctc gacatcattg aattcatact caacaccgac tccaagaga acacggcaga 1920
cagcaatgaa atcaacacag gagccacaag acttagcacg acaatctacc aacttgagtc 1980
caaaacaaca gaaacaagca aggaaaatag tggaccagct aacaaaaatc gacagtttg 2040
ggcatcacac gaacgtgcc aagagacaaa agatagaaat gtaaatcaga agactgtaca 2100
gggaggatat aggagaggaa gcagcccaga tagtagaact gagactatgg tcaactgagg 2160
aatctccaga agcagcccag atcctaacaa tggaaaccaa atccaggaag atattgatta 2220
caatgaagtt ggagagatgg ataaggactc tactaagagg gaaatgcgac aatttaaaga 2280
tgttccagtc aaggtatcag gaagtgatgc cattcctcca acaaaacaag atggagacgg 2340
tgatgatgga agaggcctgg aatctatcag tacatctgat tcaggatata ccagtatagt 2400
gactgccgca aactagatg acgaagaaga actccttatg aagaacaaca ggccaagaaa 2460
gtatcaatca acaccccaga acagtgacaa ggaattaaa aaagggagtg gaaggccaaa 2520
agacacagac aaacaatcac caatattgga ctacgaactc aactccaaag gatcgaagaa 2580
gagccagaaa atcctcaaag ccagcacgaa tacaggagaa ccaacaagat cacagagtgg 2640
atcccagggg aagagaatca catcctggaa catcctcaac agcgagagcg gcaatcgagc 2700
agaatcaaca aaccaaacc atcagacatc aatctcggga cagaaccaca caatgggacc 2760
aagcagaaca acctcagaac caaggaccaa gacacaaaag acggatggaa aggaaagaga 2820

```

```

ggacacagaa gagagcactc gatttacaga aagggcgatt acattattac agaatcttgg 2880
tgtaatccaa tctgcagcaa aattagacct ataccaagac aagagagttg tgtgtgtggc 2940
gaatgtccta aacaatgcag atactgcac aaagatagac ttcctagcag gtttgatgat 3000
aggagtgtca atggatcatg atgtcaaatt aaatcagatt cagaacgaga tattaagttt 3060
aaaaactgat cttagaaga tggatgaatc acatagaaga ctaattgaga atcaaaaaga 3120
acaattatca ctgatcacat cattaatctc aaatcttaaa atcatgacag agagaggagg 3180
gaagaaggac caaccagaac ctagcgggag gacatccatg atcaagacaa aggcaaaaga 3240
agagagaata aagaaagtca ggtttgacct tcttatggaa acacagggca tcgagaaaaa 3300
catccctgac ctctacagat caatagagaa aacaccagaa aacgacacac agatcaaadc 3360
agaaataaac agattgaatg atgaatccaa tgccactaga ttagtaccta gaagaataag 3420
cagtacaatg agatcactaa taataatcat caacaacagc aatttatcat caaaagcaaa 3480
gcaatcatac atcaacgaac tcaagctctg caagagtgat gaggaagtgt ctgagttgat 3540
ggacatgttc aatgaggatg tcagctccca gtaaacggcc aaccaagggt caacaccaag 3600
aaaaccaaca gcacaaaaca gccataaga gaccatccca acacaccgaa ccaatcaaca 3660
catacaaaag atcttttagat catagatgac taagaaaaac ttaggatgaa aggactgatc 3720
aatcctccaa aacaatgagc atcaccagct ccacaatcta cacattccca gaatcctctt 3780
tctccgagaa tggcaacata gagccgttac cactcaaggt caatgaacag agaaaggcca 3840
tacctcatat tagggttgtc aagataggag atccgcccac acatggatcc agatatctgg 3900
atgtcttttt actgggcttc tttgaaatgg aaaggtcaaa agacaggtat gggagcataa 3960
gtgatctaga tgatgatcca agttacaagg tttgtggctc tggatcattg ccacttgggt 4020
tggttagata cactggaat gatcaggaac tcctacaggc tgcaaccaag ctcgatatag 4080
aagtaagaag aactgtaaag gctacggaga tgatagtta cactgtgcaa aacatcaaac 4140
ctgaactata tccatggtcc agtagattaa gaaaagggat gttatttgac gctaacaagg 4200
ttgcacttgc tcctcaatgt cttccactag atagagggat aaaattcagg gtgatatttg 4260
tgaactgcac agcaattgga tcaataactc tattcaaaat ccccaagtcc atggcattgt 4320
tatcattgcc taatacaata tcaataaatc tacaagtaca tatcaaaaca ggaattcaga 4380
cagattccaa aggagttagt cagattctag atgaaaaagg tgaaaaatca ctaaatattca 4440
tggttcattc cggttgatc aaaaggaaga tgggtagaat gtactcagtt gaattattga 4500
agcagaagat tgagaagatg agattattat tctcattggg attagttgga gggatcagct 4560
tccacgtcaa cgcaactggc tctatatcaa agacattagc aagtcaatta gcatttaaaa 4620
gagaaatctg ctatccccta atggatctga atccacactt aaatttagtt atatgggcat 4680
catcagttga aattacaaga gtatagtcaa tctccagcc ttcattacct gggaattca 4740
gatactacct aaacatcata gcaaaagggg tcgggaaaat cagacagtaa aaccaacaac 4800
cctgacatcc aacactgcaa atcaggctac ccacaggaga aaaatcaaaa acttaggatc 4860
aaagggatca ccacaaaccc cggaacacag ccaaaccaac caacacacaa atcacagaca 4920
aaaaggaaaa ggcatgcaa agaccgagaa caagcagaac gcacacaacc aagcagagga 4980
aagccaaagc ccgcatcca caaacacacc aacaatccta caaacaagca ccaaaataga 5040
ggtcaaaaga caaagagcat cagatatgac catcacaacc ataatcatag ccatactact 5100
aatacccta tcattctgtc aaatagacat aacaaaactg caacgtgtag gtgtattagt 5160
caacaatccc aaaggcatga aaatttcaca aaattttgaa acgagatacc tgatattaag 5220
tctgataccc aaaatagaga attcacactc atgtggggat caacagataa accaatacaa 5280
gaagttattg gatagattga taattcctct atatgatgga ttaaaattac aaaaagatgt 5340
aatagtagta agtcatgaaa ccataataa tactaatctt aggacaaaac gattcttttg 5400
agagataatt gggacaattg cgatagggat agccacctca gcgcaaatca ccgcagcagt 5460
cgctcttgtc gaagctaaac aggaagggtc agacatagaa aaactcaaag aagctataag 5520
agacacaaac aaggcagtac aatcgattca aagttctgta ggtaacctaa ttgttgacgt 5580
taaatacagtt caagactatg tcaacaatga aattgtacct tcaatcacia gattaggctg 5640
tgaagcagca gggttacaat tgggaattgc actgacacaa cattactcag aattaacaaa 5700

```

tataatttgggt gataatatag gaacactgaa agaaaaaggg ataaaattac aggggatagc 5760
atcggttatat catacaaaca taacagaaat atttactact tcaacagttg accaatatga 5820
tatttatgac ctattattca ctgaatcaat caagatgaga gtgatagatg ttgatttgag 5880
tgattactca attactcttc aagtttagact tcctttatta actaaactat caaatactca 5940
gatttataaa gtagattcta tatcatacaa catccagggc aaagagtggg atattcctct 6000
tcccaatcac atcatgacaa aaggggcttt tctaggtggg gctgatatta aagaatgcat 6060
agaggcattc agcagttata tatgtccttc tgatccaggt tatatatata atcacgagat 6120
agagaattgt ttatcagggg acataacaca gtgtcctaag actgttggtta catcagatgt 6180
gggtaccacga tacgcgtttg tgaatgggtg attaattgca aactgcataa caactacatg 6240
tacatgcaat ggaattgaca atagaattaa tcaatcacct gatcaaggaa ttaagatcat 6300
aacacataaa gaatgccagg taatagggtat aaacggaatg ttattcaata ctaatagaga 6360
agggacatta gcaacttata catttgatga cattatatta aataactctg ttgcacttaa 6420
tccaattgat atatctatgg aacttaacaa ggcaaaacta gaattagaag aatcgaagg 6480
atggataaag aaatcaaatac aaaagttaga ttccgttgga agttggtatc aatctagtgc 6540
aacaatcacc ataatacatg tgatgataat aattctatctt ataatacaata taacaattat 6600
tgtagtcata atcaaattct atagaattaa ggggggaaat caaacgaca aaaacagtga 6660
gccgtatata ctgacaaata gacaataaga ctatacacga tcaaatatag aaagtacaaa 6720
aaacttagga acaaagttgt tcaacacagc agcagcgaac agacccaaag gcagcgcaga 6780
ggcgacaccg aacccaaaaa tggaattattg gaaacacaca aacagcaca aaaacaccaa 6840
caatgaaacc gaaacaacca gaggcaaaca cagtagcaag gttacaaata tcataatgta 6900
caccttctgg acaataacat caacaatatt attagtcatt ttataatga tattgacaaa 6960
cttaattcaa gagaacaatc ataataaatt aatgttcgag gaaataagaa aagaattcgc 7020
ggcaatagac accaagattc agaggacctc ggatgacatt ggaacctcaa tacagtcagg 7080
aataaataca agacttctca caattcagag tcatgttcaa aactatatcc cactatcact 7140
aacacaacaa atgtcagatc tcagaaaatt tatcaatgat ctaacaaata aaagagaaca 7200
tcaagaagtg ccaatacaga gaatgactca tgatagagggt atagaacccc taaatccaga 7260
caagttctgg aggtgtacat ctggttaacc atctctaaca agtagtccta agataagggt 7320
aataccaggg ccaggtttat tagcaacatc tactacagta aatggctgta ttagaatccc 7380
atcggttagca atcaatcatt taatctacgc ttacacctct aatcttatca cccagggctg 7440
tcaaaatata gggaaatctt accaagtact acaaatagggt ataattacta taaattcgga 7500
cctagtacct gattttaaatac ccagagtcac acatacattt aatattgatg ataataggaa 7560
atcttgctct ctggcactat tgaatacaga tgtttatcag ttatgctcaa caccaaaagt 7620
tgatgagaga tccgattatg catcaacagg tattgaggat attgtacttg acattgtcac 7680
taataatgga ttaattataa caacaagggt tacaataaat aatataactt ttgataaacc 7740
gtatgcagca ttgtatccat cagtaggacc aggaatctat tataagggtta aagttatatt 7800
tctcgatat ggaggtctag agcatgaaga aaacggagac gtaatatgta atacaactgg 7860
ttgtcctggc aaaacacaga gagactgtaa tcaggcttct tatagcccat ggttctcaaa 7920
taggagaatg gtaaaactcta ttattgttgt tgataaaggc atagatgcaa cttttagctt 7980
gaggggtgtg actattccaa tgagccaaaa ttattgggga tcagaaggaa gattactttt 8040
attaggtgac agaataatac tatatactag atccacaagt tggcacagta aattacagtt 8100
aggggtaatt gatatttctg attataataa tataagaata aattggactt ggcataatgt 8160
actatcacgg ccaggaaatg atgaatgtcc atgggggtcat tcatgccag acggatgtat 8220
aacaggagtt tacactgatg catatccgct aaacccatcg gggagtgttg tatcatcagt 8280
aattcttgac tcacaaaagt ctagagaaaa cccaatcatt acctactcaa cagctacaaa 8340
tagaataaat gaattagcta tatataacag aacacttcca gctgcatata caacaacaaa 8400
ttgtatcaca cattatgata aagggatttg tttcatata gtagaaataa atcacagaag 8460
tttgaatacg tttcaaccta tgttattcaa aacagaagtt ccaaaaaact gcagctaaat 8520
tgatcatcgc atatcggatg ccagatgaca ttaaaagaga ccaccagaca gacaacacag 8580

```

gagatgatgc aagatatataa ggaataataa aaaacttagg agaaaagtgt gcaagaaaaa 8640
tggacactga atcccacagc ggcacaacat ctgacattct gtaccctgaa tgtcacctca 8700
attctcctat agttaaagga aaaatagcac aactgcatac aataatgagt ttgcccacac 8760
cctacgatat ggatgatgat tcaatactga ttattactag acaaaaaatc aaactcaata 8820
aattagataa aagacaacg tcaattagga aattaagatc agtcttaatg gaaagagtaa 8880
atgatcttgg taaatacacc tttatcagat atccagaaat gtctagttaa atgttccaat 8940
tatgtatacc cggaattaat aataaaataa atgaattgct aagtaaagca agtaaaacat 9000
ataatcaa at gactgatgga ttaagagatc tatgggttac tgtactatcg aagtttagcat 9060
cgaaaaatga tggaaagta at tatgatatac atgaagatat tagcaatata tcaa atgttc 9120
acatgactta ccaatcagac aaatgggtata atccattcaa gacatgggtt actattaagt 9180
atgacatgag gagattacaa aaagccaaaa atgagattac attcaatagg cataaagatt 9240
ataatctatt agaagaccaa aagaatatat tgctgataca tccagaactc gtcttaatat 9300
tagataaaca aaattacaat gggatatata tgactcctga attgggtacta atgtattgtg 9360
atgtagttga agggaggtgg aatataagtt catgtgcaaa attggatcct aaattacaat 9420
caatgtatta taaaggtaac aatttatggg aaataataga tggactattc ctgaccttag 9480
gagaaagAAC atttgacata atatcactat tagaacgct tgcattatcg ctcatcaca 9540
ctcatgacc ggttaaacag ctgaggggg ctttttttaa tcacgtgtta tcagaaatgg 9600
aatcaatatt cgcagctgag tgtacaacag aggaaatacc taatgtggat tatatagata 9660
aaattttaga tgtattcaaa gaatcaaca tagatgaa atgcagaaatt ttctctttct 9720
tccgaacttt tggacaccct ccattagagg cgagtatagc agcagagaaa gttagaaagt 9780
atatgtacac tgagaaatgt ttgaaatttg atactatcaa taaatgtcat gctatttttt 9840
gtacaataat tataaatgga tatagagaaa gacatgggtg tcaatggcct ccagttacat 9900
tacctattca tgcacatgaa tttatcataa atgcgtacgg atcaaattct gccatcatcat 9960
atgaaaatgc tgtagattat tataagagct tcataggaat aaaatttgac aagtttatag 10020
agcctcaatt ggatgaagac ttaactat atatgaaaga taaagcatta tccccaaaga 10080
aatctaactg ggacacagtc tatccagctt caaacctgtt ataccgcaat aatgtgtctc 10140
atgattcacg aagattgggt gaagtattta tagcagatag taaatttgat cccaccaag 10200
tattagatta cgtagaatca ggatattggc tagatgatcc tgaattta atctcatata 10260
gtttaaaaga gaaagaaata aaacaagaag gtgactttt tgcaaaaatg acatacaaga 10320
tgagagctac acaagtatta tcagaaacat tattggcgaa taatatagg aaatccttcc 10380
aagagaatgg gatgggttaa ggagaaattg aattactcaa gagactgaca acaatatcta 10440
tgtctggggt tccgcggtat aatgaggtat acaataattc aaaaagtcac acagaggaac 10500
ttcaagctta taatgaatt agcagttcca atttatcttc taatcagaag tcaagaagt 10560
ttgaatttaa atcaacagat atatacaatg atggatacga aaccgtaagc tgcttcttaa 10620
cgacagatct taaaaaatat tgtttaaatt ggaggtatga atcaacagct ttattcggtg 10680
atacttgtaa tcagatattt gggttaaagg aattatttaa ttggctgcac cctcgcttg 10740
aaaagagtac aatatatgtt ggagatcctt attgcccgcc atcagatatt gaacatttac 10800
cacttgatga ccactctgat tcaggatttt atgttcataa tcctaaagga ggaatagaag 10860
ggttttgcca aaagttagtg aactcatat ctatcagtc catacattta gcagctgtca 10920
aaatcggtgt aagagttact gcaatgggtc aaggggataa tcaagccata gctgttacca 10980
ccagagtacc taataattat gattataagg ttaagaaaga gattgtttat aaagatgtgg 11040
taagattttt tgattctttg agagaggtta tggatgatct gggatcatg ctcaaactaa 11100
atgaaactat aataagtagt aaaatgttta tatatagcaa aaggatatac tatgacggaa 11160
gaatccttcc tcaggcggtta aaagcattgt ctagatgtgt tttttggtct gaaacaatca 11220
tagatgagac aagatcagca tcctcaaactc tggcgacatc gtttgcaaag gccattgaga 11280
atggctactc acctgtattg ggatattgat gctcaatctt caaaaatc caacagttgt 11340
atatagcact tggaatgaat ataaatccaa ctataaccca aaatatttaa gatcaatatt 11400
tcaggaatat tcattggatg caatatgcat ctctaattccc tgctagtgtc ggaggattta 11460

```

```

atttatatggc catgtcaagg tgttttgtca gaaacattgg agatcctaca gtcgctgcat 11520
tagctgatat taaaagattt ataaaagcaa atttgttaga tcgaggtgtc ctttacagaa 11580
ttatgaatca ggaaccaggc gagtcctcct ttttagactg ggcttcagac ccctattcat 11640
gtaacttacc acaatctcaa aatataacca ccatgataaa gaatataact gcaagaaatg 11700
tactacagga ctcaccaaac ccattactat ctggattatt tacaagtaca atgatagaag 11760
aggatgagga attagctgag ttcctaattg acaggagaat aattctccca agggttgctc 11820
atgacatttt agataattct cttactggaa ttaggaatgc tatagctggg atgttgata 11880
caacaaaatc actaattcga gtagggataa acagaggagg attaacctat aacttattaa 11940
gaaagataag caactatgat cttgtacaat atgagacact tagtaaaact ttaagactaa 12000
tagtcagtga caagattaag tatgaagata tgtgctcagt agacctagcc atatcattaa 12060
gacaaaaaat gtgatgcat ttatcaggag gaagaatgat aaatggactt gaaactccag 12120
atcctttaga gttactgtct ggagtaataa taacaggatc tgagcattgt aggatatgtt 12180
attcaactga aggtgaaagc ccatatacat ggatgtattt accaggcaat cttaatatag 12240
gatcagctga aacaggaata gcatcattaa gggtcctta ctttgatca gttacggatg 12300
agagatctga agcacaattg gggatatca aaaatctaag caaacagct aaggctgcta 12360
taagaatagc aatgatatat acttgggcat ttgggaatga cgaaatatct tggatggaag 12420
catcacagat tgcacaaaca cgtgcgaact ttacattaga tagcttaaag attttgacac 12480
cagtgcacac atcaacaaat ctatcacata ggtaaaaaga tactgctact cagatgaaat 12540
tttctagtac atcacttatt agagtaagca ggttcacac aatatctaata gataatatgt 12600
ctattaaaga ggcaaatgaa actaaagata caaatcttat ttatcaacag gtaatgttaa 12660
cagggttaag tgtatttgaa tatctattta ggtagagga gtagacagga cataacccta 12720
tggtcatgca tctacatata gaggatggat gttgtatcaa agagagttac aatgatgagc 12780
atatcaatcc ggagtctaca ttagagttaa ttaaataccc tgagagtaat gaatttatat 12840
atgataagga ccctttaaag gatatagatc tatcaaaatt aatgggtata agagatcatt 12900
cttatacaat tgacatgaat tactgggacg acacagatat tgtacatgca atatcaatat 12960
gtactgcagt tacaatagca gatacaatgt cgcagctaga tcgggataat cttaggagc 13020
tggttgtaat tgcaaatgat gatgatatta acagtctgat aactgaattt ctgacctag 13080
atatactagt gtttctcaaa acatttggag gggtactcgt gaatcaattt gcataatacc 13140
tttatggatt gaaaatagaa ggaagggatc ccatttggga ttatataatg agaacattaa 13200
aagacacctc acattcagta cttaaagtat tatctaagtc actatctcat ccaaaagtgt 13260
ttaagagatt ttgggattgt ggagtttga atcctattta tggctctaact actgctagtc 13320
aggaccaagt taagcttgct ctctcaattt gcgagtactc cttggatcta tttatgagag 13380
aatggctgaa tggagcatca cttgagatct atatctgtga tagtgacatg gaaatagcaa 13440
atgatagaag acaagcattt ctctcaagac accttgcctt tgtgtgttgt ttacagaga 13500
tagcatcttt tggaccaaatt ttattaaatc taacatatct agagagactt gacgaattaa 13560
aacaatactt ggatctgaac atcaaagaag atcctactct taaatatgtg caagtatcag 13620
gactgttaat taaatcattc ccctcaactg ttacgtatgt gaggaaaact gcgattaagt 13680
atctgaggat tcgtggcatt aatccgctg aaacgattga agattgggat cccatagaag 13740
atgagaatat cttagacaat attgttaaaa ctgtaaatga caattgcagt gataatcaaa 13800
agagaaataa aagtagttat ttctggggat tagctctaaa gaattatcaa gtcgtaaaaa 13860
taagatccat aacgagtgat tctgaagtta atgaagcttc gaatgttact acacatggaa 13920
tgacacttcc tcaggaggga agttatctat cacatcagct gaggttattt ggagtaaaca 13980
gtacaagttg tctgaaagct cttgaattgt cacaaatttt aatgagggaa gttaaaaaag 14040
ataaagatag actcttttta ggagaaggag caggagctat gttagcatgt tatgatgcta 14100
cactcgggtc tgcaataaat tattacaatt ctggtttaa tattacagat gtaattgggtc 14160
aacgggaatt aaaaatcttc ccatcagaag tatcattagt aggtaaaaaa ctaggaaatg 14220
taacacagat tcttaatcgg gtgaggggtg tatttaattg gaatcccaat tcaacatgga 14280
taggaaatat ggaatgtgag agtttaatat ggagtgaatt aaatgataag tcaattgggt 14340

```

WO 01/04320

```

tagtacattg tgacatggag ggagcgatag gcaaatcaga agaaactggt ttacatgaac 14400
attatagtat tattaggatt acatatttaa tcgggggatga tgatgttgtc ctagtatcaa 14460
aaattatacc aactattact ccgaattggt ctaaaatact ctatctatac aagttgtatt 14520
ggaaggatgt aagtgtagt tcccttaaaa catccaatcc tgcctcaaca gagctttatt 14580
taatttcaaa ggatgcttac tgtactgtaa tggaacccag taatcttggt ttatcaaaac 14640
ttaaaggat atcatcagta gaagaaaata atctattaaa atggataatc ttatcaaaaa 14700
ggaagaacaa cgaatgggta cagcatgaaa tcaaagaagg agaaagggt tatgggataa 14760
tgaggccata tcatacagca ctgcaaattt ttggattcca aattaactta aatcacttag 14820
ctaaagaatt tttatcaact cctgatttaa ccaacattaa taatataatt caaagtttta 14880
caagaacaat taaagatgtt atgttcgaat gggccaatat cactcatgac aataaaagac 14940
ataaattagg aggaagatat aatctattcc cgcttaaaaa taagggaag ttaagattac 15000
tatcacgaag attagtacta agctggatat cattatctt atcaaccaga ttactgacag 15060
gccgtttccc agatgaaaaa tttgaaaata gggcacagac cggatatgta tcattggctg 15120
atactgattt agaactttta aagttattat caagaaatat tgtcaaaagt tacaagaac 15180
acataggatt aatatcatac tggtttttaa ccaaagggt caaaatacta atgaaactta 15240
tagggggagt caaactacta ggaattccca aacagtacaa agagtttag gatcgatcat 15300
ttcagggtta tgaatatgat aatgaatttg atattgatta atacataaaa acaaaaaata 15360
aaacacctaa tcctctccca ttcaattcca acaaaatgaa aagtaagaaa aacatataat 15420
atacatatac caaacagagt tttctcttg tttggt 15456

```

<210> 24

<211> 12

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sequence
flanking site for introduction of Bsi W1 site for
rHPiV3 JS

<400> 24

tagacaaaag gg

12

<210> 25

<211> 12

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sequence
flanking site for introduction of Bsi W1 site for
rBPIV3 Ka

<400> 25

aagatataaa ga

12

<210> 26
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking site for introduction of Bsi W1 site for
rHPiV3 s

12

<400> 26
tagacgtacg gg

<210> 27
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking site for introduction of Bsi W1 site for
rBPIV3 s

12

<400> 27
aagacgtacg ga

<210> 28
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking Sgr A1 site in rHPiV3 sFBHNB

12

<400> 28
cgcaccggtg ca

<210> 29
<211> 12
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sequence
flanking Bsi W1 site in rHP1V3 sFBHNB

<400> 29
aagacgtacg gg

12

<210> 30
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking Sgr A1 site in rBP1V3 sFHNNH

<400> 30
tccaccggtg ta

12

<210> 31
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence
flanking Bsi W1 site in rBP1V3 of sFHNNH

<400> 31
tagacgtacg ga

12